normal proportions and in normal position. The difficulty of diagnosing anencephaly and congenital goitre has already been referred to.

CASES OF INCOMPLETE DILATATION OF THE CERVIX.

In discussing occipito-posterior positions I referred to the slow dilatation of the cervix in such cases. In my series of 68 cases of contracted pelvis in which forceps had been unsuccessfully applied, the cervix was still incompletely dilated in 33 of the cases when the patient was admitted to hospital. In two of this group the dilatation was of such slight degree that it was difficult to imagine how forceps could ever have been applied to the foetal head. In all these cases there was a definite cause for the delayed labour and the slow dilatation of the cervix, but in 151 of all the cases reviewed by Dr. Miller there was no evidence of any abnormality; yet forceps had been applied before the cervix was fully dilated. In the 34 cases which I contributed to this group one mother died from rupture of the uterus, five had a morbid puerperium, but the remainder had an uneventful recovery. There are probably many of these women who will still require treatment in gynaecological wards, because traction through an undilated cervix causes not only tearing of the cervix, but also forced descent of the cervix with subsequent uterine prolapse. However, it is when we consider the results to the child that the picture is blackest. Of the 34 children 15 were stillborn, and several of the survivors showed extensive bruising at birth.

I think we must agree that the application of forceps before complete dilatation of the cervix is a most dangerous procedure; yet we see how often it appears to have been practised. Efficient ante-natal care should identify the cases where labour is likely to be protracted and dilatation slow, but it is only effective teaching which can prevent the too early application of forceps in normal cases. From my experience as a teacher I know that on questioning students there is hardly one who does not remember to state that the cervix must be fully dilated before forceps are applied.

The point on which I think teachers and textbooks do not lay sufficient emphasis is that manual dilatation of the cervix is seldom justified, and, even when it is attempted, is very often incomplete. I have even been informed by a young graduate, negotiating for the admission of his "failed forceps" case to hospital, that he had first used gentle traction with the forceps to secure dilatation of the cervix. The patient eventually delivered herself spontaneously.

CONCLUSION.

In estimating how far the cases of "failed forceps" could have been prevented by efficient ante-natal care, I have tried to show that almost all of the real abnormalities could have been identified during the course of pregnancy, and particularly in the last month. There is only a very small group of foetal abnormalities of the developmental type which cannot be detected before the onset of labour. Ante-natal supervision, however, cannot prevent the too early application of forceps in normal cases.

When abnormalities have been detected they may either be corrected before the onset of labour, as in occipito-posterior, breech, and transverse presentations, or sent to a hospital or similar institution where appropriate treatment can be carried out in the most favourable circumstances at the correct stage in labour.

It may be asked whether the ante-natal supervision to which I have referred can be efficiently carried out by the family physician. In many cases it can, but there must always be a considerable proportion of cases in which accurate diagnosis of pelvic disproportion or abnormal presentation is difficult, even for the hospital physician with special training. He may require the help of x rays to establish his diagnosis. It would appear, therefore, that an efficient obstetric service would require well-equipped consultative ante-natal centres to which the cases presenting difficulties to the family physician can be referred. Again, work of this kind is far too responsible and difficult for midwives, and I believe that a midwife

should not be authorized to undertake the care of any pregnant woman without provision being made for the patient to be examined carefully by a medical practitioner before labour is due.

REFERENCES.

- ¹ *Zeit. f. Geburtsh. und Gynäkol.*, 1927, xci, 3, p. 504.
- ² *Proceedings Edinburgh Obstetrical Society*, 1926, p. 133.

UNSUCCESSFUL FORCEPS CASES.*

BY

WILLIAM FLETCHER SHAW, M.D., CH.B.,

Professor of Clinical Obstetrics and Gynaecology, University of Manchester; Honorary Gynaecological Surgeon, Manchester Royal Infirmary.

In the last few years more and more attention has been focused upon maternity practice; meetings of various bodies outside the profession have discussed the subject, often with great ignorance of the facts, and necessarily, from the composition of these meetings, the blame for the present high mortality in childbirth has been thrown upon the members of our profession.

In the great majority of instances these charges are an injustice, and a slight knowledge of the facts would prevent them from being made; but in some directions obstetric practice leaves much to be desired, and it behoves us, as a profession, to take stock of our methods and see if it is not possible to raise the standard. It is for this reason only that we have brought forward this subject for discussion before a meeting composed largely of general practitioners. We have stated our views as members of the staffs of maternity hospitals, and we hope that general practitioners will fully state their views, and that the discussion will bear fruit. In similar discussions in the past it has often been stated that the consultant speaks only from his experience of hospital work, and compares his own results in ideal circumstances with those of men who have to conduct their cases in circumstances far from ideal. The fact is overlooked that few consulting obstetricians reach the staff of their hospital without an extensive experience of maternity work in the very poor quarters of their cities, and therefore are fully alive to all these disadvantages. One advantage, and one only, had we over the general practitioner—we were doing obstetrical work only, and had no anxiety about other cases being neglected when we were engaged for a long period over one maternity case. If this is a factor in bad midwifery, and I feel sure it is one of the most important, it is better to face the facts and see if a remedy cannot be found.

The aspect of maternity work we are considering to-day is one of the greatest blots on our obstetrical escutcheon. The figures given by Dr. Miller occurred in the practice of three large maternity hospitals, but these hospitals were chosen at random, and there is no doubt that equally convincing figures could be produced from every other maternity hospital in the country. The facts are there and are incontrovertible. Dr. Miller has shown the types of cases which result in forceps failure, and has analysed the causes and discussed the diagnosis and treatment. He has also detailed the appalling results to mother and child, results which could have been avoided with a little more care and knowledge.

One of the most striking figures in this series is the large number of craniotomies performed. In the last twenty years the number of Caesarean sections performed in each obstetrical unit has enormously increased, and the first thought on seeing these figures is that many of these children could have been saved by this operation. In considering this alternative treatment we must remember in the first place that the majority of the children were dead or badly damaged when the patient was admitted to hospital, but a small number were alive and were deliberately sacrificed to save the mother.

Caesarean section performed upon a "clean" patient is now a safe operation with a very low mortality, but Caesarean section performed upon a patient after attempts

* Read in a discussion in the Section of Obstetrics and Gynaecology of the Annual Meeting of the British Medical Association, Cardiff, 1928.

at forceps delivery is a very different matter, and is followed by a dreadfully high maternal mortality. Eardley Holland¹ in 1921, in a collection of 3,374 cases from the large maternity hospitals of the British Isles, showed that the mortality in cases in which the operation was performed before the onset of labour was only 1.6 per cent., whereas in the cases in which it was performed after attempts at forceps delivery it had risen to 26 per cent. With the introduction of Caesarean section through the lower uterine segment and delivery of the placenta and cord through the cervix and vagina after closure of the uterine incision, as suggested by Munro Kerr, there is probably less risk of infection, and he has published a number of such cases safely delivered. It is too early to write dogmatically upon this, but this method does seem to give much greater safety from infection in "suspect" cases, and there is a growing appreciation of its value.

Another factor which renders these cases more liable to sepsis is that many of the women are admitted with lacerations of the cervix and vagina produced by attempts to deliver them with forceps; in many others the patient is so shocked by the previous manipulation that an abdominal section is quite out of the question.

In the majority of cases admitted after attempted delivery Caesarean section is out of the question, owing to death of or damage to the foetus, definite infection of the mother, shock, or severe lacerations of the cervix or vagina; and in the small remaining group of suspect cases where none of these definite contraindications appear there is a justifiable difference of opinion. Some operators, and a growing number, will operate through a lower uterine incision with delivery of placenta and membranes by the vagina, whilst others consider the risk of sepsis too great even by this method.

When the dangers of Caesarean section were overcome and the operation leaped into popularity it was expected that the number of craniotomies would reach a vanishing point, but this is far from being the case. In 1906 in St. Mary's Hospital, Manchester, the number of Caesarean sections was only four, whilst there were fifteen craniotomies; in 1925 the number of Caesarean sections had jumped to 132, but the craniotomies, instead of diminishing, had increased to thirty. Craniotomy, nowadays, is seldom performed except upon cases sent in after intervention, and these figures are a strong indictment upon one class of obstetrical practice.

Pubiotomy is another operation which may be employed in this type of case, and a few operators do employ it, but it has many disadvantages and has never become popular. The only type of case in which it is useful is one in which the pelvis has a degree of contraction which just prevents extraction of the head, but which will just allow the head to pass after it has been widened by the comparatively small amount obtained by the operation. If this amount is miscalculated there is great danger of severe laceration of the soft tissues.

Dr. Hendry has pointed out how most of these cases could have been avoided by detailed ante-natal care.

What we have now to consider are the main causes underlying this state of affairs and the best method of removing these causes. There is little doubt that apathy is the main reason—the general failure of both the profession and the laity to realize the risks of labour and that special knowledge is necessary for the skilful treatment of abnormal cases—and the crowding of obstetrical teaching into any part of the curriculum not already occupied by medicine and surgery. These faults are due as much to the apathy and ignorance of the lay public as to that of our profession, and it behoves us, in seeking a remedy, to stress the importance of the education of the laity.

In the minds of the laity labour is a natural function—if it ends successfully it is no credit to the doctor or the nurse; if anything goes wrong then the attendants must be to blame. A practical outcome of this is the very small remuneration paid for attendance upon a maternity case. A doctor may attend a case for twenty-four or forty-eight hours, may be up one or two whole nights, may completely dislocate the whole work of his practice for some days, during which time he is worn out with anxiety for this

patient and the others he is neglecting, and yet when all is over he receives in remuneration less than he would for removing tonsils or any minor operation, and in credit very little beyond what is due to his pleasant and sympathetic manner, as he has only been observing a normal function.

If anything abnormal arises he is expected to recognize it and treat it; the fact that he is a qualified medical man makes him, in the opinion of most of the laity, an expert obstetrician capable of tackling any obstetrical problem. If, in his modesty, he recognizes that the abnormality requires someone more experienced than himself, he feels this means a loss of prestige in the eyes of the patient and her friends and that he will be blamed for the extra expense incurred.

Until the general public realizes the responsibility entailed by every maternity case, no matter how straightforward, and the anxiety, dislocation of work, and broken rest which many cases involve, and until the general public realizes the skill and patience necessary for the conduct of the simplest labour—increased a hundredfold in cases of delay and difficulty—and is prepared to pay fees commensurate with the time and anxiety involved, so long will the temptation remain to hasten delivery by every available means.

This subject of fees applies to the nursing as well as to the medical profession, and it is most noticeable at examinations of the Central Midwives Board to find that all the better class of candidates are taking this diploma merely to help them to secure senior appointments and with no intention of practising midwifery.

The fault, however, does not lie entirely with the lay public; we as a profession must take a considerable share of the blame for the state of affairs which can allow such a series of cases to be brought forward. In the past midwifery has not been taken sufficiently seriously. Until 1888 it was not compulsory for a student to attend any cases of labour before qualifying, and even now it is necessary for him to spend only one month in a maternity hospital, whereas he must spend six months dressing in a surgical ward. After that date he had to be signed up for twelve and later for twenty cases, but it was possible to attend these without seeing any abnormality, and so he entered practice without any realization, other than theory, of the serious difficulties to be encountered, and, never having seen obstetrical operations performed, could not realize that a trained obstetrician would perform these operations more skilfully and with much less risk to mother and child than he could do in his own bungling way.

When we remember the haste with which we completed our twenty cases, the surroundings in which many had to reside while attending these, the fact that many students never saw a member of the honorary staff of the hospital, or even an abnormality treated by a resident, can we wonder that midwifery was looked upon as a bore and every case of labour as a nuisance which must be terminated as quickly as possible, and especially that the student should soon consider himself, as did his neighbours, as capable of conducting a complicated case as any specialist? The month's residence in hospital which is now compulsory has improved this, and it must be impossible for any student to spend this time in hospital without seeing a considerable number of complications treated by the members of the medical staff; but this time should be extended.

It is the proud boast of many men in practice that they have attended a certain number of thousands of labour cases and carried out all obstetrical operations, and therefore know as much as, or more than, any young consultant, though they overlook the fact that the consultant will probably have more experience of difficult cases concentrated into one year of hospital life than the practitioner can acquire in a lifetime, and, moreover, learns to perform these operations under men already skilled in performing them.

It is a great pity that there is this feeling of antagonism to young consulting obstetricians; it does not apply to young surgeons or physicians or even to young gynaecologists, and it would be all to the advantage of a young

practitioner if he would call to his aid in midwifery difficulties a highly trained specialist who would show him how best to deal with the problem.

The fact that the young graduate is not a specialist in obstetrics and has not had experience in performing all obstetrical operations is often laid as a charge against the competence of his obstetric teachers. In the past there was perhaps some truth in this, though the chief fault lay in the regulations which allowed the student to attend a certain number of cases without being in a maternity hospital and without coming in contact with the teachers; but that is altered now, and during the month's residence there is every opportunity to hold classes, and although the irregularity of the admission of abnormal cases makes it impossible for the senior members of the staff to demonstrate the operative treatment in all cases, they can hold classes on the most important clinical subjects.

There has been a great revival in clinical teaching in the last few years, and every large teaching maternity hospital endeavours to make the best use of its opportunities, and in the month at their disposal give each series of students a sound clinical foundation for their future work; but it must be emphasized that there are limits beyond which they cannot go, and even with more time at their disposal it is impossible to make every student a specialist, if for no other reason than that the number of abnormal cases admitted even to the largest maternity hospital would not provide sufficient material. If a graduate wishes to have special experience in operative obstetrical work he must be prepared to spend some time as a house-surgeon in a maternity hospital.

In teaching clinical obstetrics special emphasis must be laid upon certain sections of the work, and if this is done the graduate of the future will be turned out capable of taking charge of normal cases, of recognizing abnormalities, and of treating the minor ones, and with a sound clinical foundation upon which to build with future experience. The following four points are of importance and require special emphasis.

1. Ante-natal Work.

More and more emphasis is now laid upon the importance of the constant supervision of the pregnant woman. If this is universally done the great majority of the cases of eclampsia will be eliminated, defects in the measurement of the pelvis will be detected and appropriate treatment undertaken, and defects in the position of the child recognized and rectified. If this had been universally carried out in these three cities one large group of cases—the contracted pelvis—would have been eliminated from our tables, and these women saved much mutilation and have acquired living children.

It is almost unbelievable how little attention is given to this in some quarters. One of the chief reasons is the modern tendency of women of the working class to engage a midwife, who calls in a doctor only in cases of difficulty. This is the wrong way round; every pregnant woman should be under the care of a qualified medical practitioner, and if no abnormality is found the actual labour could be attended by a midwife. Even with this defect it is difficult to imagine the mentality of a man who applied forceps to a patient with a history of four previous confinements, two of which ended in craniotomy and two in Caesarean section; and yet such a case occurs in the list from my hospital.

Even the health authorities in many centres hardly realize the special training required for the work, and appoint to the charge of their ante-natal clinics young newly qualified practitioners without previous experience in this work, and with little or no chance of acquiring it in the future, as they will not see any of these cases in labour and so learn from their own mistakes.

2. Antisepsis.

It is impossible to overstress the importance of this, and it is impossible to shut one's eyes to the fact that with many practitioners this is of a most sketchy variety. I cannot help thinking that the almost universal use of coal-tar derivatives has had a bad effect. Used in sufficient

strength to be really antiseptic they are hard upon the hands, and so the tendency is to use them in insufficient strength.

If thorough antisepsis had been carried out there is no reason why these unsuccessful forceps cases should show a higher morbidity than other forceps cases in the hospital, but the figures quoted by Dr. Miller show these to have a much higher incidence than those treated primarily at the hospital.

3. First Stage of Labour.

In a straightforward case this is undoubtedly the most trying to all concerned. The patient and her friends naturally desire its curtailment, and use what influence they can upon the doctor to terminate it. This, no doubt, accounts for the large number of cases in our list in which forceps were applied before the cervix was dilated, the majority of which delivered themselves after time had been given in hospital for the cervix to dilate. These cases admitted after forceps failure possessed stout cervices which would not tear, but for each of these there must have been a large number in which delivery was accomplished by tearing through the cervix.

It is impossible for any practitioner to resist the importunities of the patient and her friends unless he has a good clinical experience of this stage of labour, and can assure them that all will be well with a little patience. The old method of attending a certain number of cases gave no experience in this stage, but it is being emphasized more and more to-day. In my own opinion—and I think all teachers will agree with me—a student learns much more from attendance upon one case from the earliest part of the first stage to the completion of labour than upon a large number observed only at the end of the second stage.

4. Importance of making a Definite Diagnosis before applying High Forceps.

Forceps applied when the head is on the perineum often relieves the patient of much suffering, and with many practitioners this is frequently done with great benefit, but forceps applied while the head is at the brim should be a rare operation, much rarer than it was in the past, as Caesarean section has removed a large group which otherwise would require this operation, and it should be performed much less frequently than it is.

A rule should be made never to apply forceps with the head at the brim merely because of delay in labour; a definite diagnosis of the cause of delay should be made, and if the practitioner cannot make this diagnosis he should call in the help of one with more experience.

Occipito-posterior positions are very difficult to diagnose, and those with the greatest experience will acknowledge that they frequently fail to make this diagnosis until the cervix is sufficiently dilated to allow a hand to be passed through the external os and the ear of the child felt; this is an infallible method, and if only it had been carried out before applying forceps this very large group in our series would have been eliminated. The treatment of occipito-posterior cases is simple, as Dr. Miller has pointed out; if only they are recognized, but to apply forceps at the brim with the head in that position is to court disaster.

If only the rule of making a definite diagnosis before applying forceps at the brim had been carried out practically all the cases in our series would have been eliminated, and many other women, in whom delivery was accomplished, would have been saved from mutilation. With this rule faithfully observed forceps would not have been applied before full dilatation of the cervix, while cases with contracted pelvis and occipito-posterior positions would have been recognized and easily treated.

This is a subject which behoves us, as a profession, to study carefully; it has been brought forward in no spirit of carping criticism, but in the hope that a free discussion between so many of us interested in obstetrical work, but looking at it from many angles, will result in practical advancement.

REFERENCE.

¹ *Journ. Obstet. and Gynaecol. Brit. Empire*, vol. xxviii, p. 351.