LIONEL COLLEDGE AND SIR C. BALLANCE: ANASTOMOSES BETWEEN THE RECURRENT
LARYNGEAL AND PHRENIC NERVES.

FIG. 14.

FIG. 15.

FIG. 16.

A. WATT, R. C. J. MEYER, AND A. CAMPBELL: OPERATION FOR PITUITARY TUMOUR. (See p. 748.)

FIG. A.—Showing position of the tumour.

SYDNEY SMITH: IDENTIFICATION FROM A FINGER-TIP REMOVED BY A BITE. (See p. 757.)

FIG. 1.—Tip of finger removed by bite.

FIG. 2.—Finger-print of right index finger of accused.

typical cross striated muscle fibre is dependent on the impulses from the central nervous system to set it into activity.

The Stimulation of the Cortex Cerebri.

Before the monkey was killed the right cerebral cortex was stimulated with the faradic current, a Sherrington single electrode being employed for the purpose (see Fig. 16).

DESCRIPTION OF PHOTOMICROGRAPHS ON PLATE.

FIG. 14.—Photomicrograph of muscle bundles in right half of diaphragm ($\times 700$). Three years and four months after recurrent laryngeal-phrenic end-to-side nerve anastomosis: showing that this method of operation has left no permanent effect on this half of the diaphragm.

FIG. 15.—Photomicrograph of muscle bundles in left half of diaphragm ($\times 350$). Two years after recurrent laryngeal-phrenic end-to-end nerve anastomosis. The distal end of the phrenic nerve was not anastomosed to any other nerve. No transverse striation is shown in the photograph. On examining many other sections faint striation of a muscle bundle was occasionally seen, and this perhaps was due to the fact revealed at the autopsy that below the point of section of the phrenic nerve a small nerve from the brachial plexus joined the phrenic nerve in the thorax.

FIG. 16.—Drawing of right half of brain of the monkey (natural size). A, Faradic stimulation of the cortex at this point caused elevation of the upper lip on the left side. B, Stimulation of the cortex at B caused abduction of the vocal cords. Unfortunately the point of the cortex at which the maximal adduction movement of the vocal cords occurred was not marked. It was in front of the area which on stimulation caused abduction of the vocal cords. The adduction movement was much stronger than the abduction movement, which was distinct from the abduction movement associated with the respiratory rhythm.

The cortical areas stimulated correspond to those originally defined by Semon and Horsley. The interest of the cerebral stimulations carried out by us is that the nerve impulses passed along the phrenic nerve, then through the phrenic-recurrent laryngeal end-to-end anastomosis to the recurrent laryngeal nerve and intrinsic laryngeal muscles.

Semon and Horsley³ showed that unilateral stimulation invariably produces bilateral effect. They at first thought that there was no abduction centre in the brain cortex, but they demonstrated that direct excitation of the accessory nucleus in the medulla always evoked abduction of the vocal cords. They found no difficulty in evoking complete bilateral adduction movement of the vocal cords by stimulating an appropriate area of the cerebral cortex. If the stimulation is prolonged the "besoin de respirer" overcomes the influence of the artificial stimulus and evokes a powerful abduction movement of the vocal cords. They explained these diverse effects in this way: Phonation (that is, adduction) is a purposive or volitional act, while abduction is essentially related to the so-called automatic processes of organic life—that is, abduction serves inspiration.

In the later paper these authors show that there is a definite area on the cortex of the brain, stimulation of which causes abduction of the vocal cords. They point out that Krause (1878) was the first to localize the phonatory area in the cortex of the dog's brain. Ferrier, however, had previously reported that excitation of a certain cortical area in the dog elicited barking. Risien Russell⁴ described separate areas of the cortex of the brain in which excitation evoked the movements of abduction and adduction of the vocal cords. Risien Russell also separated the abductor from the adductor nerve bundles in the recurrent laryngeal nerve—the abductor fibres being situated on the inner side of the nerve, while the adduction fibres are on the outer side. Stimulation of the abductor fibres produced unilateral abduction movement only, while division of the abductor fibres caused atrophy of the corresponding abductor muscle. We have been unable to separate the abductor nerve bundle in the way described by Risien Russell.

Operation in Man.

It would appear that the proper operation to perform in man in cases of paralysis of a vocal cord is recurrent laryngeal-phrenic nerve end-to-side anastomosis. This has been successful in the one case in which this operation has been carried out.⁵ Independent of the cases in man in which by stab or gunshot wound, or during an operation, the recurrent laryngeal nerve has been divided, and in which an immediate anastomosis should be effected, we are of opinion that if a vocal cord (or cords) has been para-

lysed for three months with no sign of improvement in the condition, a recurrent laryngeal-phrenic nerve anastomosis should be forthwith carried out.

Note.—Figures showing regeneration of the phrenic and recurrent laryngeal nerves after similar anastomoses had been effected were published in the *British Medical Journal*, March 26th, 1927 (see Figs. 5 and 6 in plate facing p. 555).

The experimental nerve anastomoses were carried out at the National Institute for Research, Mount Vernon, Hampstead. The microscopic specimens and photomicrographs were prepared by Miss Glascock and Mr. Stewart at the Royal College of Surgeons.

REFERENCES.

- ¹ See *British Medical Journal*, March 26th and April 2nd, 1927. ² *Ibid.*, April 2nd, 1927. ³ *British Medical Journal*, 1889, and *Trans. Roy. Soc.*, 1890, B. ⁴ *British Medical Journal*, 1892, and *Proc. Roy. Soc.*, 1895. ⁵ *British Medical Journal*, July 30th, 1927, p. 158.

OPERATION FOR PITUITARY TUMOUR.

REPORT OF A CASE.

BY

ANDREW WATT, M.B., F.R.C.S., R. C. J. MEYER, M.D., F.R.C.S.,

AND

ANDREW CAMPBELL, M.B., F.R.C.S.,

GENERAL HOSPITAL, JOHANNESBURG.

(With Special Plate.)

A EUROPEAN, aged 23 years, complained that during the last five or six years his left eye had been noticeably weaker than the right. In November, 1926, he suffered from severe headaches when reading for an examination. In February, 1927, when marking the register, he found that when he thought the pen was touching the book it was actually about an inch away from it; he was unable to see with the left eye alone. There was no complaint of headache, no vertigo, and no vomiting. At the beginning of April Dr. Meyer was consulted, and the patient was sent into hospital for investigation under Dr. Andrew Watt.

The previous history showed that the patient had fallen off a horse in 1918, but though unconscious for some time he suffered no other ill effects. In 1919 he had influenza. In 1920 he was hit over the left eye by a cricket ball. There is no family history of nervous or eye trouble, and the father, mother, brothers, and sisters are all well. He smokes ten to fifteen cigarettes daily and is a total abstainer. His condition on admission to hospital was recorded as follows.

The patient is a well-built, healthy looking individual, about 5 ft. 8 in. tall. There is no abnormality in his gait and the grips of both hands are normal. There is no tremor; the motor functions and the various reflexes are normal, as is sensation. The bowels are regular and the appetite is good; there is no frequency of micturition, and urine analysis reveals nothing abnormal. The heart and lungs are sound, and the Wassermann reaction is negative.

Examination of the cranial nerves shows the sense of smell to be normal; vision is impaired, pupils react to light and accommodation, there is no ocular paresis, no nystagmus, and no strabismus. There is no anaesthesia of the face or scalp. The functions of the seventh nerve are normal, and there is no impairment of taste. The movements of the tongue and swallowing are normal. The basal metabolic rate is normal.

Blood sugar before glucose meal (200 grams)	0.11 per cent.
" " 30 minutes after glucose meal ...	0.16 "
" " 60 " " " " " " " "	0.14 "
" " 90 " " " " " " " "	0.13 "
" " 120 " " " " " " " "	0.14 "

Sugar was not found in the urine before the glucose meal, but a trace was present two hours after. The cerebro-spinal fluid was under a pressure of 250 mm. water. The globulin was not in excess of the normal, and the sugar percentage was 0.09. The sodium chloride figure was 720 mg. per 100 c.cm. No characteristic curve was obtained with gold solution. A cell count yielded a total of two lymphocytes per c.mm.

A blood count taken on June 18th was as follows: haemoglobin 95 per cent.; colour index 0.98; red cells 5,810,000; leucocytes 8,000; polymorphonuclears 52 per cent.; large mononuclears 2 per cent.; lymphocytes 46 per cent. The red cells showed slight anisocytosis with a very moderate degree of polychromasia.

The patient was referred to Dr. Campbell for examination of the ear, nose, and throat, and for the vestibular tests. Both drumheads were retracted, and he could hear a whisper at over 16 feet on both sides. Tuning-fork tests were normal. There was no spontaneous nystagmus, no spontaneous past-pointing, and

no inco-ordination; the body balance tests gave normal reactions, and diadokokinesia was present.

Vestibular Tests:

Turning to right gave an after nystagmus to left lasting 25 seconds.

Turning to left gave an after nystagmus to right lasting 35 seconds.

Turning to right past-points 6 inches to right with left arm.

Turning to right past-points 4 inches to right with right arm.

Turning to left past-points 10 inches to left with right arm.

Turning to left past-points 6 inches to left with left arm.

Cold syringing the left ear at 68° F. elicited a rotary nystagmus to the right in 45 seconds and past-pointing to left with both arms. Cold syringing the right ear at 68° F. elicited a rotary nystagmus to the left in 25 seconds and past-pointing to right with both arms. It was concluded that a supratentorial lesion was present.

OPERATION NOTES BY DR. CAMPBELL.

The diagnosis of a pituitary tumour was established and confirmed by beautiful stereoscopic films taken by Dr. Andrew Watt. The eye tests indicated pressure on the chiasma, perhaps more on the left than on the right side. This finding was corroborated by the vestibular tests. (For skiagram see Fig. A in special plate.)

On June 13th, 1927, under rectal anaesthesia, given by Dr. F. B. Mudd, a submucous resection of the nasal septum was performed in the usual way after removal of both middle turbinates and a slice of the right inferior turbinate which happened to be in the way. The nasal mucous membrane was sprayed with 1 ounce of 10 per cent. cocaine and adrenaline solution twenty minutes before operation. The anterior wall of the sphenoidal sinuses was thin and easily removed. The sphenoidal sinuses came into view, and a rounded tumour was found to occupy the upper part of the cavities. (See Fig. 1.) There was no septum between

FIG. 1.

the sinuses, but there were one or two spicules of bone which indicated its site.

A small piece of bone was removed from the most dependent part of the tumour by means of a gouge, and the rest of the tumour was uncapped with suitable forceps. There was little bleeding. The mucous membrane of the sphenoidal sinus was removed.

The result at this stage was that the tumour covered by dura was occupying the upper half, or almost half, of the sphenoidal sinuses. On removal of the very thin bone there was no dropping downwards of the tumour. A syringe introduced into the growth did not draw liquid, but gave the impression that the tumour was solid. No packing was introduced into the sphenoidal sinus. The only communication to the nose was through the sphenoidal ostia. The muco-periosteum covering the anterior wall of the sphenoidal sinuses had been reflected intact. A large piece of the nasal cartilage was slipped back between the septal flaps, and each side of the nose was packed with a glove finger into which gauze had been introduced, so as to keep the septal flaps lightly together. The operation lasted one and a half hours. After leaving the theatre there was some respiratory difficulty, and the patient did not come round for six hours.

On the following day there was a suspicion of an acute appendicitis, but this pain was probably due to the rectal ether. The nasal packing was removed in twenty-four hours. A mild antiseptic spray was used for another twenty-four hours, after which the nose was gently irrigated with saline solution thrice daily. The recovery from the first stage of the operation was uneventful. In a week he was up and walking about the ward. No improvement in eye symptoms followed at this stage.

On June 27th, under open ether anaesthesia and cocaine spray to the nasal mucous membrane, the operative field was reopened through the nasal septum. It was noticed that a large part of the cartilage which had been replaced had been absorbed. The sphenoidal sinuses were easily

entered. The tumour had not descended, and the whole sinus and dura were covered with granulations, which bled considerably. There was no difficulty in making a crucial incision of the dura, though the growth was probably also incised in the attempt. No descent of tumour was observed during the operation; and there was no suggestion that the growth was anything but solid. There was no escape of any liquid other than blood; bleeding after incision of the dura was easily controlled. The operative field was closed by bringing the septal flaps together without any packing in the sphenoidal sinus; fingers were again utilized to keep the flaps in apposition. Towards evening the patient was rational and appeared to be doing well. On the following day, though he had passed a good night, he did not seem quite rational. He was able to recognize the surgeon and nurses, but answered "Yes" to all questions and nothing else; by the evening he failed to recognize anyone. The temperature rose to 102° F., and we feared that meningeal symptoms were commencing, though there was no other indication of this.

On June 29th he woke up perfectly rational and rapidly improved, with a gradual drop in the temperature, which became normal on June 30th. After this he gave us no anxiety; he was allowed to get up a little on the eighth day after the second operation, by which time he was conscious of improvement in his eyesight. He was discharged from hospital on July 12th.

FIG. 2.—Visual fields, showing condition at dates before and after operation. Left vision=fingers at 2 metres; 25 mm. white. Right vision=6/5; 5 mm. white.

It was thought that a decompression by two stages was a safer procedure than by one operation. No thought of any attempt to remove the tumour was entertained. It is, however, possible that if the tumour descends and fills up the sphenoidal sinuses so that further decompression is not possible, then we may with greater safety remove the anterior muco-periosteal wall of the sphenoidal sinuses, leaving the posterior edge of the nasal septum *in situ*, and thus expose the tumour mass through windows which will remain permanently open. The tumour itself may then be attacked surgically or made to shrink by electro-coagulation.

The main danger of the trans-sphenoidal route is sepsis, laying bare the field of operation to invasion by organisms from a possibly septic nose or nasopharynx.

An attempt to remove the tumour at the first operation would probably, almost certainly, have resulted in infection of the intracranial structures and death; whereas by gradual stages, and as the occasion demands, this three-stage removal has much to recommend it from the point of view of safety. No attempt was made even to remove a small piece of tumour for the microscope because of the possibility of infection. No purpose would be served by an accurate knowledge of the pathology of the tumour. If it were found to be malignant nothing could be done, and if it proved to be innocent we could still do nothing more for the present. It is possible that in the descent of the tumour it may become strangulated by the narrow bony neck through which it must pass; this will not occur until the sphenoidal sinus is completely filled.

The fields taken by Dr. Meyer on October 4th, compared

with the fields before operation, show a considerable improvement in the patient's eyesight, especially in the right eye. There is a small perforation in the nasal septum, and there are a few adhesions between the septum and the lateral nasal wall. The anterior walls of the sphenoidal sinuses are easily inspected. There is no evidence of sepsis in the nose.

The patient was first seen in April, 1927, examined thoroughly and treated by surgical operation in June, and reported on after examination on October 4th.

NOTES BY DR. R. C. J. MEYER.

Objective examination showed no abnormality as regards the outward appearances, motility, and pupils; the Wernicke pupil reaction was demonstrated later. The fundus was healthy, but there was slight hypermetropic astigmatism; the subsequent wearing of glasses stopped all headaches. Right vision = 6/6, J.1. Left vision = hand reflex centrally, but fingers at 3 metres nasally. This called for examination of fields, the broad results of which are indicated in the diagrams. Fig. 3, A (right eye);

FIG. 3.

shows the right upper outer quadrant lost; the patient was quite unconscious of right quadrant loss of field until examination demonstrated it to him. In Fig. 3, B (left eye), the temporal hemianopia is seen to cross the vertical middle by about 15 degrees. The patient was quite conscious of loss of vision of left eye, which drove him to seek medical aid.

The examination suggested the presence of a pituitary tumour involving the left side much more than the right, but later on under portion of chiasma. I handed the case over to the physician, radiologist, and surgeon, but the patient could only come for observation (May 28th).

Two operations were performed—one on June 13th and the other on June 27th. On July 13th, at the request of the surgeon, fields were again taken (there being no change in any other condition of the eyes). The right upper quadrant scotoma had now diminished to a smaller sector. (See Fig. 4, A.) The left scotoma had also diminished. (See Fig. 4, B.)

FIG. 4.

On September 1st a friend stated that the patient was "quite better," which meant, at any rate, a decided improvement. (N.B.—Patient lives in the country and over a hundred miles away.)

FIG. 5.

On October 4th the patient unexpectedly arrived for examination. This showed the right scotoma to be still further diminished. The left scotoma was also diminished somewhat. Right vision = 6/5, J.1. Patient does not feel the scotoma, and this does not exist for luminous objects at all now. Certainly a most striking improvement. (See Fig. 5, A.) The left disc shows great pallor of nasal side—that is, atrophy is showing actually at the disc. (See Fig. 5, B.)

THE AFFINITIES BETWEEN RHEUMATISM AND TUBERCULOSIS.

BY

W. CAMAC WILKINSON, M.D., F.R.C.P.,
LONDON.

TUBERCULOSIS connotes the morbid changes caused by living tubercle bacilli in the living tissues of man and animals. As the tissues react they present lesions of endless variety in degree, kind, and time. The outstanding and characteristic lesion is the tubercle with its epithelioid and giant cells, but there are also lesions as acute as those of septicaemia or pyaemia, and, on the other hand, as chronic as those of chronic rheumatism. Indeed, it seems likely that many so-called "rheumatic" affections, which do not yield to antirheumatic treatment, are really manifestations of chronic tuberculosis. Even joint affections accompanied by pain and swelling may be tuberculous in nature.

The brilliant investigations of Liebermeister, with the help of his new acetic acid-antiformin method, seem to establish the frequency of chronic inflammation free from any tubercles as a distinct and even independent lesion of tuberculosis. These chronic inflammatory lesions are, indeed, often the end-results both of acute lesions and of true histological tubercle, but Liebermeister holds that they may be independent of either of these lesions and are common in many organs and parts. They have been overlooked because death does not occur in this chronic stage, and there is no chance of seeing them. Such lesions may be the *fons et origo* of the multifarious disturbances of function occurring in the early stages of chronic tuberculosis. The lesions are the most chronic that occur in tuberculosis, and cause mainly mild and transient symptoms. Thus it may be impossible after a long lapse of time to trace them to their true cause or fix the time of their occurrence. Living tubercle bacilli may be found in these chronic lesions, and evidence supports the view that these quiescent lesions may be reactivated. These lesions are most common in the secondary stage of chronic tuberculosis, when tubercle bacilli, in small numbers, more or less isolated, and certainly not in clumps or masses, are circulating in the blood, before tertiary disease occurs in the apex of the lung. Amongst other manifestations, tuberculosis may cause lesions suggesting rheumatism of joints with painful swellings, in some cases with erythema nodosum, in others with erythematata, pains also in muscles and in one or more joints, pains in the chest without signs of pleurisy, pains in the neck and in the eyes or head, disturbance of the heart's rate and rhythm—even pains in nerves (sciatica). Liebermeister has demonstrated the presence of these chronic inflammatory lesions notably in lymphatic glands, serous membranes, even in the meninges, in the lungs, liver, kidney, spleen, veins, skin, eyes, and larynx. May not these lesions be responsible for the various symptoms that may and do occur and pass away in the early stages of chronic tuberculosis, and especially of chronic apical tuberculosis?

For the most part these lesions elude us, and we can only bring them to light by means of tuberculin. Twenty years ago I suggested the possible existence of these lesions because doses of tuberculin excited pains in various organs and parts in a manner that seemed to indicate a focal reaction. For many years I have observed pains in special muscles and joints, time after time in the same part after each dose—in the hip, or knee, or little finger, or thumb, or heel; in the back or shoulder; in bones or muscles, such as severe cramps in the calves of the legs; pains also in the epigastrium, doubling up the patient after each dose, and even boring or lancinating or cutting pains in the shoulder radiating down the arm, or deep pain in the chest. There are strange individual idiosyncrasies, because in the same patient, after injections of tuberculin, the same parts are apt to suffer, though the site of injection varies. These pains, in fact, may be related to chronic inflammatory lesions and may be really focal reactions. One has great opportunities of studying these pains, their varieties and situations, after injections of tuberculin. Naturally there is no other way of investigating them.