a few cases. The tongue was nearly always coated at the dorsum and red at the margins. Tonsillitis and pharyngitis were fairly common. No parotitis was met with. Nausea and vomiting were usual at the commencement, but rarely persisted throughout the disease. Constipation was the rule, and was followed by diarrhoea at the terminal crisis. Dysenteric symptoms were present in a few cases throughout the malady. No changes were observed in liver or spleen that could be attributed to dengue.

Urinary System.—Albuminuria was common but transient. One case of acute nephritis was reported. Chronic kidney disease is likely to end by uraemia as the result of an attack of dengue. Diabetic patients when attacked sometimes developed a fatal coma, even during convalescence. Epididymitis, orchitis, cystitis, etc., were not met with.

Complications.—These were rare. Those met with were: pharyngitis, rhinitis, gastritis, enteritis, nephritis, and thrombosis.

Sequelae.—Itching of body, irritation of palms and soles, mental depression, vertigo, motor paresis, peripheral neuritis, staccato speech.

Children were just as liable to be attacked by dengue as adults. Males and females were affected equally. No deaths occurred among children.

Convalescence may be short or long—one day to three months. Patients lose weight, and usually suffer from mental depression, debility, loss of appetite, bradycardia, migraine, and neuritis. Relapses are rare.

Diagnosis.

In this season (December) in Egypt dengue has to be diagnosed from the following diseases:

(a) Malaria: by examination of blood films, size of spleen and liver, time incidence of febrile attacks, and rapid pulse.

(b) Measles: starts less suddenly; presence of Koplik's spots, coryza, absence of pains, rapid pulse, and deep red rash.

(c) Influenza: is usually accompanied by respiratory involvement, rapid pulse, and absence of rash.

(d) Scarlet fever: diagnosed by presence of throat symptoms, rash, rapid pulse, absence of pains, and blood examination.

(e) Small-pox: in the first three days of dengue it is very difficult to differentiate it from small-pox. Later on the course of the malady soon manifests its nature. History of vaccination may help.

(f) Articular rheumatism: diagnosed by history of malady, presence of acid sweats, and joint lesions.

The treatment is symptomatic and prophylaxis involves antimosquito measures.

REFERENCES.

¹ *British Medical Journal*, June 27th, 1925, p. 1185. ² *Ibid.*, September 11th, 1926, p. 489.

DUPLICATION OF THE SPINAL CORD.

BY

F. PARKES WEBER, M.A., M.D., F.R.C.P.

AN excellent review of the known cases of partial doubling of the spinal cord was written in 1906 by A. Bruce, Stuart McDonald, and J. H. H. Pirie¹; for the reference to it I am indebted to Dr. Kinnier Wilson. Nevertheless, the following case perhaps deserves a note.

The patient, a well-developed man, aged 51, was admitted to hospital on March 29th, 1928, with a spastic parietic ataxic gait and signs suggesting combined degeneration of the spinal cord. He had a pale, sallow face; the liver was considerably enlarged, and the spleen could be felt, reaching just below the ribs. The gastric contents after a test breakfast contained no free hydrochloric acid. The erythrocyte count was 3,392,000 and the haemoglobin was 70 per cent. The white cells were 17,800 (polymorphonuclears 74 per cent., lymphocytes 17 per cent., monocytes 9 per cent.). The Wassermann reaction was negative. There was a little irregular pyrexia, and apparently a remnant of pneumonia in the middle part of the right lung, which possibly explained the presence of a polymorphonuclear leucocytosis. According to the history obtained the patient had noticed something wrong with his feet in August, 1927, and since December, 1927, he had been taking an "ox-liver juice" twice daily for anaemia. The result of the ox-liver treatment was said to have been good, till weakness in his legs developed ten days before admission.

In the hospital, in spite of liver diet, etc., the weakness in the legs increased, paradoxical incontinence of urine and cystitis, with bedsores, developed, and death occurred on April 26th.

The necropsy and microscopical examination confirmed the diagnosis of combined degeneration of the spinal cord; but during the macroscopical examination it was noticed that, close along the left side of the lower part of the spinal cord, there was another cord-like body of the same consistence as the cord, but smaller (about 1 cm. in diameter) and without nerve roots proceeding from it. Its upper end arose from the substance of the cord by a base somewhat over 1 cm. in diameter, the uppermost part of its insertion being $7\frac{1}{2}$ cm. above the terminal point of the conus terminalis. At its lower (free) end it tapered to a point, which was $1\frac{1}{2}$ cm. above the lower end of the conus terminalis.

Transverse microscopical sections of this macroscopically tumour-like appendage of the spinal cord showed it to be quite obviously a fairly well-developed duplicate cord with a central canal, etc. This was kindly confirmed by Dr. J. G. Greenfield, who also confirmed the fact that the duplicate cord showed well-marked signs of combined degeneration, just as did the main spinal cord (from which the cauda equina arose).

When I saw the long polypoid tumour-like appendage lying along the left side of the lower portion of the spinal cord, I at first thought that I had made a mistake in the diagnosis, and that the patient's paraplegia, etc., might have been due to a tumour; but microscopical examination, as already stated, completely cleared up the nature of the tumour-like structure, proving it to be a congenital abnormality, a doubling of the lower part of the spinal cord. One may wonder whether in this case lipiodol would have been partially arrested at the site of origin of the duplicate cord had it been used to ascertain the site of a possible tumour.

REFERENCE.

¹ A Second Case of Partial Doubling of the Spinal Cord, *Rev. Neur. and Psych.*, Edinburgh, 1905, iv, p. 6.

THE ETIOLOGY OF MONGOLISM;

WITH A CASE OF MONGOL TWIN.

BY

HUBERT ARMSTRONG, M.D. VICT. AND LIVERP.,

SENIOR HONORARY PHYSICIAN, ROYAL LIVERPOOL CHILDREN'S HOSPITAL.

THE occurrence recently, in my hospital practice, of a case of twins, one of whom was a Mongolian imbecile, the other not, stimulated investigation into the etiology of that distressing and increasingly common type of mental deficiency. But the *ultimate* cause of Mongolism is, I suppose, as yet unknown, nor can I find much to elucidate it.

The peculiar history so often obtained—namely, that the child is either the first of a very young mother, or, as often happens, is born many years after the previous one, and most frequently of a mother nearing the menopause—warrants the assumption that this developmental error is in some way due to immaturity or exhaustion of her procreative powers.¹ At what stage in the development of the ovum or foetus this makes itself felt has been the subject of a good deal of argument.

The problem is this: Is Mongolism inherent in the unfertilized ovum, or is it acquired after fertilization? A third proposition might be that Mongolism is imparted to the ovum in fertilization, were it not that the history appears to point to the maternal factor so constantly. As a generality it may be assumed that parents are approximately the same age, the father being on the average a year or two the elder. This would make the odds even as between the two parents, were it not that there is no evidence, known to me, that the children of elderly men tend to Mongolism otherwise than when their wives are approaching the menopause. Nevertheless, this possible factor cannot be entirely brushed aside without consideration.

An argument in favour of the post-fertilization embryonic theory is that Mongols are so often the subjects of other congenital abnormalities. Of these the commonest are malformations of the heart, followed a long way behind by congenital cataract—defects which obviously cannot