Remarks

ON

URTICARIA: ITS PATHOGENESIS AND ETIOLOGY.*

BY

RUPERT HALLAM, M.D.,

Honorary Physician in Charge of the Skin Department,
Royal Infirmary, Sheffield.

THE subject of urticaria is one of considerable importance, for the complaint is not infrequently met with both in private and in hospital practice, and also because of the distress it occasions its unfortunate victims. To the dermatologist it is of peculiar interest, for it exemplifies, perhaps better than any other skin complaint, the close relationship between the science of cutaneous medicine and the other branches of medicine. It serves to impress upon us the fact that a change in the skin is often the expression of some pathological process which is taking place in another part of the body, and that its elucidation and treatment require a knowledge of the latest advances in biology and physiology.

The signs and symptoms of the condition are fortunately easily recognized, although the wheals vary enormously in shape and size. Occasionally, it is true, there may be a confusion with the bullous rashes should the horny layer of the urticarial wheals be raised in the centre by an excess of serum, but this is an infrequent occurrence. On other occasions the oedema of a limb or other part of the body may be so pronounced as to earn the distinctive name "angioneurotic oedema," a term which may preferably be replaced by that of "giant urticaria."

It is convenient, perhaps, to follow the line of the majority of textbooks and divide the subject into four main groups: (1) factitious urticaria, (2) acute urticaria, (3) chronic urticaria, (4) papular urticaria of childhood; but to obtain an accurate conception it is necessary to realize that it is not a disease, but a symptom attributable to one of many causes.

Urticaria Factitia, or Dermographism.—This peculiar response of the skin to local trauma may or may not be concomitant with urticaria vulgaris. Lewis, in his classical work on the mechanism of wheal production, states that both this form and the wheal secondary to internal causes is due to the release of a histamine-like substance from the cells of the skin. Why the person with urticaria factitia acquires this exaggerated susceptibility of the cells of the skin is not clear. The condition is met with in apparent good health. On the other hand, the incidence is higher in epileptics and alcoholics. I recently saw well-marked dermographism begin during the course of secondary syphilis.

Acute Urticaria.—It is common knowledge that this state usually follows the ingestion of some particular article of diet, and is often accompanied by more or less severe constitutional symptoms such as vomiting, diarrhoea, and a general feeling of malaise. The rash is usually profuse, and sometimes even bullous. It may involve the mucous membrane of the lips, tongue, or glottis. The latter complication, however, is rare, but cases are recorded in which tracheotomy has been necessary. Fortunately this drastic remedy may now be obviated by the hypodermic injection of adrenaline.

Chronic Urticaria.—It is this chronic, recurring stage which exercises the skill and patience of the physician, for the duration may be measured in terms of weeks and months or even years, and it may be due to an embarrassing variety of causes.

Papular Urticaria.

Urticaria in the true sense of the word is not often met with in childhood, but the clinical entity known as papular urticaria, or lichen urticatus, bears such a close relation-

ship to chronic urticaria that it is usually classified with it. There are, however, clinical, anatomical, and etiological distinctions between the two. The disease is only met with in childhood, and is characterized by the successive crops of small urticarial wheals, in the centre of each of which is a small papule. The wheal disappears within a few hours, but the papule remains for about a week. The causation is obscure, but there is a curious factor common to all, to which I have drawn attention in previous communications—namely, that the sufferers recover immediately they are admitted to hospital, even though no treatment of any description be given. The rash returns after their discharge. Based on this experience, I have been engaged in numerous experiments during the last few years with a view of ascertaining whether there is a specific antigen responsible for the complaint, but so far have failed to detect one.

Before discussing the etiology of urticaria in detail I think it is expedient to consider briefly the present-day conception of

The Histopathology of Wheal Formation.

The actual microscopical changes have been accurately described by Gilchrist,¹ Lewis,² and others. The former excised a portion of the affected skin at varying intervals after wheal formation, and found a local congestion of the cutaneous vessels, an escape of serum through the capillary walls, and also definite signs of inflammatory change in the later stages. As previously mentioned, Lewis has shown that this phenomenon of wheal formation is exactly reproduced by an injection of histamine.

Change in the blood in the form of retarded coagulability and deficiency of calcium content were described by Wright,³ but other observers have not always been able to confirm either change. During recent years it has been recognized that the victim of this reaction of the skin is often one who is temporarily or permanently the subject of some peculiar metabolic change, which has produced an increased susceptibility to irritants. Also, that this change is a factor in the close relationship between eczema, asthma, hay fever, and urticaria; for the same individual is not infrequently a sufferer from more than one of these. The nature of this instability or sensitiveness, which is often referred to as a state of allergy, has given rise to much experimental research, but as yet the various conjectures are hypothetical. It is sufficiently clear, however, that although an organ such as the skin or the lungs shows clinical manifestation of the hypersensitive state, it must not be assumed that the particular organ is the sole site of the allergic state. It is possibly governed by highly complex changes in the cells of the individual or in the colloid stability of the blood. Allergy is found as a family disposition. The Jews are particularly subject to it; the North American Indian is not, and apparently it may be hereditary or acquired.

The Relationship to Anaphylaxis.

The conception that urticaria might be a manifestation of the anaphylactic state was advanced by Wolff-Eisner⁴ soon after the peculiar phenomenon of anaphylaxis was first observed in animals. This theory has received considerable support, and it is generally acknowledged that, although there is a wide difference between the condition of an animal which has been reinjected after an interval of ten days with some foreign protein and a patient suffering from urticaria, yet there is a close resemblance between the two states. This is particularly evident in urticaria following serum injections, but Vidal⁵ and others have shown a similarity between the alimentary type of urticaria and anaphylactic shock, for in both there is a fall of blood pressure, leucopenia, moderate eosinophilia, and sometimes albuminuria. Again, Boycott⁶ found that a flea does not produce a wheal after the first bite unless his host has been previously bitten by the same species of flea. A case described in *La Presse Médicale* last year also illustrates the similarity. The patient, after undergoing moderate exertion for a few minutes, not only suffered from a profuse urticarial rash, but in addition showed signs of dyspnoea, vertigo, profound cardio-vascular disturbance, and collapse.

* Made in opening a discussion in the Section of Dermatology at the Annual Meeting of the British Medical Association, Cardiff, 1928.

An examination of the blood revealed changes usually associated with anaphylactic shock.

In some cases urticaria follows the ingestion of or contact with a substance to which the patient has not been previously exposed; therefore anaphylaxis does not always serve to explain the reaction, and we have to fall back on the term "sensitization"—the inexplicable predisposition or idiosyncrasy to react to stimuli which have no effect on a normal individual. There is evidence to show that the antigen may be ingested, inhaled, or absorbed through the skin, and yet produce the same results. Also that the so-called sensitiveness exists in the serum, for Prausnitz and Küstner⁷ injected the serums of hypersensitive patients into the skin of normal individuals and found that the site of the injection was temporarily sensitized to the same substance. This was confirmed by A. Walzer and M. Walzer,⁸ who noted that the serum possessed this activity for some months after it was withdrawn from the sensitized person. The susceptibility may be transmitted by a female guinea-pig to her offspring (Rosenau⁹). Another observation which has an important bearing on the treatment is that when once acquired the sensitiveness to a particular antigen appears to be followed by sensitiveness to other antigens.

May this reaction of the skin be of some protective significance? This point is raised by Memmesheimer¹⁰ in relationship to eczema, but there is little to support this hypothesis. The histology of the wheal certainly shows that cell damage takes place, and therefore cell products escape into the general circulation. We know that immunity follows an attack of measles and scarlet fever, diseases in which the skin is particularly affected; but we are equally cognizant of the fact that in many other diseases involving extensive areas of skin no protection is acquired.

Etiology.

Although the conception of the biological changes associated with an attack of urticaria has changed considerably in later years, it is interesting to note that the dermatologists of the middle of the last century fully recognized the diversity of the causes of the reaction. In every textbook, ancient and modern, there is a long list of both external and internal causes, and almost every observer is able to add to the list. Hebra noted that the introduction of a uterine sound produced a profuse urticarial rash in one of his patients. Crocker¹¹ quotes a case in which a ring at the front door would determine an attack. We know that it may mask a scabies infection; that it follows the ingestion of some particular food or drug. Furthermore, it may be one of the earliest signs of a grave disease such as leukaemia or diabetes; also, it may be associated with focal sepsis. The latter coincidence has been so frequently demonstrated in recent years that it must be accepted. My own view is that focal sepsis plays a far greater role in the causation of chronic urticaria than food protein, and that in the past it has been assumed too readily that urticaria is the outward manifestation of food poisons.

As the chance of successful treatment largely depends on the detection of the cause, it is essential that a case of urticaria should be approached with the greatest circumspection, and also with the knowledge that there is no specific remedy. A detailed history will be required; the skin will need careful inspection to exclude the possibility of local irritants. Protein sensitization tests must be considered, although these have not given the assistance that was anticipated from them. Each system will require examination, for none is above suspicion. The key to the mystery may be the examination of the blood. In a recent article F. R. Menegh¹² claims that disease of the biliary tract was an etiological factor in 50 per cent. of his cases. A high incidence of urticaria has been described in patients with syphilis, and I have met with this coincidence on several occasions, but antispasmodic treatment has not prevented the recurrence of the wheals.

It is sufficiently obvious that there are innumerable possibilities to be considered, and that the physician who successfully detects the offending abnormality has good reason to congratulate himself. That trauma plays an

important part as an exciting cause was demonstrated as long ago as 1888 by Jaquet, who found that when a limb of a patient suffering from generalized urticaria was protected the wheals did not arise on the skin of that limb.

Desensitization.

Should the offending antigen be detected, desensitization may be attempted by giving minute, increasing doses until tolerance is established. On the assumption that urticaria is a manifestation of the anaphylactic state French dermatologists have recently advocated the ingestion of 0.5 gram of peptone one hour before each meal.

Empirical Remedies.

Calcium salts have had a long innings, but in my experience the results have been singularly disappointing, and in my opinion confidence in the supposed virtues of this drug have many times been responsible for an inadequate examination of patients. Urticaria has been noted in hypothyroidism; therefore thyroid extract has recently been prescribed as a remedy; one author, indeed, proclaims it as a specific. I do not go so far with him, but believe, in the unhappy event of the actual cause not being ascertained, that thyroid extract is worthy of a trial. Papular urticaria is influenced neither by dieting nor drugs, but, as already mentioned, the patient obtains a respite if he be removed from his own environment. True, the rash often recurs when the patient returns home, but I have noted a permanent benefit in a number of instances, and recommend a trial when practicable.

REFERENCES.

- ¹ Gilchrist: *Journ. Amer. Med. Assoc.*, 1896, 27, 1222. ² Lewis: *The Blood Vessels of the Skin and their Responses*. ³ Wright: *Brit. Journ. Dermatol.*, 1896, 7, 82. ⁴ Wolff-Eisner: *Derm. Zeit.*, 1907, 14, 512. ⁵ Widal: *Soc. Méd. des Hosp.*, February 13th, 1914. ⁶ Boycott, A. E.: *Journ. Path. and Bact.*, 1912-13, 17, 110. ⁷ Prausnitz and Küstner: *Centralbl. f. Bakt.*, 1921, 86, 160. ⁸ Walzer and Walzer: *Amer. Journ. Med. Sci.*, 1927, 659, 279. ⁹ Rosenau and M. Anderson: *Bull. Hyg. Lab. U.S. Pub. Health*, No. 50, April, 1909. ¹⁰ Memmesheimer: *Monograph Haut Esophylaxis*. ¹¹ Crocker: *Diseases of the Skin*, 1893. ¹² Menegh: *Journ. Amer. Med. Assoc.*, 1928, 90, 668.

BIOCHEMICAL INVESTIGATIONS IN ALLERGIC CONDITIONS.*

BY

H. W. BARBER, M.A., M.B., F.R.C.P.,

AND

G. H. ORIEL, M.A., M.D.

OUR communication is concerned almost entirely with certain biochemical investigations carried out by one of us (G. H. O.) in what may be termed allergic conditions. The word "allergy," in the sense originally employed by von Pirquet, has a narrower significance than others have given to it, for, as Darier points out, although including all forms of acquired hypersensitiveness, it excludes inborn hypersensitiveness or idiosyncrasy. We do not, however, consider this distinction of practical importance, or, in fact, valid, and with Doerr, Zinsser, Wells, Duke, and others, we employ the term to indicate almost all forms of hypersensitiveness.

Allergy is beyond question the most important biological and medical problem that exists or ever has existed, for it represents the pathology of the reaction of man and the lower animals to their environment—to the air they breathe, the physical agents, such as light, heat, and cold, to which they are exposed, the food they eat, and the various parasitic organisms which may invade them.

Moreover, in our opinion the discussions that have arisen as to whether or not the so-called allergic conditions, such as hay fever, asthma, and urticaria, should be termed anaphylactic reactions, or whether they differ essentially from true anaphylaxis as produced in animals experimentally, or as it may occur accidentally in man, are profitless. The important fact is that many of the

* Read in a discussion on urticaria in the Section of Dermatology of the Annual Meeting of the British Medical Association, Cardiff, 1928.