and the nervous, excitable, worrying type is especially apt to throw an excessive amount of nervous energy into whatever it does, accommodation included. Being anxious and over-intent about their lessons, they accommodate for the book nearer than it is, draw closer, therefore, and have to accommodate still further, overdo that, and get closer still. With their big power of accommodation they keep this up for a while. The so-called accessory muscles of accommodation (orbicularis, corrugator, frontalis) are reflexly put into strong action, and this helps to aggravate the ache in those regions. Later the effort becomes too great to be sustained. Convergence gives out, I think, first (letters are said to run into each other), and accommodation fails; the whole process is followed by headache and severe mental fatigue following on the expenditure of nervous energy. These children are of the type who have "night terrors." The treatment of these cases is the treatment of the child rather than its eyes. The cause of the trouble has to be explained to mother and teacher, and in a certain degree to the child, in order that its habits may be altered. The family doctor's help is obtained, and a daily dose of bromide may be of more use than a pair of spectacles. Our reputation will suffer if we limit our examination to the correction of half a dioptré of hypermetropia in an individual with 14 dioptrés of accommodation. It will be enhanced by the explanation that glasses are not necessary, and by the subsequent improvement in the child's health.

REFERENCES.

¹ N. Bishop Harman: *Infant Eyesight in Education, School Hygiene*, 1920. ² James Kerr: *Fundamentals of School Health*. ³ T. H. Whittington: *Metropolitan Asylums Board Reports*, 1922 and 1925-26.

THE KIDNEYS IN PNEUMOCOCCAL INFECTIONS.

BY

A. VICTOR NEALE, M.D., M.R.C.P.LOND.,

RESIDENT MEDICAL OFFICER, CHILDREN'S HOSPITAL, BIRMINGHAM.

CONFLICTING records appear in the literature concerning the relationship to nephritis of invasions of the body by the pneumococcus of Fraenkel.

Sir Thomas Horder¹ in a recent address remarked that it had been suggested that pneumococcal infections were responsible for a large proportion of haemorrhagic nephritis cases, but he himself was unable to confirm the statement from his own experience.

Abrahams,² discussing the results of 558 cases of typical lobar pneumonia occurring in the Aldershot Command during the war, shows that although there were 61 deaths and 116 developed complications, only 2 cases developed nephritis, giving the extremely low figure of 0.35 per cent.

Parsons,³ discussing the etiology of nephritis in childhood, believes "acute catarrhal desquamative or tubular nephritis" to be a condition occurring in varying degree as a result of acute infections, such as septicaemia or pneumonia, and being principally due to the circulating bacterial toxins, the renal lesion being largely recovered from when the cause is removed; but severe cases may pass on to a subacute form to constitute nephrosis (Volhard and Fahr). Several workers have attempted to associate the pneumococcus with lipoid nephrosis. Volhard⁴ observed the frequency of termination of the nephrosis cases by a pneumococcal peritonitis; and Stolz⁵ in one case found bacteria resembling Fraenkel's organism in the walls of the glomerular capillaries, the tubule cells, and the intertubular capillaries, and in another case in the intertubular connective tissue. He is of opinion that the coccus in the kidney causes an inflammation that is at first acute, but soon takes on a chronic form and results in injury to the tubules. It is suggested, however, that the germ may be a secondary invader. Variability in the incidence and degree of kidney lesions may be related to the particular virulence of the pneumococcal infection, or even to a special group type of the pneumococcus.

It would seem, therefore, that some authorities attach significance to pneumococcal infections as a source of acute nephritis, or as a starting point in the genesis of certain degenerative lipoid processes in the renal tissue (nephrosis). Others have less regard for this suggestion. However, I have made some investigations into records and certain cases which will be considered.

An analysis of the urinary findings in 287 cases of acute lobar pneumonia in adults has yielded the following:

No. of Cases.	Urine.
Group A: 10=3.4% ...	{ Albumin +++.
	{ Red blood cells +.
	{ Granular and tube casts +.
Group B: 32=11% ...	{ Albumin +.
Group C: 107=37% } 48% ...	{ No blood cells.
	{ No casts.
Group D: 138=47.5% ...	{ Albumin -.
	{ No blood cells.
	{ No casts.

In all the cases in Group A the urinary abnormality commenced between the third and tenth day after onset of the lobar pneumonia. In none of the above cases was there any history or clinical signs of previous renal disease.

Three other cases have been studied in which a previous history of nephritis was given, and which briefly were as follows.

Case 1.—Nephritis onset two years ago. Apparently mild. Acute lobar pneumonia then occurred and ran the usual course. A few days after the onset of the pneumonia the urine showed features of an acute exacerbation of the nephritis albuminuria ++++. Casts and blood cells +. Marked improvement in the renal condition quickly occurred after the febrile crisis.

Case 2.—Acute nephritis several weeks before, but rapid improvement under treatment. Later, onset of acute lobar pneumonia, which ran a normal course, ending by crisis.

Urinary Findings.

	Albumin.	Casts.	Blood Cells.
4 days after onset of pneumonia ...	++	+	+
6 " " " (crisis) ...	+	+	+
8 " " " ...	+	-	-
15 " " " ...	+	-	-

Case 3.—Known to have chronic nephritis. Urine before onset of pneumonia: albumin +; casts few; blood cells nil. Patient had an attack of acute lobar pneumonia, with usual course and crisis.

Urinary Findings.

	Albumin.	Casts.	Blood Cells.
4 days after onset of pneumonia ...	+++	++	++
6 " " " (crisis) ...	+++	++	+
10 " " " ...	++	+	-
20 " " " ...	+	+	-

The morbid anatomical conditions of the kidneys in 42 fatal cases of acute lobar pneumonia were investigated. In all the cases no previous renal disease had been suspected or detected.

- Normal in 17 cases (40.5 per cent.).
- Normal, but signs of arterio-sclerosis, in 3 cases (7.1 per cent.).
- Cloudy swelling in 17 cases (40.5 per cent.).
- Acute inflammatory changes in renal parenchyma in 3 cases (7.1 per cent.).
- Chronic nephritis with superadded recent acute inflammatory changes in one case.
- Small cortical haemorrhages in one case.

It is the aim of the clinician to attempt to correlate a particular clinical feature with a particular anatomical lesion. But in the kidney extreme difficulties have presented themselves, and to-day many exceptions are found to certain views previously held. This is to be anticipated where such an anatomically complex tissue is concerned, and where probably the different elements, although having in themselves particular functions, cannot be considered as physiological entities, but only as units in cyclic action. Osler⁶ suggests that glomerular damage is the chief feature in "pneumococcal" nephritis, and Muir⁷ agrees.

An attempt to correlate the urinary contents with the probable renal damage may be made. The meaning of albuminuria occurring in a febrile state needs explanation.

It is unlikely that undamaged renal tissue would allow such an important plasma constituent to escape and be lost in the urine. Parsons³ considers "febrile albuminuria" indicative of the mildest form of renal inflammation—a form in which clinically there is no oedema, but which shows changes in the tubule cells on morbid anatomical examination.

A comparison between the urinary findings and the morbid anatomical findings may be seen in the following table:

	Urinary Findings.	Necroses.
Nephritis		7.1%
Albumin +, casts +, blood +	4.5%	
Cloudy swelling		40.5%
Albumin +, casts -, blood -	48.0%	
Normal		47.6%
Albumin -, deposit -	47.5%	

This table shows a comparison between renal and urinary findings in two series of cases. The number of cases showing normal findings at necropsy and normal urine findings is practically the same. Similarly, the acute nephritis figures closely approximate. The remaining large group, in which albumin, in variable degree, is the only abnormal feature in the urine, agrees closely with the number showing simple cloudy swelling ("parenchymatous degeneration" of MacCallum). The significance of this change in the cells is regarded as one of the earliest indications of a disintegrative effect. There seems no doubt, however, that such cells can resume a normal anatomical and physiological state. Authorities state that acute nephritis may occur at any stage of the disease in acute lobar pneumonia. In four cases occurring in the present series the condition was detected on the fourth, fifth, sixth, and eleventh days of the illness. In each case a simple albuminuria appeared before the presence of casts and blood cells. No relation to the length of the febrile period or other single clinical feature was found. In each instance the abnormal urinary findings only lasted for a maximum period of one week, and quick recovery appeared to be the rule. In the convalescent stage each patient conformed to the normal in response to renal function tests. Pneumococcal infection would appear to act adversely on kidneys which have suffered from a previous attack of nephritis, but even in these instances the exacerbation is probably only a temporary matter, and return to a satisfactory condition may be anticipated in most cases.

Investigation of 102 cases of pneumococcal disease other than lobar pneumonia showed:

Normal	46 = 45%
Albuminuria alone	45 = 44%
Albumin, casts, and blood cells	11 = 10%

Morbid anatomical findings in a similar series (65 cases) showed:

Normal	25 = 38.5%
Cloudy swelling of parenchyma	31 = 47.0%
Pyæmic abscess	1
Nephritic changes	8 = 10%

Comparison of these results with those obtained in cases of pure lobar pneumonia shows that the readings are similar. No greater incidence of renal involvement is indicated; the degree and duration are apparently the same, and an early recovery of any renal lesion also occurs. It would also appear, therefore, that the particular focus or even extent of pneumococcal infection bears no relation to any consequent renal damage.

A remarkable constancy in the low incidence and the benign clinical features, *qua* renal symptoms and signs, was observed. In no case were any detectable cedema or uræmic symptoms found. Twenty-one cases of clinically pneumococcal infection in children under 7 years of age showed:

Normal	5 = 23.7%
Simple albuminuria	12 = 57.2%
Albuminuria with casts and blood cells	4 = 19.0%

This small series suggests that a somewhat higher incidence of renal involvement occurs in the young patient. Clinically, however, as in the adult, a minority of signs occur and quick recovery is usual.

On clinical grounds the pneumococcus would appear to be a benign organism in relation to the etiology of nephritis. Immediate and ultimate prognosis of renal damage in infections due to Fraenkel's organism would appear to warrant optimism. That profound toxæmia occurs in many cases of pneumococcal disease is not doubted, but the views here suggested are in accordance with a low renal susceptibility thereto.

Conclusions.

1. In acute lobar pneumonia nephritis is uncommon, but mild recoverable degenerative changes in the renal parenchyma are not uncommon.
2. Nephritis occurring in lobar pneumonia is of a mild type, of short duration, and apparently recovery is the rule.
3. Similar conditions occur in other pneumococcal infections.
4. The benign features of renal involvement in pneumococcal infection are similar in children and adults.

REFERENCES.

¹ Horder: *British Medical Journal*, June 4th, 1927, p. 993. ² Abrahams: *Lancet*, 1920. ³ Parsons: *British Medical Journal*, August 28th, 1926, p. 367. ⁴ Volhard, Berlin, 1918. ⁵ Stolz: *Med. Klin.*, 1922. ⁶ Osler: *System of Medicine*, 1926. ⁷ Muir: *Pathology*, 1926.

EXTREME EMPHYSEMA INVOLVING THE GREATER PART OF THE BODY:

REPORT OF TWO CASES.

BY

JOHN A. C. MACEWEN, M.B., C.M., F.R.F.P.S.,
SURGEON, ROYAL INFIRMARY, GLASGOW.

WHILE surgical emphysema is fairly often met with in connexion with fracture of the ribs, it is, so far as my experience goes, generally fairly localized in a small area surrounding the site of injury. The only case seen by me which was comparable in extent and severity to those now reported occurred in Sir William Macewen's wards in the Western Infirmary, where, in addition to the chest, the head and neck were involved, and the emphysema extended down over the abdomen to the scrotum, which was greatly distended, resembling a large hydrocele.

The cases now reported occurred at about two months' interval, the second being admitted before the first was dismissed. Both men were critically ill on admission and for a considerable period thereafter, rendering thorough examination, even by x rays, impossible.

The first, a miner, aged 44, was crushed by a fall of stone in the pit, and complained of great pain on the right side of the chest on admission, and inability to move the right arm, probably due to fracture of the right clavicle, with injury to the brachial plexus. He had a curious audible crepitus on respiration, which appeared to originate over the sternum. Emphysema had already appeared over the right chest, and on applying strapping the emphysema spread rapidly over the right neck and face, the eye being completely closed. Soon the whole face, neck, chest, and abdomen were involved, the tissues becoming greatly distended, and the skin tense and glossy. The patient was quite unrecognizable. The scrotum was ballooned up. He had great difficulty and pain in breathing; the pulse was good, but the temperature was subnormal. In spite of incisions made to allow of escape of the air, the emphysema extended, and next day the arms, and the legs down to the ankles, had become involved. Breathing was now laborious, and the patient was cyanosed; the temperature was elevated and the pulse rapid and soft.

After some days the emphysema began to subside very gradually, so that, three weeks later, it was very markedly less, the face now being free, although the thighs were still affected.

Although in this case fracture of the ribs was diagnosed clinically from the emphysema over them and the curious crepitant sound, no evidence thereof was found on x-ray examination, the fracture of the clavicle being the only definite x-ray finding.

The second case, a man of 47, a forester, fell from a tree, and was admitted in a collapsed condition, with an obvious fracture of the right humerus about the insertion of the deltoid, a dislocation of the same shoulder, which was entirely masked by the emphysema which had already occurred, extending from the fourth rib (which was fractured, together with the second and third ribs), to the angle of the jaw, and over the shoulder. A crepitant sound was also heard in this case, appearing to arise from the sternal region. About three hours after admission the