stress on these points at a discussion held at the Royal Society of Medicine last year.

To fail to distinguish between a distended ileum, a pelvic caecum, and a dilated rectum is a radiological crime of the first magnitude. The clinical significance is in each case so different that the matter cannot be left in doubt. A caecum in its normal place can nearly always be seen separately from a distended ileum by manipulation in the oblique position; where the caecum is pelvic the matter presents difficulties. But a picture which is confusing at six or eight hours is often quite clear somewhat later. A pelvic caecum is fairly common; a distended ileum rare. In the few cases in which doubt remains after the meal examinations a full opaque enema will usually clear the matter up. A simple manoeuvre serves to differentiate the rectum; a small injection, or even a glycerin suppository, will, if the mass is rectal, cause its partial or total disappearance.

As to the x-ray diagnosis of adhesions, manipulation under the screen is very unreliable. Often a picture taken a few hours later will show that an apparently immovable portion of gut has in fact moved considerably. Suspected adhesions about the colon can, however, often be put to the test by an opaque enema. A pelvic caecum is in itself of little importance, but one which does not rise on injection should always be regarded seriously, especially if it contains dense residue twenty-four hours after the main meal has passed on. Such a caecum usually has behind it an unhealthy appendix, which may give no localized signs or symptoms, and yet be responsible for gastric or duodenal disturbance. Hence the undesirability of x-ray examination confined to the upper part of the alimentary tract.

The demonstration of a genuine pelvic caecum, with a pool at twenty-four hours, and a probably infected appendix, can be accomplished only by x rays. A chronic appendix in the ordinary position could no doubt always be detected clinically, but, in fact, is often overlooked. The reason is, I think, that pain in such cases is very sharply localized, and the radiologist, palpating a visualized organ, is unlikely to overlook what may easily be missed when the hand is unaided by sight. I have heard it said that heavy x-ray gloves will cause tenderness almost anywhere if used for palpation. Apart from the fact that strictly localized tenderness is alone of diagnostic value, the statement simply is not true. I have palpated with x-ray gloves thousands of patients in the past eighteen years, and I can certify that only a small percentage complained of pain in the right iliac fossa, and in a still smaller percentage was tenderness localized in the appendix. It is, of course, a fallacy to declare an appendix necessarily free from infection because it cannot be incriminated by radiological methods.

Mr. Paterson's last criticism is of the value of x-ray evidence in demonstrating the healing of a gastric ulcer. The facts are simple. The ulcer makes a wide-mouthed pocket which fills with barium. With proper technique this pocket can be shown at will. If a time comes when the same process several times repeated fails to show the pocket, its mouth must be closed. And when, a year or two later, the symptoms return, the demonstration is complete, for the pocket is found again to be present. Here I am, for once, prepared to be dogmatic, and to say that seeing is believing.

In conclusion, I cannot allow to pass without adverse comment Mr. Paterson's dictum that if the x-ray findings do not support the clinical signs and symptoms they should be disregarded. This I consider to be a most dangerous statement, not, perhaps, as Mr. Paterson means it, but certainly as it will be interpreted by many who read his remarks. For example, if the history and symptoms call attention to the digestive system, and the radiological examination reveals changes in the lungs suggestive of tuberculosis, are these findings to be disregarded? This is no fanciful example, but has happened several times in my own practice; the course of events subsequently proving that the dyspepsia was secondary to the lung trouble. To say that there is no such thing as a purely radiological diagnosis is to play with words. No physician practising radiology makes his report with-

out taking into consideration the patient's history and symptoms. History and symptoms *plus* x-ray findings often leave no practical doubt as to the patient's complaint; just as history and symptoms *plus* physical examination may in other cases be virtually conclusive.

So far my argument has been, in the language of another profession, that there is no case to go to the jury. But I am pleased to be able to close on a note of agreement with Mr. Paterson. I do believe in co-operation between surgeon and radiologist. In particular I think it essential that both should obtain the history of the case. A patient will often tell quite a different story to separate catechisms, and it is important that these histories should be correlated.

ERRORS IN THE INTERPRETATION OF RADIOGRAMS OF THE CHEST:

THEIR CORRECTION BY TELE-RADIOGRAPHY.

BY

FF. ROBERTS, M.D., M.R.C.P.,

ASSISTANT PHYSICIAN (IN CHARGE OF THE X-RAY DEPARTMENT),
ADDENBROOKE'S HOSPITAL, CAMBRIDGE.

(With Special Plate.)

In the ordinary method of radiography of large and deep parts of the body such as the chest, when the x-ray tube is situated at 2 ft. or less from the film, considerable errors are introduced owing to the divergence of the rays. Though these errors are known to exist, their magnitude and importance are, in this country at any rate, not sufficiently recognized. It is the purpose of this article to show how misleading such errors may be, and how by the simple method of tele-radiography they can be reduced to a minimum.

The errors are of two kinds: (1) errors of position, (2) errors of size; in addition there is (3) faulty definition.

1. ERRORS OF POSITION.

Fig. 1 illustrates the projection upon the film of the ordinary postero-anterior view of the chest, the central

FIG. 1.—Postero-anterior projection through the upper part of the lung, showing displacement. See text.

ray passing through the mid-line of the body at the level of the third costo-sternal articulation. A and B are two points lying in the same horizontal plane, on the anterior and posterior surfaces of the lung respectively. As projected upon the film (A'B') A appears to be higher than B. Moreover, a point, C, lying at a lower level than A and B, is projected above A but below B. Fig. 1 (plate) shows the postero-anterior view of the chest, the distance between the tube and the film being about 2 ft. On the chest four coins of equal size have been placed. Coin A lies on the anterior wall just below the inner end of the clavicle; B lies on the posterior wall opposite the fifth rib in exactly the same horizontal plane as A. The central ray is at the level of the third costo-sternal articulation. It will be seen that B appears to be supraclavicular, whereas its true position is well below the apex of the lower lobe. The two coins C and D are placed on the anterior and posterior walls of the chest respectively in the same horizontal plane (level of the third costo-sternal articulation), through which the central ray passes. They are both $3\frac{1}{2}$ in.

from the mid-line. It will be seen that 'D' is projected well to the outer side of C.

This displacement is very little realized, and is of the greatest importance in the correlation of the radiological with the clinical findings. In a radiogram taken in this way the part of the lung field which is projected above the clavicle is not the true supraclavicular part; in other words, it is not the view which the observer would get were the patient standing in front of him with a transparent chest wall. In viewing the supraclavicular part the observer is looking into a section of lung shaped like a truncated cone, the narrower posterior end of which points downwards and towards the middle line so as to include the upper part of the lower lobe. The plane of the lung, which is just below the clavicle, is projected partly above and partly below the clavicle. The upper part of the lung is, in fact, distorted in much the same way as the roof of a house in an architect's elevation. Similarly when he is viewing the costo-phrenic angle he is looking not directly backwards, but upwards and towards the middle line, and when he is viewing the lateral wall of the chest he is looking slightly inwards. The radiogram as a whole is in reality the integration of an infinite number of parts, each of which is the projection of a cone of lung tissue. The farther any part is from the film and from the mid-line the greater is the distortion which it suffers. Even when he realizes this fact the observer finds it very difficult to transform the visual into an adequate mental image.

This important point can perhaps be expressed more simply in the following way. Let us suppose once more that the chest wall is transparent. In order to obtain a naked-eye view corresponding to the radiogram the observer would have to stand not in front of but behind the patient, with his eye in the position occupied by the x-ray target—that is to say, at the level of the third costo-sternal articulation and about one foot away from the posterior surface of the body. This fact was first pointed out by Cotton.¹

The worst feature of this error is that different parts of the lung are displaced unequally; the greater the distance from the central ray, the greater is the displacement. The consequence is that a small change in the position of the central ray alters the projection considerably, magnifying the error on the side from which the central ray has moved, and diminishing it on the other. If the central ray passes through the fourth instead of the third costo-sternal articulation the error will become greater in the apical region but less at the base. No two radiograms of the same chest, therefore, present the same appearance unless the central ray has the same position in both cases.

FIG. 2.—Projection through horizontal section through the chest, showing magnification.

(Fig. 2) be situated 8 in. from the film, the tube-film distance being 2 ft. The projected diameter is X'Y'. Let O be the position of the tube. It is obvious that—

$$\frac{X'Y'}{XY} = \frac{OX'}{OX} = \frac{24}{16}$$

$$X'Y' = \frac{3}{2} XY$$

and the area of the projected cavity is nine-fourths or two and a quarter times the true area.

In other words, from the clinical point of view the radiological appearance always exaggerates the case. Provided the error were always of the same magnitude it would not matter so much, since those who have to interpret the radiogram would know it and learn to make mental allowance for it. This, however, cannot be done, since the error varies with the distance from the film of the structure concerned.

3. FAULTY DEFINITION.

The blurring of the edges of the two coins on the posterior wall of the chest in Fig. 1 is very obvious. It is, of course, due to the penumbra formed by the blunt focus of the tube. No tube can be made with a point sufficiently fine to avoid it. But even if a point focus could be obtained, faulty definition would still occur, and for this reason. In the normal and abnormal structures found in the lung we are dealing with fine differences in the depth of shadows. Let the triangle A (Fig. 3) represent the varying depth (not size) of such a shadow tapering to a point. Let the shadow be projected in the form A'.

FIG. 3.—For explanation see text.

The inevitable result will be that the grading of the shadow will be, so to speak, more prolonged. In other words, relative opacity will be exaggerated and blurring will result. The image exhibits a greater gradation of shadow than the object warrants.

ORTHODIAGNAPHY AND TELE-RADIOGRAPHY.

In order to avoid the errors above described two methods are employed. These are orthodiagraphy and tele-radiography. In the former the patient stands behind a fluorescent screen the position of which is fixed. The tube can be moved laterally and vertically in such a way that the central ray can be projected through any part of the chest in a plane at right angles to the screen. Using a very small diaphragm, the operator with one hand moves the tube, while with the other he sketches point by point the outline of the organ examined. Since each point is the projection of the central ray, the outline thus obtained represents the true size of the organ. The sketch is then transferred to tracing paper. Alternatively a pencil moving automatically with the tube is used. This method, which was devised by Moritz and developed by Groedel, is now extensively employed on the Continent, particularly in Germany and Austria. It has the advantages of rapidity and cheapness, since films are not required. It has two great disadvantages: first, while it provides an admirable method of recording bold outlines such as the borders of the heart, it is impossible by this means to follow fine details; secondly, the record thus obtained is subjective and not objective.

Tele-radiography simply consists in increasing the distance between the film and the tube so that the errors due to divergence are reduced to a minimum consistent with practice. The distance usually employed is 2 metres, or about 6 ft. The rays issuing from a tube at this distance may for practical purposes be considered to be parallel. This method, which was first suggested by Kohler, is coming into increasing favour, but in my opinion has in this country not yet received the attention which it merits. One reason for this is the incidental result of the improvements which have been made in the construction of x-ray apparatus. Such improvements have, rightly enough, had for their main object the protection of the operator. Most screening stands are so constructed that the patient must stand close up to the tube. But while this arrangement is admirable both from the point of view of protection and for the purpose of a visual examination, it is not suitable for obtaining a true picture. The old objection that a long exposure is necessary is in these days of powerful plant no longer valid.

My own procedure is as follows. Using a screening stand in which the tube and screen move together, a screen examination is first made. The patient comes out of the screening stand and sits on a stool fitted with a film

FIG. I.—Postero-anterior view of chest, taken at 2 ft. A, B, C, and D are four similar coins, A and C being on the front wall, B and D on the back wall. B is situated directly behind A, and D directly behind C. Note magnification and displacement of B and D. B, though opposite the upper part of the lower lobe, appears to be above the clavicle.

FIG. II.—The same, but at 6 ft. instead of 2 ft. Note the correction in size and position of B and D.

FIG. III.—Cavity in the left lung taken at 2 ft. The cavity is indicated by cross-lines.

FIG. IV.—The same, but at 6 ft.

FIG. V.—Cavity in its true projected size (unreduced) from FIG. III.

FIG. VI.—The same (unreduced) from FIG. IV.

holder (Fig. 4). This is a very simple piece of apparatus which was put together in a few minutes. It consists of an ordinary office stool 2 ft. 6 in. in height, above one side of which projects a vertical board 2 ft. in height and 1 ft. in breadth, supported on a pillar which is screwed to the stool. The lower edge of the board is 9 in. above the seat. The patient sits facing the board with his legs protruding beneath it, one leg on each side of the pillar. The film in its cassette having been placed upon the board, the patient presses his chest against it with his arms folded behind the board. This position not only immobilizes him, but also has the incidental advantage that it causes the maximum abduction of the scapulae, with, in consequence, the least possible obstruction of the lung fields by these bones.

FIG. 4.—Stool and film stand used for tele-radiography.

The stool is placed so that the film is 6 ft. from the tube. The positions of the tube and stool have been previously adjusted so that the central ray passes through the mid-line of the body at the level of the third costo-sternal articulation. These positions are standardized once and for all, allowance being made for varying heights of patients. Suitable marks are made on the upright and arm of the tube-stand, while the position of the stool is marked on the floor. Tube and stool can therefore be adjusted rapidly in the correct position.

For patients who have to be examined in the recumbent position, the best method is to put the patient on a stretcher on the floor and to use a vertical stand with the tube as high as the pillar will allow. It is not necessary to turn the patient over on his face, for the following reason. The ordinary close-up antero-posterior view of the chest (the rays passing from front to back) is unsatisfactory, since the heart and great vessels lying nearer the front than the back wall of the chest cast an exaggerated shadow. With the tube at 6 ft., however, this exaggeration is reduced to a minimum. In fact, there is very little difference at 6 ft. distance between the antero-posterior and the postero-anterior view. It will therefore be seen that the method is very simple, the only trouble being involved in the initial standardization of the position of stool and tube.

Fig. 11 is the same chest as Fig. 1, but taken in this way. It will be observed that coin B is almost directly behind A and approximates closely to it in size; D, however, is slightly to the outside of C, owing to the fact that the patient is turned slightly to his left. This can readily be determined by comparing the positions of the two sterno-clavicular articulations in relation to the edges of the vertebral column. It will be seen that the inner end of the left clavicle is slightly farther out than that of the right. Had the patient been sitting quite square D would have been almost directly behind C.

Again, taking the case of a structure situated 8 in. from the film, let XY be the actual diameter of the structure and X'Y' the diameter of its shadow. We have

$$\frac{X'Y'}{XY} = \frac{OX'}{OX} = \frac{6 \text{ ft.}}{5 \text{ ft. } 4 \text{ in.}} = \frac{9}{8};$$

$$\text{while } \frac{\text{area of } X'}{\text{area of } X} = \frac{81}{64}$$

The exaggeration in area is therefore about nine-sevenths, as compared with nine-fourths in the close-up view. It will further be observed that the outlines of coins B and D, instead of being blurred as in Fig. 1, are now as well defined as those of A and C.

Space prevents my showing many instances of the various conditions in which the superiority of the tele-radiogram is manifest. I shall confine myself to one case, illustrated in Figs. III and IV. These show in the left lung an egg-shaped ring shadow, probably due to a cavity. The position and extent of the cavity are indicated by cross-lines. Fig. III was taken at a distance of 2 ft., and Fig. IV at

6 ft. It will readily be understood that in the process of reduction the difference has become considerably diminished. I have therefore reproduced in Figs. V and VI the cavity in its true projected size; Fig. V is at 2 ft., Fig. VI at 6 ft.

Enough has been said to show how misleading the ordinary radiogram of the chest may be. Owing to the error in size the film always shows the case to be more serious than it really is, while the error in position must often lead to considerable discrepancy between the radiological evidence and the evidence deduced from physical signs. It is also clear that, as records of the progress of a case—for example, the changing size of a tumour, cavity, or aneurysm—successive radiograms lose much of their value unless they are taken under the same standard conditions. It seems to me to be highly desirable that for comparative purposes a conventional standard should be adopted, the two constants being the distance of the tube from the film and the level of the central ray. This, of course, is not to say that other views are unnecessary. I should be the last to minimize the importance of special views, such as the Kreuzhohl view suggested by Fleischner for revealing interlobar effusions, but I feel sure that the adoption of a standard tele-radiogram would be a step forward in the radiology of the chest.

I have to thank Mr. Mann, radiographer at Addenbrooke's Hospital, for his skilful technical assistance.

REFERENCE.

¹ Cotton, W.: *Archives Röntgen Ray*, 1902.

AN UNUSUAL CASE OF URETERAL STONE.

BY

HERBERT WILLIAMS, M.B., F.R.C.S.,

SENIOR HONORARY SURGEON, ROYAL SOUTHERN HOSPITAL, LIVERPOOL; AND

C. THURSTAN HOLLAND, D.L., M.Ch., F.R.C.S.,

LECTURER ON RADIOLOGY, LIVERPOOL UNIVERSITY.

(With Special Plate.)

THE investigation of this case presents many features of interest both from the urological and radiographic points of view. The short history of the case is as follows.

Miss M., aged 46, in May, 1927, had a sudden pain in the left loin lasting from four to five hours, accompanied by frequency of micturition. The temperature was normal.

In October, 1927, she had very severe pain in the left loin; she was writhing in agony and could not keep still. This was associated with a rigor, sweating, and violent retching. There was no great frequency of micturition, but the urine was acid and contained a slight amount of albumin, pus cells, staphylococci, and coliform bacilli. In ten days she was well again.

In November, 1927, she had a similar attack to the last, but it was more severe and lasted a fortnight.

Between the attacks she had persistent frequency of micturition and sharp pain in the urethra after the act, persisting for varying times afterwards.

In December, 1927, an x-ray examination was made elsewhere. Various shadows were found in the pelvic region, and it was also suggested that there might be a stone in the pelvis of the left kidney.

On February 2nd, 1928, C. T. H. examined her and took the usual films to cover the whole stone area. These radiographs showed that there was no stone in either kidney, and both kidneys appeared to be normal in size and position. In the pelvic area were shown the same shadows which had previously been found, and all of them were in exactly the same position. (See radiograph, Fig. 1.) This showed that no stone had passed in the interval from the left kidney. It was obvious that the only shadow to be considered was the rounded one on the left side of the pelvis. This was not in the usual line of the ureter, it was round in shape, and it was in exactly the same position in two radiographs taken at intervals of a few weeks. C. T. H. very definitely decided against it being a shadow of a ureteral stone.

Now came the first piece of luck. C. T. H. was out when the surgeon in charge of the case (H. W.) came in to see