- De Haen, A.: *Theses sistentes febrium divisiones*, 1760, p. 23.
Hirsch, A.: *A Handbook of Geographical and Historical Pathology*, New Sydenham Society, 1883, i, p. 171.
Huxham, J.: *Dissertation on the Malignant Ulcerous Sore Throat*, third edition, 1759.
Ingrassias, J. P.: *De tumoribus praeter naturam*, Tom. I, cap. i, p. 194, Naples, 1553.
Lützhoft, F.: *Bibliothek for Laeger*, 1919, cxi, 479.
Malfatti, J.: *Neues Journal der praktischen Heilkunde*, 1801, v, p. 120.
Morton, J.: *Pyretologia*, cap. v, Geneva, 1696.
Noirot, M. L.: *Histoire de la scarlatine*, 1847.
Parsons, A. C.: Some administrative aspects of scarlet fever, Ministry of Health, *Reports on Public Health and Medical Subjects*, No. 35, 1927.
Payne, J. F.: *Masters of Medicine: Thomas Sydenham*, 1900.
Piorry, P. A., et Lheritier, D.: *Traité des altérations du sang*, 1840.
Rayger: *Observationum medicinalium centuria*, 1691.
Richter, R.: *Arch. f. Gesch. d. Med.*, 1908, i, p. 161.
von Rosenstein, R.: *Diseases of Children*, English translation by Spearman, 1776.
Sanné, A.: *Dictionnaire encyclopédique des sciences médicales*, 1879, art. scarlatine.
Sennert, D.: *Epist. med.*, Cent. II, Ep. 18.
Trousseau, A.: *Clinique médicale de l'Hôtel-Dieu de Paris*, 7e éd., 1885, i, pp. 148, et seq.
Welch, W. W., and Schamberg, J. F.: *Acute Contagious Diseases*, 1905.
Willan, R.: *On Cutaneous Diseases*, 1808.
Withering, W.: *An account of the sore throat or scarlatina anginosa, particularly as it appeared at Birmingham in 1778, 1793.*

THE LIGHT SENSE IN MINERS' NYSTAGMUS.*

BY

R. J. COULTER, M.B., B.CH., F.R.C.S.I.,
Ophthalmic Surgeon, Royal Gwent Hospital.

I AM fully aware that this communication is incomplete and inconclusive, but I have some hopes that it will open up a fresh aspect of the subject with which it deals, and induce others to investigate the changes which occur in the light sense in miners' nystagmus, and possibly find in it some help in elucidating the mysteries of that most complicated condition.

My attention was first directed to the subject by a paper read by Mr. Percival at the Oxford Congress,¹ in which he stated that in all cases of miners' nystagmus which he had examined he had invariably found the light minimum greatly increased, while their light difference was very little if at all greater than his own. At the same meeting he demonstrated the simple apparatus which he used in making the examination; it consists of black-and-white discs with sectors of white and black of varying sizes on them, which, when rotated on pins, show grey circles of varying intensity. My tests have been made with this apparatus, but recently I have checked the light difference with Young's discs, and find that it is easier to work with them and that the results are similar, so far as I can judge. Unfortunately Young's portfolio does not contain a light minimum test.

The accompanying table is founded on the records of

The Light Sense in 100 Cases of Miners' Nystagmus.

LIGHT MINIMUM.				LIGHT DIFFERENCE.			
Total.	No Miners' Nystagmus.	Miners' Nystagmus or Neurosis.		Total.	No Miners' Nystagmus.	Miners' Nystagmus or Neurosis.	
200	8	4	29	100	51	26	25
100	51	25					
50	31	12	19	50	35	14	21
25	7	3		25	11	5	
12.5	3	1	2	12.5	3	0	3
	100	45	55		100	45	55

Light Minimum.—Normal or practically normal=59 per cent.; reduced=41 per cent.

Light Difference.—Normal=51 per cent.; reduced=49 per cent.

100 cases, and the first thing that strikes one is that my results are exactly the reverse of Percival's, as they show reduction in both the light minimum and the light difference in about 50 per cent. of the cases. This naturally puzzled me, and on thinking the matter over I came to

the conclusion that Percival's observations must have been made on fresh cases, while mine were made on old ones, which were sent to me for examination with a view to ascertaining whether they had recovered. I wrote to Mr. Percival, and he kindly replied, stating that this suggestion was correct, and that he agreed that both light senses are reduced in old cases. I have not been able to verify the observations on fresh cases, but if it is true that they have an increased light minimum which later turns into a deficiency it is a remarkable phenomenon. Percival attributes the early increase to an increase of endogenous stimulus, and considers that this also accounts for the intolerance of light, but this latter may persist after the light minimum has become reduced. It would seem that either the increased endogenous stimulus must be sufficient to overcome the reduction in light sense which becomes manifest after it has passed off or that the reduction is something fresh which develops after the men have ceased working underground.

Coming to my cases, and adopting Percival's nomenclature and standard, only 8 per cent. had a normal light minimum, but 51 per cent. had the next grade labelled as 100, while in the remaining 41 per cent. the light minimum was definitely reduced. I have divided these cases into two groups according to whether I considered from their other signs and symptoms that they were still suffering from miners' nystagmus, or neurosis arising out of it, or had recovered. Of course this is only an opinion to be taken for what it is worth, but it indicates that, although 50 per cent. of the cases with full light minimum still showed evidence of the disease, and one of those with the lowest grade measurable by the test was free from symptoms, there was a definite increase in the proportion of those still suffering from miners' nystagmus to those who had recovered as the light minimum diminished.

Dealing with light difference I found a similar condition. It was normal in 51 per cent. and subnormal in 49 per cent., while the proportion of cases with signs or symptoms of miners' nystagmus to those without them tended to increase more definitely than in the case of light minimum.

I am aware how inadequate these records are, both in numbers and in the manner in which they have been worked up, but in spite of this I suggest that possibly there are some lessons to be learnt from them. The first is, unfortunately, the negative one, that I have not found any evidence that examination of the light sense will help us in definitely deciding whether a man has recovered from miners' nystagmus. Another arises from a consideration of the possible causes of the symptom. At first sight it would seem to be obviously a part of the neurosis which is the true cause of most of the incapacity arising out of the disease, but it is possible that this may not be so in all, or even in the majority, of the cases. During the war Ransome Pickard and G. W. Lloyd² found that a number of men who were invalided suffered from a mild form of scurvy with night-blindness, due to deficiency of vitamin from insufficient supply of vegetables, and had symptoms somewhat resembling those of miners' nystagmus, and Barton states that the blood of men suffering from the disease is deficient in haemoglobin, the average being 75 per cent. of normal. It may seem a far-fetched theory that some of the prolonged cases of miners' nystagmus are really suffering from vitamin deficiency, but these men cannot be in a position to afford expensive food, and it is possible that they may be using margarine, which is notoriously deficient in vitamin D. I throw out the suggestion as indicating a line on which investigations might be carried out with regard to the general condition of the sufferers from this most serious disease, on the lines suggested by Mr. Freeland Fergus, with some possibility of doing something for the patient. I cannot refrain from expressing my opinion that in spite of all the discussions which have taken place on miners' nystagmus, and all the money which has been spent on compensation and deciding whether individuals are entitled to it or not, the poor patient has been most shamefully neglected, and very little attention has been paid to his general condition or to the measures which ought to be adopted to improve it.

REFERENCES.

- ¹ *Trans. Ophth. Soc.*, v, xl, p. 323. ² *Proc. Roy. Soc. Med.*, 1920 (Section of Medicine), p. 43.

* A paper read in the Section of Ophthalmology at the Annual Meeting of the British Medical Association, Cardiff, 1928.