vein. The present specimen, I believe, shows this point. I cannot help thinking but that the coiled up clot in the heart came from a systemic vein, yet one finds evidence of quite definite *antemortem* clotting in branches of the pulmonary arteries.

Discussion.

The features of this case are:

1. The lack of proper massage and movement, thus favouring stasis of the blood, which many workers have shown tends towards thrombosis.¹
2. The lack of evidence of thrombo-phlebitis before the embolism occurred.
3. The length of time which the patient lived, presumably with this large foreign body thrashing about in the heart, the only symptoms being the sickness and the vague feeling of something wrong.
4. The remarkable size of the clot.

REFERENCE.

- ¹ Lister, W. A.: *Lancet*, 1927, i, 111.

SOME OBSERVATIONS ON THE ERYTHROCYTE.

WITH SPECIAL REFERENCE TO PUNCTATE BASOPHILIA, DIFFUSE POLYCHROMASIA, AND RETICULATION.

BY

W. E. COOKE, M.D., F.R.C.P. Ed., M.R.C.P.
(From the Pathological Department, Wigan Infirmary.)

FROM the time that Ehrlich (1885) noted the three distinct basophilic conditions in erythrocytes controversies have waged on their character and significance; but for many years the trend of opinion has been towards the view that punctate basophilia, diffuse polychromasia, and reticulation are expressions of the same process.

Cesaris-Demel (1907) concluded that punctate basophilia and reticulation were identical, and Key (1921) agreed that the basophilic substance in the three conditions was the same. Brookfield (1928) believes that the stippled cells in lead poisoning are immature cells (reticulocytes and polychromatic cells) altered by the toxic action of lead. His

FIG. 1.—Erythrocytes showing spongioplasm after the lipid membrane has been altered or dissolved. ($\times 1,000$.)

results tend to confirm the work of Hawes (1909) and Key (1924), in which the close numerical relationship between stippled and polychromatic cells and reticulocytes was definitely established.

The nature of the staining substance is still undetermined, nor has any explanation been offered as to why the different appearances occur. Pappenheim (1907) and Biondi (1908) independently propounded the view that the stainable substance was a dye-precipitate upon the surface of the lipid membrane, but, later, Pappenheim (1919) rather altered this idea, and suggested that the lipid component of the spongioplasm was the reacting substance.

That the substance is not chromatinic is beyond doubt. The staining reaction with methylene blue-eosin preparations and with methyl green-pyromin, and the fact that

none of the conditions is visible by dark-ground illumination or ultra-violet light, are among the arguments against punctate basophilia, polychromasia, and reticulation being nuclear in origin.

Gruner (1913) suggested that the reticular substance was not nuclear but plasmatic in character, because in embryos and young animals its abundance was not associated with the appearances of changes in the nucleus. I have confirmed this observation in cases of Addison's anaemia during normoblastic crises, in von Jaksch's anaemia, and in myeloid leukaemia. In these diseases there does not appear to be any relationship between the younger normoblasts and the older normoblasts and the amount of reticular substance.

The present work is an attempt to elucidate these problems. The fact that haemoglobin gives a characteristic reaction with benzidine and hydrogen peroxide was utilized to the exclusion of all other stains. Blood films were made on 3×1 microscope slides and allowed to dry in the air before treatment. By altering the quantities of the reagents* it is possible to disturb the apparent homogeneity of the erythrocyte. For the better appreciation of the early effects we will take the final results first. Fig. 1 is a photomicrograph of erythrocytes from which the haemoglobin has been more or less completely discharged. Benzidine in alcohol and hydrogen peroxide were used in such strengths that when flooded on to the film a greenish precipitate was immediately produced and floated to the surface; the reaction was allowed to continue for twenty minutes. If the corpuscles in the figure are examined with a hand lens an irregular meshwork is seen, and the spaces between the strands have the appearance of pores. The outline of the cells is irregular, but the mean diameter is increased. The probable sequence of events is that the lipid membrane has been oxidized and dissolved, and a slow fixation has brought into evidence a spongy stroma through the interstices of which the reagents have penetrated. Reaction between the benzidine and hydrogen peroxide and haemoglobin has taken place with solution and escape of the end-product. Having seen the end-result, all kinds of patterns, depending upon the varying degrees of penetration of the envelope and the amount of haemoglobin

FIG. 2.—Erythrocytes showing finely granular punctate basophilia. ($\times 1,000$.)

released, would appear possible. This is, in fact, true. Every type of punctate basophilia and reticulation may be duplicated.

Fig. 2 is an example of punctate basophilia of the fine type. If the cells are examined with a lens it will be apparent that the reaction has not progressed to the stage of Fig. 1. There are areas in which the granules are sharply defined; in others, however, they are blurred in outline, suggesting that the haemoglobin has been reached but different stages of the reaction have taken place. The granules may be fine as in this figure, or coarse as in some of the cells in Fig. 4, or they may be irregular in shape and aggregated into small clusters as in Fig. 3.

At a further stage the haemoglobin has escaped through

* The reagents are benzidine 0.01 per cent. in 80 per cent. alcohol, and hydrogen peroxide.

the pores and bridged over intervening strands of the stroma. Areas have coalesced in an irregular manner and the appearance of reticulation results, as shown in some of the cells in Fig. 4.

All the forms of reticulation described by Brookfield (1928), from the unfragmented reticulum to the bar form, may be reproduced. Fig. 5 is an example of the bar form. The reticulocytes in Fig. 4 suggest that the next stage

results strengthen the belief that the stainable substance is not nuclear in character, but is probably a haemoglobin compound, and that the three basophilic conditions of the erythrocyte are qualitatively the same and vary only in degree. The second deduction is that the conditions, when obtained by the usual staining methods, would point to increased permeability or a defect in the lipid envelope of the erythrocyte rather than to youthfulness, although

FIG. 3.—The clumped form of punctate basophilia showing granules of irregular shape. ($\times 1,000$.)

FIG. 4.—Punctate basophilia, reticulation, and diffuse polychromasia (x). ($\times 1,000$.)

FIG. 5.—The bar form of reticulation. ($\times 1,000$.)

must be diffuse polychromasia. The cell marked X in Fig. 4 illustrates the condition up to a point, but there is a rim of unaffected material around the central stained area. This is probably due to the time given for the reaction being too short.

Comments.

This preliminary note is merely intended to place on record the fact that punctate basophilia, diffuse polychromasia, and reticulation may be produced in any erythrocyte. The time is not opportune to enter into the discussion which these results open up, but two deductions may, I think, be made. The first, that with our previous knowledge of the staining reactions the above

there may be, and probably is, a close relationship between punctate basophilia, diffuse polychromasia and reticulation, and immaturity.

I take this opportunity of expressing my thanks to my friend C. F. Hill for his invaluable assistance with the photomicrographs.

REFERENCES.

- Biondi, C.: *Fol. Haematol.*, 1908, vi, 205.
- Brookfield, R. W.: *Journ. Path. and Bact.*, 1928, xxxi, 2, 227.
- Cesaris-Demel, A.: *Fol. Haematol.*, 1907, iv, Suppl., 1.
- Ehrlich, P.: *Charite-Annalen*, 1885, x, 136.
- Gruner, O.: *Biology of the Blood Cells*, 1913, 75.
- Hawes, J. B.: *Boston Med. and Surg. Journ.*, 1909, clxi, 493.
- Key, J. A.: *Arch. Int. Med.*, 1921, xxviii, 511.
- Idem: *Amer. Journ. Physiol.*, 1924, lxx, 86.
- Pappenheim, A.: *Fol. Haematol.*, 1907, iv, Suppl., 46.
- Idem: *Ibid.*, 1919, xxiv, 1.

BILATERAL OBSTRUCTION OF THE CENTRAL RETINAL ARTERIES.*

BY

FRANK JULER, M.A., M.B., F.R.C.S.,

Surgeon, Royal London Ophthalmic Hospital; Assistant Ophthalmic Surgeon, St. Mary's Hospital.

My excuse for drawing attention to this subject is that Dr. T. G. Wynne Parry published a report of a similar case in the *British Medical Journal* last February, and commented upon the rarity of the condition. As a matter of fact, a number of bilateral cases have been reported, a few of simultaneous onset, a larger number with an interval between the incidence of the condition in the two eyes. The notes of my patient are as follows:

A man, aged 50, came to Moorfields on November 20th, 1920, having lost the sight of the left eye suddenly two days before. The vision was reduced to bare perception of light, white opacity of the retina at the posterior pole of the fundus was already present, and the macula showed the typical cherry red spot. The arteries were narrow. Two weeks later the white opacity of the retina had disappeared. The right eye with spherical—2D had 6/9 vision and its retinal arteries showed slight sclerosis. Dr. Gordon Holmes reported that the blood pressure was high, 260/156, and that chronic renal disease was present.

On April 5th, 1922, he came again, having lost the sight of the right eye suddenly five nights before. Its vision was reduced to finger-counting at one metre. The right disc was atrophic, the arteries were narrow with apparent obliteration in places, and the lumina of the veins were much interrupted. There was a cherry red spot at the macula, and some whitish specks in that region.

* A paper read in the Section of Ophthalmology at the Annual Meeting of the British Medical Association, Cardiff, 1928.

Cause of the Obstruction.

The work of Coats and others has made it more than probable that most of the cases occurring in elderly folk are due to endarteritis or thrombosis in the central artery itself, and that an embolus is unusual. Embolism undoubtedly does occur, and is probably the correct explanation of the condition when it is present in young adults who are the subject of mitral stenosis, congenital heart lesions, etc. Spasm of arteries, haemorrhage into the nerve sheath, and pressure on the nerve itself are other possible, but not so well authenticated, causes. In a number of the cases occurring in young folk no cause can be detected from a general examination of the subject.

Obstruction Occurring during Sleep.

Frequently the condition of blocked central artery or vein has been noticed first on waking from sleep, and it has been concluded that, in the arterial as in the venous cases, this was suggestive of a thrombosis. In cases occurring in elderly persons with arterial disease this is likely to be the explanation, but it is possible for embolism also to occur during sleep.

MacWilliam has recently drawn attention to the not infrequent occurrence of cerebral haemorrhage, anginal attacks, and death during sleep, and he has explained this by his experimental observations on the blood pressure in this state. He finds that there are two entirely different conditions in question in sleep: (1) sound sleep, with lowering of blood pressure, (2) disturbed sleep, dreaming, etc., which may be attended with remarkable elevations of pressure.

In sound sleep, lowering of the systolic blood pressure to the extent of 15 to 30 mm. at the end of two hours