GANGLIONECTOMY IN THE TREATMENT OF SEVERE POLYARTHRITIS AND OSTEO-ARTHRITIS

BY

ARCHIBALD YOUNG, B.Sc. ; M.B., C.M. ; F.R.F.P.S.G. ; F.A.C.S. (HON.) ;

DOCT. HON. CAUSA (STRASBOURG)

REGIUS PROFESSOR OF SURGERY IN THE UNIVERSITY OF GLASGOW

It is generally agreed that the severer forms of arthritis, whether of the primary rheumatoid type or of the nature of osteo-arthritis, primary or secondary, present a prognostic and therapeutic problem often of the utmost difficulty. In a certain proportion of cases the disease proves intractable, no matter what line of treatment is adopted, and the condition progresses relentlessly to a stage of extreme disablement.

The exact percentage of intractable cases would be assessed variously by different authorities. A recent writer, Dr. C. W. Buckley,¹ taking a more hopeful view than most, would place it at something less than 10 per cent. ; my own opinion is that the percentage is considerably higher. In any case, as Buckley observes: "To the lay mind the prospect before the patient attacked by arthritis of any kind is one of pain, crippling, and deformity, with little or no prospect of relief." And though he goes on to state his belief that such a gloomy prognosis is not justified, provided suitable treatment is instituted early and continued perseveringly, I am bound to say that his comparatively optimistic view is not in line with such experience as I have had, nor in keeping with the views of many of my colleagues.

In a discussion at a meeting of the Royal Medico-Chirurgical Society of Glasgow in 1933² I ventured to suggest that, in view of their somewhat gloomy therapeutic outlook in many of these cases, the physicians might well turn more readily and more frequently towards their colleagues, the surgeons, who could offer them and their patients, in at least a certain proportion of otherwise intractable cases of chronic polyarthritis, something more than a mere visionary hope of relief. I described then certain personal experiences I had had within the immediately preceding years in two cases of distressing and seriously disabling chronic rheumatoid polyarthritis of hip, knee, and ankle, which had been relieved almost beyond belief by the operation of ganglionectomy.

One patient was a young woman who had been bedridden for almost two years before her operation, and who after surgical treatment (lumbar ganglionectomy) was able to walk out of the ward without assistance. The other case was that of a middle-aged man whose lower limbs in particular were profoundly affected ; he suffered intense pain on the slightest attempt at movements of his joints, and non-surgical treatment had entirely failed to provide any relief. A lumbar ganglionectomy operation gave him immediate relief from pain, and he soon recovered a wide range of movement.

I claimed, therefore, that in a certain proportion of cases of chronic arthritis surgery could offer something substantial in the way of alleviation, and even of cure, which was worthy of the consideration of the physicians. Such a claim I am still prepared to maintain.

Recorded Results of Operative Treatment

At the recent International Surgical Congress in Cairo a discussion took place on the subject of lumbar sympathectomy. As British rapporteur I had the honour of making one of the opening contributions, in which I dealt with the various more important applications of the procedure.

After reviewing many of the conditions for which lumbar ganglionectomy has been more generally employed, such as Raynaud's disease, thrombo-angiitis obliterans, arteriosclerosis, and combinations of these, I went on to discuss its application in the treatment of certain painful and disabling chronic diseases of the joints, such as periarticular rheumatism, rheumatoid arthritis, and even certain cases of osteo-arthritis. The results achieved in certain cases of profoundly disabling polyarthritis I characterized as among the most striking illustrations of the value of the procedure. To quote from the summary of my paper:³

"The most striking result of all was achieved in six cases of profoundly disabling polyarthritis, of a rheumatoid type, the completely bedridden and crippled patients being relieved from pain and restored to the possibility of a useful life."

In an earlier paper, delivered to the Royal Medico-Chirurgical Society of Glasgow in 1931,⁴ I had drawn attention, not merely to the value in certain doubtful joint conditions of the procedure of periarterial sympathectomy—and cited illustrative cases—but also to the possible efficacy of ganglionectomy in the treatment of certain painful and chronic joint conditions. Up to that time (1931) I had myself had no personal experience of ganglionectomy in the treatment of these serious joint conditions, but I pointed out that Adson and his co-workers had already reported a number of cases in which a substantial degree of benefit had been achieved.

Adson's first case was reported in 1928 in the *Proceedings of the Staff Meetings of the Mayo Clinic*⁵ amongst other places.

A young woman was presented at a staff meeting on November 21st of that year by Messrs. Rowntree and Adson. She had consulted Dr. Rowntree originally by reason of a crippling polyarthritis, particularly of the joints of the lower limbs, and, more as an experiment than anything else, a bilateral lumbar ganglionectomy was performed in June, 1926. Careful studies had been made previously of her vasomotor reactions, and she was found to have a high vasomotor index. It was thought, therefore, that the ganglionectomy procedure, by inducing a favourable vaso-dilatation, might beneficially affect the condition of the disabled joints. The operation was followed by a result so favourable that Rowntree characterized it at the time as "in many ways the most remarkable result I have seen in my experience in medicine, remarkable not only from the standpoint of the patient's

relief, but from the standpoint of the innumerable questions suggested to us as physicians."

The patient had an uneventful convalescence. There was at once a very definite improvement, as well as definite relief from pain, and increase in the surface temperature of the feet and ankles. Three weeks later a bilateral operation for union was performed successfully. The patient was soon able to walk again without the slightest difficulty, and the toes, which had been previously deformed and distorted, could be extended forward in a manner previously altogether impossible.

In view of this favourable result other cases were submitted to the same procedure, and in 1929 Messrs. Rowntree and Adson were able to report their sixth case.⁶

This patient, also a young woman, was perhaps not so badly crippled as the first one, but almost all her joints were to some extent involved except the hips. X-ray examination showed periarticular arthritis of the elbows, wrists, ankles, and knees, with destructive arthritis and deformity of the phalangeal joints of the right hand. Medical measures had been tried perseveringly.

A bilateral lumbar ganglionectomy in June, 1929, was followed (after an initial, somewhat stormy, post-operative course) by very remarkable improvement in function, a great improvement in the warmth of the lower limbs, and the abolition of pain.

Adson and his co-workers continued to employ the operation in suitable cases of polyarthritis with generally satisfactory results. In view of their favourable experience it is somewhat remarkable how little attention seems to have been paid by other surgeons to the promise of this new line of experimental therapeutics.

Fraser, writing in 1931,⁷ made only the briefest reference to the possible value of ganglionic sympathectomy in the treatment of polyarthritis. He recorded one case in a girl aged 23, who had a pronounced polyarthritis of the upper extremities. A cervico-thoracic ganglionectomy was carried out on the left side, with a tolerably favourable result, especially as regards relief of pain; but evidently not a result so remarkable as to make Fraser particularly enthusiastic.

A few surgeons here and there have tested the operation in suitable cases, but it cannot be said that it has been put to any widespread or general test, or has met with general acceptance. Thus in the report of Mr. Paterson Ross to the Association of Surgeons of Great Britain and Ireland in May, 1935, which represented a summary of a collective inquiry made by the association into the results of sympathectomy, there was no mention at all in the tables presented to the association of any cases of polyarthritis or indeed of any form of arthritis submitted either to the periarterial procedure or to ganglionectomy. This omission seemed to me so remarkable in view of my own experience in recent years that, in the subsequent discussion on that occasion, I drew particular attention to it, and referred to several cases of my own in which I had already put the matter to the test, and in which the results had been of a highly satisfactory nature.

Note.—It should perhaps be stated here that, while the tabular summary of the collective inquiry into the results of sympathectomy as presented to the association by Mr. Paterson Ross—a copy of which was in the hands of all the Fellows present at the meeting, and a copy of which is before me as I write—contained no reference to the employment of sympathetic ganglionectomy in cases of arthritis, there is the following brief reference to it in the paper contributed by Mr. Paterson Ross to the October, 1935, number of the *British Journal of Surgery*.⁸ In this paper were presented the various tables included in the summary, with Mr. Ross's comments thereon: "*Chronic Arthritis*. Sympathetic ganglionectomy was performed in five cases of chronic arthritis, associated with excessive sweating and vasomotor disorders in the affected limb. In three patients the arm-joints were involved, and two of them derived great benefit from the operation. In the other two patients arthritis affected the lower limb, and in one

of them the treatment was successful in relieving pain and restoring function."

In the interval between the delivery in 1931 of my earlier paper already referred to and the Cairo congress of 1935 I had dealt with six cases of extreme disabling polyarthritis, involving many joints. Two of these were the cases I referred to in the opening part of this present article, and which I cited at the meeting of the Royal Medico-Chirurgical Society of Glasgow in November, 1933. The other four were additional. I reported all six in detail at the Cairo congress. As a result of my experience in this group of cases I had now definitely reached the position that I felt able to promise to these extremely crippled and disabled sufferers a degree of relief from pain, and even the prospect of an amount of functional recovery, that previously I could hardly have believed possible.

Factors Concerned

What the exact rationale of the therapeutic process may be it is not at present possible to say with any degree of certainty. The simplest explanation would seem to be that the augmentation of blood supply to the limb plays the chief part in determining the satisfactory remedial effect. Such a view is rendered all the more probable by the observation made by Rowntree, Adson, and their co-workers⁹ that the cases most likely to respond beneficially are those in which the protein shock test is followed by a favourable vasomotor or vascular index. But much further investigation is necessary before even this view can be accepted.

Other factors must of course be taken into consideration. It is quite possible, for example, that the physiological reaction which determines the beneficial therapeutic result may be, at least in part, of a nervous character. If, as some think, there is a definite trophic element concerned in the production of chronic arthritis—even in arthritis of a rheumatoid character—this may, in part at least, explain the remedial effect produced by gross interruption of the sympathetic pathway. That still another factor may require to be taken into consideration in the search for the real explanation of the therapeutic effect is suggested by the somewhat remarkable outcome in a number of my cases, in which the beneficial result of a lumbar ganglionectomy has not been confined by any means to the joints of the lower extremities, but has affected also—and sometimes to a considerable degree—the joints of the upper limbs as well.

Leriche¹⁰ drew attention long ago to the fact that in many cases where a periarterial sympathectomy was performed on one limb—say in early cases of Raynaud's disease—a corresponding therapeutic effect might be produced at the same time in the homologous limb. Further, it is well known that, even following the simple periarterial procedure, there is produced, temporarily at least, a definite fall in the number of blood cells, both red and white, the loss of leucocytes affecting chiefly the polymorphonuclears. Similar changes in the cellular balance of the blood no doubt follow upon the operation of ganglionectomy.

It may be suggested, therefore, with some degree of confidence, that any ganglionectomy operation must substantially disturb the physiological equilibrium in directions as yet undetermined, and that certain of the remedial effects may be due to one or more of such undetermined factors. For example, the sympathetic denervation may grossly disturb the normal mechanism of liberation of adrenergic substances into the blood stream. What effect this departure from the normal may have on the tissue cells and on the articular and periarticular structures one cannot yet say. The whole matter will require, and will certainly repay, much further study.

Clinical Details

All the six cases reported at the Cairo congress were of an extremely painful and grossly disabling type. In all many joints, both of the upper and of the lower limbs, were involved. They were all regarded as cases of chronic rheumatoid arthritis. Three patients were male and three female, and their ages ranged from 14 to 49 years. With the exception of the youngest, a girl aged 14, whose symptoms had lasted for only nineteen months, and one of the male patients, whose symptoms were of just over two years' duration, all had suffered for quite a number of years (five and a half, seven, eight, and ten years).

All the six patients were extremely crippled and practically bedridden. All suffered, some more than others, from severe pain, not only on any attempted movement of the affected joints, but even when the joints were at rest. In most there was a good deal of associated spasm of the adductor and flexor muscles, particularly of the lower limbs; in several there were present already varying degrees of contracture of the hips and knees, as well as of the joints of the upper limbs. In nearly all of them certain of the larger joints were still considerably swollen, and showed not merely thickened synovial membrane but also varying degrees of effusion.

As the result of the lumbar ganglionectomy operation (in each case bilateral), all the patients were at once relieved of pain, and obtained relief almost immediately from the adductor and flexor spasm. All were able, almost at once, to carry out movements previously quite impossible, and to permit passive movements which had been previously out of the question. In five of the six there occurred the remarkable improvement in the joints of the upper limbs as well to which reference has already been made. In this connexion it may be well to say that in every one of these cases the patient was the first person to draw attention to this remarkable, and at first unexpected, sequel to the lumbar ganglionectomy procedure. Now we have learned to expect such a result.

As regards the permanence of the improvement it is perhaps too soon yet to express a dogmatic opinion, but all six cases have been under observation at intervals since the dates of the operations, and so far there has been no sign of deterioration of function or of recurrence of painful or disabling symptoms. The first patient was operated on in August, 1931, the second in September, 1932, the third in November, 1934, the fourth a month later, the fifth in June, 1935, and the sixth on October 21st, 1935.

In all cases but one the immediate result of the operation has been to convert an apparently hopelessly crippled and helpless patient into an individual restored, more or less completely, to the possibility of normal activity, and freed from pain.

The exception was that of the 14-year-old patient, a girl, many of the joints of whose upper and lower limbs were already completely ankylosed—particularly the knees; both of her femora had been fractured a little above the knee in the forcible straightening of the lower limbs before she came under our care. Even in this case, however, the improvement in her whole condition was remarkable. The swelling of the affected joints very quickly disappeared, and pain was abolished, not merely in the lower limbs, but also in the upper, so that the grosser movements of the limbs, both active and passive, could be carried out without pain, within the limits imposed by the degree of ankylosis already established.

Study of the detailed reports of these six cases which follow will serve, I believe, to convince the most incredulous of the reality of the benefit secured by the operative procedure adopted.

CASE I.—*Extreme disabling polyarthritis of upper and lower limbs, but particularly of the lower limbs, in a woman aged 27. Condition probably of rheumatoid nature. Symptoms of seven years' development. Associated extreme*

adductor and flexor contractures of the lower limbs. Ultimately almost complete disablement, so that for two years the patient had been practically bedridden. Bilateral ganglionectomy and trunk resection on September 30th, 1931, followed at once by improved circulation in feet and legs, by rapid relaxation of adductor and flexor spasm, and by diminution of the contractures of hips and knees. Power of walking soon restored, and pain banished. Practically complete restoration of useful function.

This patient was admitted to the Western Infirmary, Glasgow, on August 7th, 1931, complaining of stiffness of the lower limbs of progressive development during seven years. At the onset the pain was present in both knees for a year or two, especially in the right. The knees then improved somewhat, but a similar painful condition developed in the shoulders and in the elbows. Sixteen months previously the right hip had become painful, and six months later the left. At the same time the knees once more became affected. With the pain, in the case of each of the joints, there was also a variable amount of swelling and considerable limitation of movement. For about two years before admission the patient had not been able to leave her home, and was for a good deal of the time actually bedridden. During the final ten or twelve months both hips became so stiff that she was hardly able to move them at all.

On admission both knees were markedly contracted, as were also the hips, the thighs being flexed upon the abdomen and the legs on the thighs. Both lower limbs were strongly adducted. When even moderate force was employed to try to separate them for only two to three inches considerable pain was caused in the region of the hips and along the inside of both thighs. Any attempt to extend the knees passively from their position of flexion-contracture caused severe pain in the hips and in the small of the back. Both hips were very stiff. X-ray examination showed the presence of marked arthritic changes in both hip-joints.

On September 10th, 1931, in order to determine the nature and extent of the limitation of movement of the hips and knees, the patient was examined under a general anaesthetic. It was then found that considerable range of movement at the hips was possible. Even under deep anaesthesia, however, the adductor muscles of both thighs remained tense and contracted. These contracted muscles were forcibly stretched, and the spasm was overcome by a combination of forcible traction and local hammering of the contracted muscle bellies, with considerable improvement in respect of the possible range of passive movement.

In view of the favourable finding just mentioned, it was decided that ganglionectomy might well be beneficial. Accordingly, on September 30th, 1931, a bilateral lumbar ganglionectomy and trunk resection was performed by the trans-peritoneal route. A good deal of difficulty was encountered by reason of enlarged lymph glands, especially on the right side. The ganglia were removed separately in the case of the right chain, but the left ganglionic trunk and ganglia (second, third, and fourth left) were removed *en bloc*.

Half an hour after the operation had been completed both feet were warmer than previously, but the left was much warmer than the right. A few hours later the temperature was the same on both sides. Almost at once the patient obtained great relief from pain, and the movements of the hips and knees rapidly improved, while the adductor spasm of the thighs soon disappeared altogether. Normal healing of the abdominal wound took place; almost immediately after that was complete the patient was allowed up, and was able to get about the ward unaided. She was discharged from hospital on December 14th—that is, two and a half months after the ganglionectomy. By this time she was able to walk about quite actively, and had no pains or aches. This remarkable degree of improvement has been satisfactorily maintained since.

No observation was made in this case regarding any coincident improvement in the upper limbs.

CASE II.—*Painful and grossly disabling polyarthritis of rheumatoid type in a man aged 43. Duration upwards of five and a half years. All the larger joints seriously affected, but also the smaller joints of fingers and toes. Most profound disablement of the lower limbs, especially of the knees and hips. Marked flexion-contracture of the knees, and spastic*

contraction of adductors of thigh. Completely bedridden for four months before admission. Bilateral lumbar ganglionectomy on November 10th, 1932, followed by immediate relief of pain and rapid restoration of function in the lower limbs. Almost full use of the lower limbs quickly restored, and this improvement maintained. Improvement also in the case of the upper limbs.

A man, aged 43, was admitted to the Western Infirmary, Glasgow, on September 5th, 1932, under the care of Professor T. K. Monro. He was suffering from a severe polyarthritis, of rheumatoid type, of five and a half years' duration, which latterly had so profoundly disabled him that for some months he had been practically bedridden. Both before admission to hospital and while in a medical ward he had received the usual medical course of treatment, including protein shock therapy, without any improvement. Latterly, surgical treatment was suggested.

His illness had begun somewhat insidiously, with pain and swelling of isolated joints, the left wrist, left knee, and, in their turn, practically all the larger joints of both upper and lower limbs, including the joints of the fingers and toes. For the last three years he had been very lame, and he developed contracture deformities of both knees, which interfered with progression. In 1930 his lower limbs had been kept for six months in extension, but without effect. By June, 1932, he was quite unable to walk. There was no history of acute rheumatic fever.

On November 1st, 1932, he was transferred to my ward for surgical treatment. At this time most of the joints were swollen, and movements of the joints were much limited. Both knees were in a state of partial flexion, and were extremely stiff and painful. The hands were markedly deformed, the fingers notably ulnar-flexed. Both wrists were swollen and painful. There was great limitation of movement also at the elbows. The lower limbs were in a state of continuous adduction, and any attempt at overcoming adductor spasm caused great pain. The circulatory system generally seemed satisfactory, though there was a certain degree of clamminess and coldness of the feet.

On November 10th bilateral lumbar ganglionectomy was performed, the second, third, and fourth ganglia on each side being removed, with the associated trunks and connexions. Some difficulty was experienced by reason of two separate chains of enlarged lymph nodes on each side, which obscured at first the ganglionic chains. Once the lymph nodes had been cleared away, however, there was no dubiety in the recognition of the ganglionic chains.

Two days after the operation it was noted that the feet were much warmer and less clammy, that there were no pains in the knees, that the adductor spasm had passed off, and that the lower limbs could be moved much more freely, both actively and passively, and without pain. By December 7th—that is, a month after operation—at the time of his discharge from hospital, he was able to walk very satisfactorily; he could straighten his left knee almost fully, and his right very much better than previously.

On May 24th, 1933, about six and a half months after his operation, he reported. He was able to walk well, though using a stick. He had full and free movements of his right hip. There was slight diminution of abduction and adduction of the left hip. Movements of the right knee were normal. Movement of the left knee was still a little limited. Ankle movements were normal. It was noted also that movements of the wrist and of the fingers were substantially improved.

CASE III.—Rapidly progressing and ankylosing polyarthritis of both upper and lower limbs, of rheumatoid type, in a girl aged 14. Duration of symptoms nineteen months. Medical treatment, including protein shock therapy, of no avail, so that ultimately she was completely disabled, and most of the larger joints were ankylosed. Bilateral lumbar ganglionectomy on November 19th, 1934, was followed by the rapid disappearance of swelling of the affected joints and the abolition of pain, not merely in the lower limbs, but also, in marked degree, in the upper limbs, so that the grosser movements could be carried out, both actively and passively, without pain, within the limits imposed by the ankylosis already established.

This girl was admitted to the Western Infirmary, Glasgow, on February 7th, 1934, to the wards of Professor T. K. Monro,

suffering from a rapidly progressive polyarthritis of practically all the joints of the upper and lower limbs, the first symptoms of which had been observed in April, 1933. The girl had had her tonsils removed in 1928, but there was no joint trouble until April, 1933. Her ankles were first affected, but later the other large joints of the upper and lower limbs became involved, particularly the knees and elbows, which became greatly swollen, stiff, and painful. In the summer of 1933 she was for nineteen weeks in the Paisley Royal Infirmary where, in addition to other treatment, she had protein shock therapy, with no remedial effect.

When admitted to the Western Infirmary both elbows and knees had become permanently flexed and fixed. Extension failed to straighten the knees, and, under a general anaesthetic, the lower limbs were forcibly straightened by fracturing the femora a little above the knee.* Thereafter the lower limbs were kept in splints, in the extended position. The elbows were almost completely ankylosed, and in a very disadvantageous position, so that the girl was unable to feed herself. Any attempt at movement of the limbs, upper or lower, active or passive, caused great pain. Ultimately, though the advance of the disease seemed to have become arrested, pain continued to be a distressing feature.

The patient remained in Professor Monro's ward from February, 1934, until November, when she was transferred to my ward. At this time she was completely helpless. She was unable to alter her position in bed; the knees were markedly swollen and ankylosed; the elbows were in semi-flexion; movements of ankles and hips were impossible, partly by reason of severe pain; the lower limbs were still in splints.

On November 19th, 1934, I performed bilateral lumbar ganglionectomy by the transperitoneal route. The second, third, and fourth lumbar ganglia and connexions were removed on both sides, and this was verified later histologically. From the time of the operation there was almost at once complete abolition of pain, and rapid diminution of the swelling of the affected joints followed. The splints were removed from the lower limbs and the grosser movements of these limbs could be carried out, both actively and passively, quite without pain, though the ankylosed knees remained stiff. It was observed that not only was there a marked remedial effect in the lower limbs but there was a definite and considerable improvement also in the upper limbs, so that quite a fair range of movement at the wrists and of the fingers became possible without pain. On December 13th, 1934, three and a half weeks after the operation, the girl was discharged from my ward greatly improved.

CASE IV.—Generalized and progressive polyarthritis of the joints of upper and lower limbs, of rheumatoid type, of slow development over a period of years, in a woman aged 33. Earliest symptoms observed ten or a dozen years before, affecting first the smaller joints of the fingers and toes, but later involving also the wrists, ankles, knees, and elbows. Patient at last quite unable to walk and practically bedridden. Bilateral lumbar ganglionectomy on December 26th, 1934, followed by almost immediate relief of pain and diminution of swelling of the affected joints, and by rapid recovery of practically full movements of these. There was also observed a remarkable associated improvement in the joints of the upper limbs, and particularly in the hands and fingers.

This woman was admitted to the Western Infirmary, Glasgow, on December 14th, 1934, suffering from a very disabling polyarthritis of both upper and lower limbs, of slow development over a period of years. The onset of symptoms had been very insidious. Ten or a dozen years before the patient had had some recurrent pain and swelling of the smaller joints and of the fingers and toes, and later of the wrists, ankles, and knees. At a later stage still the elbows became involved to a slight extent. During the last two years the progress of the disease had become more marked, and particularly during the later months of 1934. For at least six weeks before her admission to hospital she had been quite unable to walk, and was practically confined to bed.

On admission there was marked swelling of the metacarpophalangeal and the first interphalangeal joints of all the fingers of both hands. Both wrists were also considerably

* For this part of the treatment I had no responsibility. It was carried out by another surgeon.—A. Y.

swollen and deformed, especially the left. Both ankles were swollen, stiff, and painful. Both knees were swollen, but the left more so than the right, and both were tending to adopt a position of flexion-contracture. Movements of all the joints were painful, and particularly movements of the joints of the lower limbs, especially the left knee. The skin of the legs and feet, particularly the left, was cold and clammy, and frequently the feet and ankles were covered with a clammy sweat. X-ray examination showed well-marked changes—regarded as being of a rheumatoid type—in several of the joints, including the astragalo-scapoid joints of both feet.

On December 26th a bilateral lumbar ganglionectomy was performed. At the operation there was found to be a very definite fleshy enlargement of the pre-aortic lymph nodes, and in consequence there was some little difficulty in identifying the ganglionic chain on both sides, as well as some difficulty from venous bleeding, which, however, was successfully checked. Almost immediately after the operation the pains in the lower limbs were completely relieved and the swelling of the joints quickly diminished. The skin became warm and dry, and the patient recovered practically full movements of the diseased joints.

A very interesting feature of the case was the associated improvement in the joint conditions in the upper limbs. The patient was the first to draw attention to this improvement. Almost from the first she was quite clear in her mind that the upper limbs were much more comfortable than they had been before the operation, and as the weeks passed she regained in remarkable degree free movement at her wrists and of her previously much crippled fingers. She was soon able to walk quite well. She was discharged from hospital on February 4th, 1935, fully six weeks after her operation. She has been seen since at frequent intervals, and the improvement has been well maintained.

Note.—It may be of interest to state here that it was as a result of the personal recommendation of this patient that the following gravely crippled patient sought similar treatment.

CASE V.—Extreme crippling polyarthritis of practically all the joints of both upper and lower limbs, of rheumatoid type, in a man aged 42. Marked deformities and muscular contractures. Duration upwards of eight years. Various non-surgical lines of treatment of no avail. Ultimately, almost complete disablement; much pain at rest and on attempted movements, active or passive, particularly of the lower limbs. Bilateral lumbar ganglionectomy on June 12th, 1935, followed at once by relief of pain, by relaxation of muscular spasm, and by rapid improvement in function of the limbs, even of the most disorganized and disabled joints. Probably a permanent improvement. Notable improvement in function of the joints of the upper limbs as well as of the lower.

This patient was admitted to the Western Infirmary, Glasgow, on June 5th, 1935, complaining of a crippling polyarthritis of the joints of both upper and lower limbs, with marked contractures. The illness began in 1927, some months after a gastric or duodenal perforation, with pain, stiffness, and swelling of both ankles. The onset was gradual. There was no acute rheumatic fever. The disease steadily advanced, so that in 1932 all the joints of the limbs had become affected, in varying degree, also the temporo-mandibular joints and the lower joints of the spine. He had medical treatment in the Royal Infirmary, Glasgow, in 1928 and 1932, with no improvement. Amongst other lines of treatment he had protein shock therapy, but without remedial effect. Latterly he had become almost completely bedridden.

On admission he was much emaciated, and there was considerable muscular atrophy, particularly of the muscles of the thighs and legs. Both hips were in a state of partial flexion; neither could be extended to any degree. Both were in a permanent position of external rotation, the right more markedly than the left; the right, in fact, was in a state of partial dislocation. Both knee-joints were swollen, and the bones were unduly prominent. Both ankles were stiff, and the feet unduly arched. The toes were in a position of marked plantar flexion, and were very stiff and deformed. The knees could not be extended beyond an angle of 45 degrees of full extension. The ankles, especially the right, were almost ankylosed. The shoulders and elbows were less affected, also the left wrist and hand, though the fingers of the latter

showed characteristic flexion deformity. The right hand was greatly deformed, all the fingers being acutely flexed upon the palm, and deflected to the ulnar side. They were practically incapable of voluntary movement. The right hand could not be used for grasping. The patient was almost always in pain, and any attempt at abducting the thighs or moving the knees, ankles, or toes caused considerable outcry. Both feet and legs were constantly clammy, wet, and cold.

An x-ray examination on June 10th, 1935, showed the following: (1) shoulders and elbows—comparatively normal; (2) hand and wrists—osteoporosis of bones and general narrowing of joint spaces with destruction of cartilage; (3) pelvis—arthritis of right sacro-iliac and right hip-joints, destruction of cartilage, and some localized bone destruction (a low-grade type of infective arthritis suggested); (4) knees—destruction of cartilage, especially of articular surface of patella, with osteoporosis; (5) ankles—well-marked destructive arthritis, joint spaces diminished or destroyed, some bony ankylosis.

On June 12th a bilateral lumbar ganglionectomy and trunk resection was performed. The ganglionic trunk on both sides was somewhat thin and the ganglia unusually small. During the first few days after the operation the patient was somewhat ill by reason of a post-operative bronchitis and a limited patchy bronchopneumonia; fortunately this cleared up quickly. As regards the general joint condition, however, the result from the very first was almost magical. Pain entirely disappeared, and the lower limbs became warm and dry. The range of movement, especially of the lower limbs, greatly improved, and even the disabled and partly luxated right hip showed improvement. Muscular spasm disappeared almost from the first, and soon even the contracted knees could be straightened to an unexpected extent. The patient was finally discharged from hospital on July 13th, greatly and probably permanently improved. A certain amount of walking was now possible, but this was somewhat limited by reason of the still existing plantar flexion of the great toes.

A particularly gratifying feature noted in this case, as in several other similar cases, was the improvement which occurred simultaneously in the joints of the upper limbs. The wrist-joints became more flexible, and finger movements—even those of the right hand—very greatly improved. In the course of a few days the patient was able to use the previously completely useless right hand for a variety of purposes hitherto quite impossible, and he developed quite good grasping power. He himself drew attention to the gratifying improvement in function in the upper limbs—all the more grateful to him because so unexpected.

CASE VI.—Generalized and progressive polyarthritis of the larger joints, of both upper and lower limbs, of rheumatoid type, of slow development over a period of two years, in a man aged 49, the subject of an old anterior poliomyelitis. Non-surgical treatment, including protein shock and vaccine therapy, of no avail. Considerable pain and limitation of movement of the affected joints. Bilateral lumbar ganglionectomy on October 21st, 1935, followed at once by complete banishment of pain and by improved range of movement. Remarkable improvement also in respect of the joints of the upper limbs.

This man was admitted to the Western Infirmary, Glasgow, on September 15th, 1935, complaining of pain and stiffness in most of his larger joints of over two years' duration. Before admission he had been seen in consultation by Mr. W. A. Mackey, who arranged for his admission. The trouble began somewhat insidiously, involving first both ankle-joints, and extending later to most of the other distal joints, including, particularly, both knees, both elbows, and both wrists. Various lines of treatment had been tried, including protein shock and vaccine therapy, but without any improvement. In other ways the man appeared healthy, and nothing was discovered in the way of material focal sepsis. He had had from time to time a slight urethral discharge, but there was no evidence that this had in any way been responsible for the production of the polyarthritis. His tonsils were healthy.

When admitted to hospital there was abundant evidence of a progressive and disabling involvement of many of the larger joints. Both ankles were still swollen, the swelling being in part, at least, periarticular. Both knees were swollen, with obvious thickening of the synovial membrane, and also a good deal of effusion into the joints. There was marked peri-

articular swelling of both wrists and of both elbows. Movements of all of these joints were, in varying degree, limited and painful, except in the case of the right elbow. The range of flexion of the left elbow was limited to 60 degrees, that of extension to 160 degrees. The interphalangeal joints and the metacarpophalangeal joints of both hands were not very obviously abnormal in appearance, but in the case of the right hand there was the beginning of a definite ulnar deviation of the fingers, and the hand grip was definitely lessened.

An x-ray examination on September 17th showed the following: polyarthritis, with associated destruction of the articular cartilage of the left knee; narrowing of joint space of the left elbow, but no actual ankylosis; early changes in the right wrist and right intertarsal articulations; early changes suggestive of rheumatoid arthritis at the carpus and at the proximal interphalangeal joints. Both legs showed considerable wasting of muscles below the knee, and on each side there was a well-marked pes cavus. These were sequels to an anterior poliomyelitis in early life, and were more marked in the left leg than in the right.

On October 21st a bilateral lumbar ganglionectomy and trunk resection was performed by the transperitoneal route. Excellent exposure of the ganglionic chain was obtained on both sides, and the usual three ganglia were removed. Many enlarged lymph nodes were found in close relation to the ganglionic chains. At the conclusion of the ganglionectomy a markedly atrophic appendix, which was attached by definitely inflammatory adhesions to the root of the mesentery, was removed. The operation was well borne.

The subsequent progress in the days immediately following operation was generally satisfactory. The only troublesome feature, which persisted for several days, was the difficulty the patient had in voiding urine; on several occasions catheterization was necessary. On the other hand, the operation was followed at once by the complete banishment of joint pains and by a very obvious improvement in the range of voluntary movements. A noteworthy feature, too, was the fact that the patient himself drew attention, within the first twenty-four hours, to the substantial improvement which had taken place in the upper limbs as well as in the lower limbs. He found, for example, that he was able to use his right hand much more freely, and with considerably enhanced power. It was also observed that not only were the feet, which had previously been moist, clammy, and cold, now warm and dry, but that the same improvement, though to a less degree, had occurred in the case of the hands.

A Seventh Case

Since the last of these was dealt with I have had occasion to treat a further case of a similar kind, and though it is true that so far the result has been less satisfactory than any of the six—indeed, all the more for this reason—it seems well that it should be reported in full with the others.

The patient was a man aged 38, who had had symptoms for fifteen years, and had been completely disabled from work for nine years. Many joints of both lower limbs and upper limbs were involved, and some of them were almost ankylosed. He was operated on in February, 1936, and obtained immediate relief from pain, as well as a considerable improvement in function; but further progress seems for the present to have been suspended. He is able to get about with considerably more freedom than previously, and has much less pain in his joints, but there seems to have been some slight recurrence of swelling, particularly of his knees, and the ultimate result is yet undetermined. The case is worthy of record, if only to emphasize the fact that not every case will necessarily respond to treatment in an equally satisfactory way.

The following is a detailed report of this seventh case.

CASE VII.—*Severe, progressive, and widely generalized polyarthritis of joints of upper and lower limbs, of rheumatoid type, in a man aged 38. Earliest symptoms appeared fifteen years before. Patient had been quite unable to work for the last nine years. Admitted to hospital in September, 1935, with a view to lumbar ganglionectomy, but a severe attack*

of appendicitis, which demanded emergency operation, made the postponement of ganglionectomy inevitable. It was not, indeed, until February 26th, 1936, that the ganglionectomy procedure was carried out. A bilateral operation was followed at once by complete relief of pain, by diminution of the swelling of the affected joints, and by substantial improvement in function. This improvement occurred in the upper limbs as well as in the lower.

This patient was admitted to the Western Infirmary, Glasgow, on September 15th, 1935, complaining of painful and stiff joints of both upper and lower limbs, of many years' duration. The condition dated back to 1921, when the patient was laid up for over two months with successive acute inflammatory conditions affecting most of the joints, which gradually became stiff; some of them had become almost ankylosed. For the last nine years he had been quite unable to work. There was no history of venereal infection. The joints affected were as follows: fingers of the right hand, swollen, stiff, and contracted; right wrist, almost ankylosed; elbow, ankylosed at an angle of 90 degrees; right knee, swollen, stiff, and movement restricted to about 30 degrees (90 degrees flexion and extension to 120 degrees); left elbow, extension limited to 140 degrees; left knee, like the right. In other respects the man appeared healthy.

He was admitted with a view to early ganglionectomy, but immediately following admission he developed an acute appendicitis with abscess formation, which called for immediate operation, and which for the time being ruled out the ganglionectomy procedure. An x-ray examination of the joints at this time gave the following result: "Polyarthritis, associated with marked narrowing of the joint spaces of both knees, and ankylosis of the right tarsus. Marked articular destruction and ankylosis have taken place in the right elbow. The right wrist-joint also partially ankylosed. No changes were made out in the spine." The man was discharged from hospital on October 9th, 1935, three weeks after his appendix operation. His joint condition was as before.

He was readmitted on February 15th, 1936. At this time further examination was made of his joints. There had been little change from the earlier note. The knees were perhaps a little more swollen, and the movements of both knees a little more restricted. The ankle joints were more definitely involved. The feet were generally covered with a cold, clammy sweat.

On February 26th a bilateral lumbar ganglionectomy and trunk resection was performed by the usual transperitoneal route, the second, third, and fourth ganglia being removed on both sides. On the right side these were removed in an unbroken chain, but on the left side the third and fourth, with the corresponding trunk and connexions, were removed separately from the second, which was unusually high, and which was united to the third by a very thin nerve strand. There were considerable adhesions in the region of the caecum.

The after-course was uneventful. From the time of the operation the man's sensations were completely altered. His feet and legs felt warm and dry, and he himself drew attention to the fact that his hands and right wrist had participated in the improvement. He was dismissed from hospital on March 22nd, when he was able to get about much more freely than formerly. The movements of all his joints were improved, notably the right ankle and both knees. He was still using walking-sticks, but it may be said that the general improvement appeared to be well maintained. He reported on April 18th. He was now able to get about much more easily, but he had some pain in his right hand, probably due to the use of his stick.

Osteo-arthritis

In my original paper before the Royal Medico-Chirurgical Society in Glasgow in 1931 reference was made to the successful result of periarterial sympathectomy in the treatment of certain doubtful joint conditions, and one case was cited of destructive osteo-arthritis of the ankle and tarsus in a middle-aged man (a case of doubtful pathology), in which an unexpectedly favourable result was obtained by the periarterial procedure. Thought, at the first, to be possibly a chronic tuberculous

lesion, the condition was found ultimately to present features inconsistent with such a diagnosis, but the exact pathology could not be determined. More as an experiment than anything else the periarterial operation was undertaken, and the result was so satisfactory that the patient, who had been before almost completely crippled, was able to return to work, and has been able to work normally since, the affected ankle being restored practically to normal.

It is admitted that, only in the exceptional case, can such a favourable result be expected in advanced cases of osteo-arthritis. Some temporary improvement may be effected, especially in respect of pain, but only occasionally can permanent benefit be achieved. It is probable that, in cases of osteo-arthritis which have reached anything like an advanced stage, not even the more severe procedure of ganglionectomy can hold out substantial promise of benefit, except perhaps in the relief of pain and in diminishing spasm.

I am able to record, however, one case of a somewhat striking character, where such a favourable result followed upon the carrying out of a lumbar ganglionectomy. The case was one of advanced, grossly disabling osteo-arthritis of the left hip-joint in a woman aged 62. Her symptoms had developed progressively during four years, until ultimately she walked only with difficulty, and suffered severe pain on any attempt at walking. The region of the left hip was swollen. There was marked objective crepitus, and the movements of external rotation and of abduction at the hip were markedly limited. X-ray examination showed advanced osteo-arthritic change, with lipping of the acetabulum and considerable increased density of the bone. So much was she crippled and so insistent her pain that the patient herself demanded that the ganglionectomy operation should be tried. Her request was granted, and the operation was followed almost at once by a manifest improvement in the circulation in the limb by almost complete disappearance of the pain, and by considerable improvement in all the movements at the hip. This improvement was maintained; when she was seen many months after, pain had quite gone and she was able to walk with comfort. Detailed notes of the case follow.

CASE VIII.—*Advanced disabling osteo-arthritis of the left hip-joint, of four years' duration, in a woman aged 62 years. Great aggravation of disability in the later period. Gross osteo-arthritic change in the hip-joint, confirmed by x-ray examination. Severe pain from the hip down the anterior and external aspect of thigh, knee, and leg, ultimately making walking almost impossible. Left lumbar ganglionectomy and trunk resection on May 22nd, 1933, followed at once by an improvement in the peripheral circulation and rapid amelioration of the painful disability. Pain entirely disappeared, and substantial improvement occurred in the movements of external rotation and abduction at the hip. Improvement maintained—probably permanent.*

This woman was admitted to the Western Infirmary, Glasgow, on April 27th, 1933, complaining of severe pain in and around the left hip, and pain shooting down from the front of the hip along the front of the thigh and the anterior and outer aspects of the knee, leg, and foot. The symptoms were of gradual but progressive development during about four years. At first pain was felt chiefly on walking, but at a later period even when at rest. As a result the patient walked only with a marked limp and required the use of walking-sticks. There was no pain in any other joint. The region of the left hip was swollen. There was as a rule no subjective sensation of creaking, but objectively crepitus could be made out. The movements of external rotation and of abduction of the left lower limb were markedly limited. There was no ascertained focal sepsis anywhere. X-ray examination showed advanced osteo-arthritis of the left hip-joint, with lipping of the acetabulum and local increased density of the bone. Elsewhere the bone texture was normal.

The patient became so crippled that, in spite of her age, she was anxious that operative treatment should be given a trial. Accordingly, on May 22nd, a left lumbar ganglionectomy and trunk resection was carried out. The operation was rendered somewhat difficult by excess adipose tissue in the anterior abdominal wall and retroperitoneally. In consequence the third and fourth ganglia were removed in one piece, with their connecting trunk. There was a certain amount of doubt as to whether the second ganglion was actually removed.

Immediately following the ganglionectomy, however, there was a manifest improvement of the circulation of the limb, the foot and leg becoming definitely warmer; the improvement was progressive and steadily maintained. The patient was discharged from hospital on June 23rd, able to walk quite freely without pain, and with much less limitation of the movements of external rotation and abduction. She was seen several times after her discharge, the latest occasion a good many months after, when her general condition was excellent, there was no pain, and she was able to walk with comfort.

Conclusion

In view of the results detailed above it is clear that by a sympathectomy procedure, such as ganglionectomy, a substantial degree of relief can be given in many cases of profoundly disabling polyarthritis, especially in those of a rheumatoid type, and even in certain cases of osteo-arthritis. Many completely bedridden and crippled patients may thereby be relieved from pain to a remarkable degree, and may even be restored to the possibility of a useful life.

While it is perhaps more particularly in those cases where the active disease has exhausted itself, or, as it were, has burned itself out, that the ganglionectomy procedure will prove most efficacious, it may well be considered whether the same procedure might not be worth trying as a definite therapeutic agent even in the earlier, more active stages of the disease. It should have the effect of producing such a degree of improvement in the peripheral blood supply and in the local temperature as to exert a valuable therapeutic influence upon the local joint conditions.

It may well be, also, that the disturbance so induced in the physiological equilibrium of the tissues of the limbs, in directions as yet undetermined, may have a beneficial influence both on articular and on periarticular structures, and may in this way alter favourably the whole therapeutic outlook.

REFERENCES

- ¹ Buckley, C. W.: *Lancet*, 1936, i, 1023, 1081.
- ² Young, Archibald: *Trans. Glasgow Medico-Chirurgical Society*, 1933-4, xxviii, 26.
- ³ Idem: *Trans. Xe Congrès de la Société Internationale de Chirurgie*. (In the press.)
- ⁴ Idem: *Glasgow Med. Journ.*, June, 1931, p. 305.
- ⁵ Rowntree, L. G., and Adson, A. W.: *Proc. Staff Meetings Mayo Clinic*, November 21st, 1928.
- ⁶ Idem: *Ibid.*, September 4th, 1929.
- ⁷ Fraser, John: *Edinburgh Med. Journ.*, 1931, xxxviii, 201.
- ⁸ Ross, J. Paterson: *Brit. Journ. Surg.*, October, 1935, xxiii, 441.
- ⁹ Hench, Henderson, Rowntree, and Adson: *Journ. Lab. and Clin. Med.*, 1930, xv, 1247.
- ¹⁰ Leriche, R.: *Nelson's Living Surgery*, iii, 787.

A. Gutman (*Thèse de Paris*, 1936, No. 313), who records three illustrative cases in men aged 58, 60, and 71 respectively, states that cancer of the male breast presents the same features as in the female. The prognosis depends chiefly on the stage of the disease at which the diagnosis is made and the variety of the malignant growth. Recurrences are frequent, and death is due to cancerous cachexia. Pulmonary metastases are the rule, and occurred in two of Gutman's patients who died, while the third was lost sight of. According to some authorities, such as Velpeau, Birkett, and Liston, the prognosis of cancer of the male breast is more favourable than that of mammary cancer in the female owing to the greater frequency in the males of the less malignant forms of cancer.