Our observations also correspond to those of Shecket, Friedmann, and Nice.³ They reported that adrenalectomized rats showed an increase in blood platelets up to 100 per cent.

TABLE VII

	Leucocytes	Erythrocytes	Blood Platelets
October 29th, 1935, before intramuscular injection of 2 ampoules cortigen	3,800	4,040,000	303,000
$\frac{1}{2}$ hour after injection...	4,000	3,800,000	250,000
1 " " " " " "	3,800	3,500,000	152,000
2 hours " " " " " "	3,500	3,200,000	170,000
4 " " " " " "	3,800	4,400,000	240,000
24 " " " " " "	5,000	3,600,000	350,000
48 " " " " " "	4,000	4,200,000	344,000
72 " " " " " "	4,000	4,200,000	366,000

Summary

The blood platelet count in the peripheral blood is increased by the hormone of the thyroid gland. The action is more constant than in the case of red blood cells. The increase usually appears twenty-four to forty-eight hours after the administration of the hormone and has disappeared within about seventy-two hours. The thyrotropic hormone acts in the same way, but its action is more protracted. In a case of chronic thrombopenia in a child a gradually increasing number of platelets was counted after repeated administration of thyrotropic hormone (single dosage, 500 rabbit units). The hormone of the adrenal cortex showed a diminishing effect on the number of blood platelets in a large number of experiments, but in several other experiments this result was not obtained.

REFERENCES

- ¹ Zondek, H.: *Deut. med. Woch.*, 1922, 31; *Klin. Woch.*, 1926, v, 876.
² Bose, P.: *Klin. Woch.*, 1924, 1367.
³ Shecket, Friedmann, and Nice: *Proc. Soc. Biol.*, 1935, xxxii, 608.

BLOOD TRANSFUSION IN CHILDHOOD*

BY

DENIS BROWNE, F.R.C.S.

SURGEON, HOSPITAL FOR SICK CHILDREN, GREAT ORMOND STREET

I have re-read my paper on this subject of some five years ago, and rather to my surprise find that I still agree with almost all of it. I think anyone else who is interested in it will have the same experience—that is to say, he will agree that blood transfusion has very rapidly reached a stage where it appears destined to remain for some time. Such valuable innovations as the prolonged drip method will of course be introduced, but in this there is no change of principle or widening of the field of application; there is merely an increase in efficiency.

However satisfactory this settling of the status of blood transfusion may be there is another aspect from which the static condition of affairs over the last ten years or so is anything but a matter for the profession to congratulate itself upon. Ten years ago all who knew anything about the subject were convinced that this method of treatment, one of the few which we can say will save lives that otherwise would be certainly lost, was far too little used. Is not the same true to-day? Consider the practice of the profession throughout England. Would anyone claim that more than 10 per cent. of those who would be benefited by a blood transfusion get one? Would anyone like to

estimate coolly the number of lives lost which could have been saved by one? Admittedly the number of transfusions given is increasing, or at least, in the absence of any proper recording of the work of the profession as a whole, we have good reason to think it is. But it is increasing very slowly, and in present conditions it seems to me that it is never likely to reach the optimum figure.

Of course if we think too much along these lines we are apt to fall into one of the last infirmities of mind, whether noble or otherwise—the construction of Utopias. Imagine, for instance, that the medical profession were to reconstruct itself—no, reconstruction is not the word, it has at present no construction whatever—were to design for itself a structure with the sole object of efficiency in the treatment of the sick. Would not one of its first tasks be the covering of the entire country to ensure that anyone could get a blood transfusion with reasonable speed? This would call for specially trained men, not necessarily doing nothing else but (this is the important thing) constantly performing transfusions. It is no good blinking the fact that a blood transfusion is now, and appears likely to remain, a complicated and fairly difficult job. If a man learns how to do his first one from a book and repeats it at the infrequent intervals afforded by any single general practice he will have difficulties and failures that he would not get if properly trained and constantly repeating the operation.

I am assuming in this Utopia that immediately blood transfusion appeared as a method of treatment it would have been investigated under strict scientific conditions; and that as soon as its value had been established its use in appropriate cases would have been incumbent upon every member of the profession without exception. Also that all the results obtained would be tabulated and practice varied as necessary according to what they revealed. And here you see the absurdity of Utopias. This would mean a discipline directed towards efficiency equivalent to that of the crew of a liner; a system of bookkeeping almost as efficient as that in a department store. It is obviously utterly beyond us.

The population would have to be taught that the suggestion of a blood transfusion does not mean that all hope is abandoned, and that the giving of blood does not need dauntless heroism. There are, however, limits to any Utopian's imagination, and getting the lay press to co-operate with us on this and other points is far outside my own.

Then again, if we believe that the injunction "Know thyself" is of any value, a useful thing for the individual to know is his own blood group, a knowledge that may save several vital hours. I suggested, in the paper I have mentioned, that a very useful item of information to be added to religion upon an Army identification disk would be the individual's blood group. I have myself an extremely defective sense of humour, and I have never been able to understand why this suggestion, springing from a certain experience both of transfusion and of recently inflicted wounds, is invariably regarded as a deliberate joke.

Another department of Utopia would deal with the purely technical problems of transfusion. It would therein be an offence to repeat a statement, now sure of universal approval in any medical gathering, that every man will get the best result by using the method he is accustomed to, whichever of the nine and sixty ways that may be. The underlying fallacy would have been discovered upon which this statement rests: the assumption that because all progress springs from variation therefore all variations should be encouraged. The difference would have been recognized between those variations which spring from inability to master the best current practice and those which arise from dissatisfaction with these methods by one who really understands them. In consequence there

* Read in opening a discussion in the Section of Diseases of Children at the Annual Meeting of the British Medical Association, Oxford, 1936.

would be a uniform standard practice, only to be changed after suggested variations had been considered, tested, and approved. This again is rather beyond us; it means a uniform standard of efficiency approaching that attained in such pursuits as hurdling, the crawl stroke in swimming, or piloting an aeroplane.

As a part of the department of technique there would be a factory in which the necessary instruments of the profession would be made. This would do away with one of our primary curses, in that our instruments are made by people who do not understand their use, and used by people who do not understand their making. If one tries to conceive a similar state of affairs in the manufacture of motor cars or shotguns one may get a suspicion that possibly there is here a chink for inefficiency to creep in so long as our present system persists. And here again I find that I have more than once been humorous without intending it. Everyone knows that if money is required to pursue some line of scientific research—for example, something to do with guinea-pigs and test tubes—it is quite easy to get hundreds of pounds to help carry out the work. But the suggestion that small amounts of money should be available for research into instruments or splints is invariably greeted with loud laughter. This really is to me a curious point; in the last resort the surgeon is a workman whose patients live or die by what he does with his hands. Yet if a single book has been published in any language upon the theory of surgical instruments or surgical technique I have yet to hear of it.

Some Points in the Technique

I may conclude by returning to practical matters. There are certain points of difference between the giving of transfusions to children and to adults that to some extent determine the technique to be used in the former. Direct transfusion becomes inadvisable for several reasons, the most important being that it needs a speed of flow, if the blood is not to clot, that is far too great for small children. This question of speed of flow is, I believe, of the utmost importance, and I find that I tend to put the blood in more slowly the more transfusions I do. It may need considerable patience to spend a full half-hour giving 50 c.cm. of blood to a marasmic infant; but if it is looked upon as corresponding to a dose of three pints in an adult the need for caution appears. I have known of two deaths and several alarms caused solely by putting the blood in at rates which would be quite suitable in an adult. If indirect transfusion is used, the citrate method seems to me by far the best available, but only if the solution is properly prepared. Dry sterile crystals made up in ampoules of 2 grams, dissolved in water specially prepared for intravenous injections, cut down reactions almost to vanishing point. A stale solution made with anything labelled distilled water will mean a very large proportion of rigors after transfusion.

Then there is the question of the propelling force for the blood. To force a viscid fluid down veins which at the best are of small calibre, and which in some of the most desperate cases are spasmodically contracted, may need great pressure. Gravity is not nearly enough, and air pressure has the very serious disadvantage of being explosive. If a connexion or cork gives way the result may be the loss of the blood, with or without the drenching of the operator, or, if there is some temporary blockage, a dangerous amount may suddenly be shot into the circulation of the recipient when it is released. It is probably because of this difficulty that such dangerous procedures as injection into the superior longitudinal sinus and the peritoneal cavity have been advocated.

Finally, for a procedure that is often an emergency one, it is essential that at a pinch it can be carried through without any assistance at all. Too many methods,

especially those of the "simpler" type, call for at least two intelligent and clean helpers. My own attempt to meet these requirements and many others which there is no space to set out have already been described. The only thing I have added to it in the last few years is a needle for withdrawing the blood which is provided with a glass mount that enables one to tell immediately the vein has been entered.

Clinical Memoranda

Ruptured Ectopic Gestation: Implantation in Interstitial Portion of the Tube

The following record of an unusual case of ectopic gestation may be of general interest.

The patient was a married woman of 38 with two children aged 15 years and 8 years respectively. She had a normal menstrual period in February, 1936. In March the period was six days late (normally three weeks and three days). She had normal loss for two days, cessation for five days, except for slight staining. Bleeding per vaginam then returned, associated with the passage of clots for a further seven days, at the end of which time she complained of weakness, pain in the shoulder, but no abdominal pain throughout. She was admitted to hospital on April 14th, fourteen days after the haemorrhage began. On admission the patient was blanched, conscious, but unable to take much notice of her surroundings. Pulse rapid and of poor tension. Examination of the abdomen showed no distension and no tenderness or guarding on palpation. On bimanual examination the uterus was found to be slightly enlarged—no palpable swelling in fornices. A diagnosis of ruptured ectopic gestation was made, largely upon the blanched appearance of the patient being more than could be accounted for by vaginal loss.

Operation.—Ether anaesthesia. During the operation 500 c.cm. citrated blood were given, followed by auto-transfusion from blood in peritoneal cavity. The right cornu of the uterus at the junction with the Fallopian tube was the site of haemorrhage due to rupture of an interstitial pregnancy which had practically severed the Fallopian tube from the uterus, leaving a circular bleeding area about $\frac{3}{4}$ in. in diameter. Bleeding was arrested by inserting two interrupted mattress sutures through the uterus on the proximal side. The Fallopian tube was divided and ligatured on the distal side.

The patient had a remittent temperature for about a week after operation, when the temperature settled down; she had a normal convalescence, and was discharged from hospital in seventeen days.

COMMENTARY

(a) *The Site.*—The interstitial portion of the Fallopian tube is the least likely to be the site of ectopic pregnancy. In this case the pregnancy was most probably utero-tubal, which may account for the long period of haemorrhage before the condition of the patient became urgent.

(b) *The Absence of Pain.*—This was most misleading, and almost confirmed the view that the condition was a uterine abortion. The only pain present had been pain between the shoulder-blades the day before admission. In a recent investigation by Folk and Rosenbloom,¹ pain was absent in only nine out of 320 cases of ectopic gestation. Free fluid in the peritoneal cavity was not determined, but I understand that a negative finding is of no value in these cases.

My thanks are due to Dr. Jeffrey Ramsay for permission to publish this case, and also for the blood examinations, both on admission and before the discharge of the patient.

A. McDOWALL, M.B., Ch.B.Ed., F.R.C.S.

Blackburn Royal Infirmary.

REFERENCE

¹ *Surg., Gynecol. and Obstet.*, February 1st, 1936, p. 228.