Electrical Aids

Electrical aids to hearing may be divided into two groups:—

1. The popular type in which there is a small carbon microphone with a small dry cell and amplifier. The receiver is designed either to wear in or over the ear, or behind the ear for bone conduction.

2. The valve amplifier for increasing the range of the microphone, and a receiver. Modern methods of manufacture allow them to be reduced in size to less than a lady's small handbag. Naturally they are more conspicuous than the ordinary telephone type because the amplifier and microphone have to be carried in the hand instead of being about the patient's person. There is no doubt, however, that the valve type gives better amplification and is comparatively free from distortion and noise.

Electrical aids to hearing have proved in the past a source of income to the manufacturers. Instruments costing about one pound to make have been marketed at fifteen to twenty pounds; the testing of these instruments has been inefficient and without due regard to the individual case. In the past some firms have not been content with Press advertising, but have employed travellers to hawk their wares on the door-step. Thousands of these aids have been sold before the profession, and more especially the otologists, have wakened to the fact that those who are hard of hearing must be protected.

It is essential that the standard of the performance of the electrical aid must be fixed so that a suitable instrument can be prescribed accurately, and, once fitted, properly tested. To-day many firms have adopted this attitude and have fitted up their own testing rooms and there try out various types of instruments on the patient. One interesting sequel of this attitude has been the lowering in price of many of the electrical aids to hearing. It is unfortunate that for business reasons, or for the sake of appearances, an inconspicuous aid is being demanded; and it must be realized that the smaller the aid the less effective it becomes, and this also must hold for bone-conduction aids. The hard of hearing do not want to parade their disability, which is often to their social and economic disadvantage, but if it were realized generally that the use of a suitable aid lessens the hearing strain and thus improves the general outlook and ability, then appearance would become of minor importance. It should be our aim as otologists not only to aid the deaf but to educate the general public in its attitude towards the user of a hearing aid.

Prescribing of Aids

In prescribing an electrical aid it must be recognized that certain hearing defects may possibly be increased by an electrical aid; for example, a mixed middle and inner ear deafness with a marked lowering of the upper tone limit cannot be a suitable case for an electrical aid, which, although it may improve the hearing temporarily, in the long run will increase the nerve defect. Similarly it is inadvisable to prescribe an electrical aid in a case of senile nerve deafness, although a non-electrical aid to hearing would lessen the constant strain. In many cases where there is a marked loss of the cochlear response amplification of the sounds of an electrical aid would only lead to confusion. These few examples go to show that aids to hearing must be accurately prescribed.

The days are now past when the public buy spectacles over the shop counter. The opticians have combined among themselves in the drawing up of a scheme of registration. Although there is no statutory recognition yet the registered optician has had some training and experience in sight testing and has a modicum of know-

ledge of the possible diseases causing defective sight. If any disease is suspected the case is referred to an ophthalmologist.

This scheme has very obvious defects—but a similar scheme applied to the prescribing and testing of aids to hearing would be far superior to the haphazard way it is done at present. Aural clinics should have similar facilities. Patients should not be left to their own devices when obtaining aids to hearing. The common experience of us all is to interview hospital patients already equipped with expensive aids to hearing which on test are absolutely useless. Lectures on diseases of the ear, nose, and throat at present contain no reference to acoustics. The audiometer and the audiograph should be more familiar to all in our specialty. The pure tone method leaves something to be desired as the rectifying property of the ear is left out of consideration—that is, the ability to supply overtones or monotonies which comes into perception of speech. Some method of testing by actual speech should be our aim where not only the loss of pure tones but also the loss of intensity of speech could be measured.

Fortunately, our knowledge on the subject is being advanced daily, so this paper is simply a plea for consideration of the whole question of the demand and the supply for aids to hearing. Now that modern science can supply better and more effective aids these must not be haphazardly foisted on unresponsive ears. I welcome the pamphlets issued for the protection of the deaf by the National Institute for the Deaf, but publications of this description should be given a wider and freer publicity.

AN UNUSUAL CASE OF JAUNDICE IN A MINER

BY

J. A. McKEON, M.B., B.S.,

ROYAL VICTORIA INFIRMARY, NEWCASTLE-UPON-TYNE

AND

H. C. BROWN, C.I.E., M.B., CH.B., D.T.M. AND H.

WELLCOME BUREAU OF SCIENTIFIC RESEARCH, LONDON

The following case illustrates the fact that an isolated positive serological test in the case of a patient suffering from jaundice is not necessarily diagnostic of Weil's disease if there is any chance of agglutinins being present in the blood owing to a previous infection. This is analogous to the significance of the absence of a rise in titre of the serum in typhoid suspects who have received a prophylactic inoculation.

It is now well known that Weil's disease occurs more frequently in certain occupations than in others; for example, in sewer workers (Fairley, 1934; Alston and Brown, 1935; Alston, 1935), fish workers (Davidson *et al.*, 1934), and miners (Buchanan, 1926; Swan and McKeon, 1935) the incidence is distinctly greater than in other occupations in which the risk of exposure to the dejecta of infected rats is less. When, therefore, jaundice occurs in a patient engaged in one of these occupations the probability of it being due to Weil's disease must naturally be considered, and every means in our possession must be brought to bear upon the differential diagnosis from other forms of jaundice.

Diagnostic Criteria

The disease which is most likely to be mistaken for Weil's disease is infectious catarrhal jaundice; when the latter occurs in epidemic form, affecting as it generally does mostly children, no difficulty will be experienced in differentiating it from Weil's disease. When, however,

an adult is attacked by the severe form of this disease characterized by haemorrhages, intense jaundice, albuminuria, and conjunctival congestion the clinical differentiation of the two diseases is by no means easy. It is in these cases that the laboratory findings are of the utmost value.

In the first place, in the absence of any conditions such as a septic focus which would affect the blood picture, the differential blood count is of great importance. In catarrhal jaundice Findlay, Dunlop, and Brown (1931) found a very distinct relative mononuclear increase. A typical count would be: polymorphonuclears 51 per cent., small lymphocytes 41 per cent., large mononuclears 3 per cent., eosinophils 4 per cent., mast cells 1 per cent. In Weil's disease, however, the most striking feature of the differential count is the polymorphonuclear increase, up to 85 or 90 per cent. being by no means uncommon.

When circumstances do not permit of the actual finding and isolation of the leptospira in Weil's disease the serological reactions are very important. Antibodies as a rule appear about the sixth day of the disease, and reach their maximum three or four weeks later. One point which must always be taken into consideration is the fact that agglutinins persist in the patient's serum for many years, and an agglutination titre, even up to 1 in 300 several years after the onset of the disease, is not uncommon (Fairley, 1934; Brown, 1935). This fact, together with the rise in agglutinin titre which occurs during the first fortnight of the disease, must always be considered in making a diagnosis of Weil's disease.

Case Record

The following account of a case of jaundice in a miner whose serum agglutinated *Leptospira icterohaemorrhagiae* to a dilution of 1 in 100 bears out these points.

The patient, aged 52, who had worked in a seam of a colliery in which cases of Weil's disease had occurred, was admitted to hospital on March 20th, 1936, complaining of jaundice, epistaxis, pruritus, dark-coloured urine, and putty-like stools.

The serum gave the following agglutination reactions on three occasions against a London human strain of *Leptospira icterohaemorrhagiae*:

1936	1/10	1/30	1/100	1/300	1/1,000
March 31st	0	+	+	0	0
April 4th	+	+	+	Trace	0
April 14th	+	+	+	0	0

It will thus be seen that there was no appreciable rise in the titre of this serum on successive examinations.

The second point to be noticed is that the titre of the serum is much lower than would be expected to be the case considering that jaundice was well marked eleven days before the first agglutination test.

The following shows a rise in titre in a typical case of Weil's disease:

1935	1/10	1/30	1/100	1/300	1/1,000	1/3,000	1/10,000	1/30,000
Sept. 3rd	+	+	+	0	0	0	0	0
Sept. 4th	+	+	+	Trace	0	0	0	0
Sept. 7th	+	+	+	+	+	0	0	0
Sept. 8th	+	+	+	+	+	0	0	0
Sept. 9th	+	+	+	+	+	+	Trace	0
Sept. 11th	+	+	+	+	+	+	+	0

The second point which we consider important in the diagnosis of the case under consideration is the differential blood count, which was as follows:

	April 1st, 1936	April 14th, 1936
Polymorphonuclears ...	48.5 per cent.	48 per cent.
Lymphocytes ...	39.0	38
Large mononuclears ...	9.0	11
Eosinophils ...	3.5	3

This is typical of the count met with in cases of catarrhal jaundice, and in that it is present in a patient in whom there is no rise in the agglutinin titre against *Leptospira icterohaemorrhagiae* as the disease progresses it leaves very little doubt that the disease in question is catarrhal jaundice, and that the low titre of agglutination is due to a previous infection with the leptospira.

We wish to thank Dr. J. C. Spence for kindly allowing us to publish this case.

BIBLIOGRAPHY

- Fairley, N. H.: *British Medical Journal*, 1934, ii, 10.
 Alston, J. M., et al.: *Lancet*, 1935, i, 806.
 Alston, J. M., and Brown, H. C.: *British Medical Journal*, 1935, ii, 339.
 Davidson, L. S. P., Campbell, R. M., Rae, H. J., and Smith, J.: *Ibid.*, 1934, ii, 1137.
 Buchanan, G.: Medical Research Council, Special Report Series, No. 113, London, 1927.
 Swan, W. G. A., and McKeon, J. A.: *Lancet*, 1935, ii, 570.
 Findlay, G. M., Dunlop, J. L., and Brown, H. C.: *Trans. Roy. Soc. Trop. Med. and Hyg.*, 1931, xxv, 7.
 Brown, H. C.: *British Medical Journal*, 1935, i, 411.

THE OTOLITHIC CATASTROPHE

A NEW SYNDROME

BY

A. TUMARKIN, M.B., F.R.C.S.ED., D.L.O.

HONORARY AURIST, BOOTLE GENERAL HOSPITAL

Ever since Ménière gave his famous account of vertigo and allied phenomena associated with disease of the ear the title of Ménière's syndrome has been used vaguely to include a very heterogeneous collection of conditions. With better understanding of labyrinthine function, however, some order is emerging from this chaos. For instance, a comparatively clear-cut entity is now well recognized in aural vertigo, a condition characterized by:

1. Progressive deafness, usually bilateral but almost invariably much more pronounced in one ear.
2. Attacks of vertigo, with allied symptoms of pallor, nausea, sweating, etc.

It is clear that the syndrome is referable to some degeneration, whether toxic, inflammatory, or abiotrophic, of the inner ear. Whether the lesion is primarily in the neuro-epithelium of the inner ear or, as Dandy has not very convincingly maintained, in the eighth nerve itself is not yet proven. That question, while important from the point of view of surgical intervention, will not influence the present argument.

Anatomy of the Inner Ear

The inner ear can be divided both anatomically and physiologically into:

1. The cochlea.
2. The semicircular canals.
3. The vestibule, further subdivided into: (a) utricle, (b) saccule.

It will be noted immediately that although the organ involved in aural vertigo has a triple, if not quadruple, nature yet the symptoms evoked from it are classified under only two headings. The progressive deafness is referred to the cochlea, and the vertigo, etc., to the canals. This discrepancy is due to the fact that until comparatively recently very little was known of the