widest part of the forearm. There was no pitting. The axillary vein was still palpable and cord-like. The superficial veins were more prominent on the right shoulder than on the left.

Summary

The aetiology of primary thrombosis of the axillary vein is discussed. One case is described, and in the appended bibliography twenty-three others are collected since the exhaustive study made in 1928 by Pearce Gould and Patey on eight of their own cases and on the thirty-one other cases reported in the literature previous to this. It is suggested that pressure on the axillary vein by constriction of a pyjama sleeve at night may cause the onset of the thrombosis in certain cases.

We are indebted to Professor H. Collinson for permission to publish this case, and to Mr. E. Pearce Gould for his kind advice.

BIBLIOGRAPHY

- Paget, J.: *Clinical Lectures and Essays*, 1875, p. 306.
 Ruge: *Med. Klinik*, 1911, vii, 1615.
 Schepelmann: *Behefte z. Med. Klin.*, 1911, vii, 23.
 Rosenthal, W. J.: *Deut. Zeit. f. Chir.*, 1912, cxvii, 405.
 Willan, R. J.: *Edinburgh Med. Journ.*, 1918, xx, 105.
 Cadenat, F. M.: *Paris Méd.*, 1920, xxxv, 253.
 Pearce Gould, E., and Patey, D. H.: *Brit. Journ. Surg.*, 1928, xvi, 208.
Cases since 1928
 Löhr, M.: *Deut. Zeit. f. Chir.*, 1929, ccxiv, 263.
 Oehme, C.: *Med. Klinik*, 1930, xxvi, 472.
 Kinzel, H.: *Beitr. z. klin. Chir.*, 1930, cl, 252.
 Horton, B. T.: *Journ. Amer. Med. Assoc.*, 1931, xcvi, 2194.
 Clute, H. M.: *Surg. Clin. North America*, 1931, xi, 253.
 Joskisch, G.: *Beitr. z. klin. Chir.*, 1931, clii, 453.
 Jakobson, L.: *Munch. med. Woch.*, 1931, lxxviii, 1017.
 Goudsmit, J.: *Nederl. Tijdschr. v. Geneesk.*, 1931, lxxv, 5480.
 Thomsen, E.: *Ugeskrift for Læger*, 1932, xciv, 810.
 Firth, D., and Mackay, R.: *Lancet*, 1932, ii, 679.
 Ross, J. C.: *British Medical Journal*, 1933, ii, 525.
 "F.R.C.S.": *Ibid.*, 1933, ii, 844.
 Taylor, C. H. S.: *Ibid.*, 1933, ii, 818.
 McIntosh, J. N.: *Ibid.*, 1933, ii, 1144.

ACETYLCHOLINE IN TOBACCO AMBLYOPIA

BY

H. CAMPBELL ORR, M.B., F.R.F.P.S.

HONORARY SURGEON, WOLVERHAMPTON AND MIDLAND COUNTIES
EYE INFIRMARY

In 1901 Parsons,¹ as a result of experimental work on rabbits, suggested that the defect in tobacco amblyopia was due in part to two causes: (1) constriction of the arterioles, which resulted in the starvation of the macular area, where the blood supply is poor; and (2) paralysis of the synapses in the cone fibres, or of the bipolars, or of both. For many years opinion favoured the degeneration of the nervous elements as the primary cause of the symptoms.² De Schweinitz stated: "The situation of the pathological process is in the optic nerve, and especially in that portion which is known as the papillo-macular bundle," but he added that the appearances were more like those of an interstitial sclerosing inflammation with secondary nerve atrophy than primary nerve atrophy with secondary interstitial change.

More recently some observers have inclined towards the opinion that the constriction of the arterioles from nicotine poisoning was the primary cause, and have based their therapeutic measures accordingly. In 1930 Pfimlin³ reported fifteen cases treated with sodium nitrite in isotonic sodium chloride solution injected intravenously, and 50 per cent. of the eyes obtained 20/30 vision in five and two-tenths weeks. In 1932 Duggan treated six cases with sodium nitrite intravenously, and eight out of twelve eyes obtained 20/30 in three to fourteen days.

In 1935 Duggan⁴ reported a series of twenty-four cases of toxic amblyopia due to excessive smoking, which he treated with a form of sodium nitrite known as nitroscleran, and reported marked improvement in the visual acuity, with corresponding reduction in the central scotoma. Twenty-one of these patients improved to 20/30 in eighteen and four-tenths days. Five of the patients actually continued smoking during treatment, but their average time of recovery to 20/30 was delayed to thirty-six days. Where the patients did not smoke recovery was completed in twelve days. These figures compare favourably with the older forms of treatment. Fuchs found the period before recovery in very mild cases to be from one to two months; Traquair cured mild cases in about six weeks, but reported that in severe cases two or three months might elapse before definite improvement began.

The following cases have been treated with acetylcholine, and this note is published so that other observers may try it and report results. The condition is becoming less frequent, and to wait for a larger number of cases

may deprive many patients in other places of the advantages of this method. All the patients were advised to stop smoking, but abstinence was not complete.

A gate porter aged 63 attended the out-patient department of a hospital on October 24th, 1934, complaining of failing vision. The right vision was 1/60, with correcting glasses 6/60; the left vision was 1/60, c. gl. 6/60. He was treated with potassium iodide until November, 1934, when he stopped attending since he had made no improvement.

I saw him through the National Eye Service on July 15th, 1935, and diagnosed tobacco amblyopia. Vision was then with glasses 1/60 in each eye, with total loss of colour vision; 0.1 gram dose of acetylcholine was injected intramuscularly, and the patient was admitted to hospital on the following day, when the vision had improved to 4/60 in each eye. He was given intramuscularly 0.1 gram of acetylcholine (B.D.H.) daily, and after the third injection the vision with glasses was in the right eye 6/36 and in the left eye 6/24 L 1. On the 19th he was discharged with right vision 6/18 partly and left vision 6/18 partly. He attended daily for his injection, and resumed his work. On the fifth day he could recognize blue and yellow, but not red or green. On the tenth day the vision in each eye was 6/18, and this was maintained until August 13th, when the vision in each eye was c. gl. 6/12; he could identify red and green, and there was no sign of any scotoma. The acetylcholine was stopped, and he was given a mixture containing potassium iodide, potassium acetate, and sodium citrate, 5 grains of each to be taken three times daily. On August 27th the vision with glasses was 6/9 and J 1 in each eye, with perfect field of vision and colours.

A carpet weaver was seen by me through the National Eye Service on June 26th, 1935, complaining of failing vision for two years. The vision in each eye was 1/60. Both disks looked pale and atrophic, and there was a scotoma for all colours. He was given an injection of 0.2 gram of acetylcholine, and was admitted to hospital. On the 27th he received 1 c.cm. acecholin (Anglo-French Drug Co.), and the vision was then: right 6/36 (L 1) and left 6/18. The blood pressure was 160 mm. before the injection, and 140 mm. half an hour afterwards. The injection was repeated each day, and on July 2nd the vision was 6/12 in each eye. He was discharged on July 5th with vision 6/9 in each eye and a good colour response. He was given a mixture of potassium iodide with potassium acetate and sodium citrate, 5 grains of each to be taken three times daily, and attended at the out-patient department three days later for an injection. The vision in the right eye had dropped to 6/18, but the left remained at 6/9. Thereafter he had one injection each week, and admitted having had an occasional smoke. On July 30th he could see his wools at work, and the vision was: right 6/12 and left 6/6. Tested with Ishihara's test on September 24th, he showed some red-green blindness.

A man aged 60 was seen as an out-patient on July 23rd complaining of failing vision for a long time. The vision in each eye was 4/60. The blood pressure was 220/110. The right eye identified yellow only and the left eye saw it as

"brassy colour." The disks looked pale and the arteries sclerotic. There was an absolute scotoma for colour in the right eye, and a central scotoma for colour in the left eye. He was given 0.5 c.cm. acecholin intramuscularly each day, and on August 2nd there was a reduction in the scotoma in each eye, with vision 6/60. Thereafter he received 0.1 gram acetylcholine every second day, and on September 2nd colours were recognized centrally. From that date his attendance averaged once a week, when he received 0.1 gram intramuscularly. On October 10th there was no scotoma, and with correcting glasses he could see 6/12 with each eye. On the first day this patient had 5 minims of acetylcholine injected subconjunctivally without showing any improvement.

A patient aged 72 was sent to me by Dr. A. V. Russell with a history of recent illness and a very high blood pressure. He complained that after his illness his vision failed and colours were blurred. The vision in each eye with glasses was 6/12, and in each eye there was a central scotoma for colours. The fundus vessels were fine and sclerotic. Dr. Russell injected 0.1 gram of acetylcholine each day for six days, when the patient was seen again. He smoked only a few cigarettes in the interval. On the eighth day he could recognize all colours except blue centrally, and the vision with glasses was now 6/6. The acetylcholine was stopped. When seen on the twenty-first day all colours were recognized centrally and the vision was unchanged.

Summary

The first three cases may be termed chronic, and show progress as follows.

Case 1.—On the eleventh day vision was c. gl. 6/18, and the patient resumed work. On the nineteenth day vision was c. gl. 6/12, and on the twenty-eighth c. gl. 6/12, all colours being identified.

Case 2.—On the sixth day vision was 6/12 and the patient fit to resume work. On the eighth day vision was 6/12 (colours good), and on the ninth 6/9.

Case 3.—On the fortieth day vision was 6/36 without glasses: colours identified. On the forty-sixth day with 1.5 dioptries correction the vision was 6/12.

Case 4.—In this acute case on the seventh day the vision with glasses was normal, and all colours except blue were identified centrally.

Conclusions

The first three cases form a very severe test for any therapeutic agent in this condition, and I think it must be agreed that the results are exceedingly good. The result in the fourth case is also very satisfactory. If these results can be produced by other observers this is shown to be the best therapeutic agent we have at present for the treatment of tobacco amblyopia. I have also been using acetylcholine in the treatment of retinitis pigmentosa, and hope at a later date to publish some notes on cases.

REFERENCES

- ¹ *Ophth. Review*, 1901, xx, 189.
- ² *Trans. Amer. Ophth. Soc.*, 1897-8, viii, 186.
- ³ Pfimlin: *Klin. Monatsbl. f. Augenh.*, 1930, lxxxv, 787.
- ⁴ *Arch. of Ophthalmol.*, 1935, xiii, 1059.

The Fédération des Unions Hospitalières de France has given place to a new organization—the Fédération Hospitalière de France. The original society, which had been in existence since 1924, was not completely national in character, excluding, as it did, Paris and the Paris region. The new society has a council of thirty-five members, five from each regional association and five from the Paris Public Assistance Board. The Federation, whose offices are at 4, Avenue Victoria, Paris, publishes every month *La Revue Hospitalière de France*. We are informed that the Minister for Public Health has promised his official patronage and the co-operation of the Ministry at the fifth International Hospital Congress, to be held in Paris in 1937. An organizing committee is to be nominated, including Dr. René Sand, M. Sarraz-Bournet, and Dr. Alter as liaison officers to the International Hospital Association.

Clinical Memoranda

Prophylaxis in Puerperal Sepsis and Pyrexia

In making a further contribution in regard to the use of calcium sulphide as a prophylactic against puerperal sepsis¹ I am now reviewing the cases that have been treated from January 1st, 1935, to April 16th, 1936. There have been admitted to this institution during that time approximately 1,850 cases for confinement and 230 cases of abortion, a total of 2,080, and my results are even better to-day than they have been at any time since calcium sulphide was used during the fourteen months previous to this review.

In commencing the use of calcium sulphide I was inclined not to overstep the pharmacopoeial dose of 1/2 grain twice or thrice daily, because there was always the possibility that with a new venture of this sort something untoward might happen to either the mother or the baby. To-day the dose of 1/2 grain is replaced by doses of 3 grains twice daily, in either pill or capsule form, preferably in pill form, because in a capsule the calcium sulphide has to be suspended in oil—the oil sometimes causing a degree of nausea. In complicated cases, such as the toxæmias of pregnancy, haemorrhage, lacerations of the soft parts, and in cases of delayed labour where a degree of shock exists, 3 grains are given four times daily.

During the period under review there have been notified to the local supervising authority fourteen cases of puerperal sepsis—all infected before admission—and recovery has occurred in every case, with one exception—namely, a case of severe pyelitis, eclampsia, and delivery by craniotomy (already referred to in my previous communication)—giving an incidence rate of 6.8 per 1,000, while other cases recorded as "puerperal pyrexia" numbered eleven in all, giving a total notifiable rate of 11.8 per 1,000, with a death rate of less than 0.5 per 1,000.

It is interesting to note that the cases of puerperal sepsis were of a mild or moderate nature, with one exception. I can also add 1,000 cases occurring consecutively in the previous year (1934), and that added to the present series would give a maternal death rate of 0.3 per 1,000, whereas the previous death rate over a period of seven years (1927 to 1933) was 5.3 per 1,000.

During the previous seven years the same precautions were taken, with the additional use of (a) puerperal anti-toxin; (b) vitamins A and D, in the form of adexolin; (c) swabs soaked in acriflavine, 1 in 750 aqueous solution, per vaginam at the beginning of the first stage and repeated during the second stage. Those forms of treatment were costly, and for antitoxin alone up to £3,300 in the years 1932-3 was spent for nearly 1,000 cases of confinement, while the cost involved for 1,500 confinements using calcium sulphide prophylactically would not exceed £100 per annum.

It may interest the reader to know that cases admitted to this institution are not selected cases; 60 per cent. of all cases are "booked" and have ante-natal supervision, but 40 per cent. are emergency cases. Lanarkshire probably has a greater percentage of complicated confinements than any area in Great Britain, and the fact that the local authority's area is one of the districts stricken with unemployment tends to handicap the work, and the malnutrition consequent on prolonged unemployment renders the patients more susceptible to puerperal infections.

¹ *British Medical Journal*, 1935, ii, 1070.