second lumbar ganglia, the coeliac ganglion, and the three splanchnic nerves causes severe pain from the shoulder to the lower abdomen, the pain disappearing after injection of the splanchnic nerves with procaine. These physiological experiments on human beings are difficult to understand unless the sympathetic has afferent functions.

2. Excessive heat stimulation of anaesthetic trigeminal areas may produce in a few seconds a stinging sensation, which is abolished after stellate ganglionectomy, a fact strongly suggestive of sympathetic afferent conduction of pain. In exceptional cases deep sensibility may return and become hyperaesthetic with spontaneous crises of pain after total trigeminal sensory root resection, even in the presence of total facial palsy. This appears clear evidence of sympathetic afferent function becoming active, though normally latent.

3. A form of neuralgic hemicrania appears to be connected with sympathetic disturbance and may be curable by stellate ganglionectomy. A preliminary test should be applied by injecting 10 c.cm. of 1 per cent. procaine around the ganglion from behind the first rib.

4. Migrainous neuralgia appears to be due to a sympathetic disturbance in part of some of the branches of the external carotid, the middle meningeal, and temporal arteries, the pain being due to vasomotor dilatation effects in the meninges. The afferent path is the recurrent meningeal branches of the first and second divisions of the trigeminal nerve, and the painful attacks may be curable by partial Gasserian block or root resection. This may be compared with the headache of true ophthalmic migraine, which is probably of cortical origin and associated with vaso-dilatation of the posterior cerebral arterial branches.

REFERENCES

- ¹ Carmichael and Woollard: *Brain*, 1933, lvi, 109.
- ² Frazier: *Amer. Journ. Med. Sci.*, 1925, clxix, 469.
- ³ Max Peet: *Arch. Neurol. and Psychiatry*, 1929, xxii, 313.
- ⁴ Davis and Pollock: *Ibid.*, 1932, xxvii, 282.
- ⁵ Foerster, Altenburger, and Kroll: *Zeit. f. d. ges. Neurol. u. Psych.*, 1929, cxxi, 139.
- ⁶ Adson: *Proc. Staff Meetings of the Mayo Clinic*, 1935, x, 623.
- ⁷ Tinel, M.: *Rev. Neurol.*, 1930, xxxvii, 1, 765.
- ⁸ Ivy and Johnson: *Univ. Penn. Med. Bull.*, 1907.
- ⁹ Harris: *Neuritis and Neuralgia*, 1926, p. 241.
- ¹⁰ Helson and Frazier: *Brain*, 1932, lv, 114.
- ¹¹ Craig: *Proc. Staff Meetings of the Mayo Clinic*, 1935, x, 362.
- ¹² Alajouanine and Thurel: *Journ. Med. Franç.*, 1933, xxii, 204.

TREATMENT OF HAEMORRHAGE

BY

IAN J. WOOD, M.D., M.R.C.P.

MARION CARTY RESEARCH FELLOW, WALTER AND ELIZA HALL INSTITUTE OF MEDICAL RESEARCH, MELBOURNE

The recent introduction by Marriott and Kekwick^{1,2} of continuous intravenous blood transfusion has provided a new weapon for the treatment of haemorrhage. The following contribution is based on a study which has been made during the past two years of the clinical and biochemical changes that are associated with varying degrees of exsanguination and the effects of transfusion of blood and glucose-saline. Patients with haematemesis and melaena have been selected, and although there is as yet not sufficient evidence to show the indications for operation in these cases it is felt that a strong plea may be advanced for continuous blood transfusion as a valuable aid in treatment. The following fragmentary observations are placed on record in the hope that they may help other workers in this field or perhaps provoke their criticisms.

Normal Blood Volume

The normal blood volume for a man of 64 kilos (10 st.) is approximately six litres (ten pints). When a haemorrhage takes place the amount lost is replaced by the tissue fluids, the rapidity of replacement depending on the ratio of the volume of available tissue fluid to the volume of blood which is lost. Small amounts, such as 500 to 1,000 c.cm., are rapidly replaced, but if more than one-half the normal blood volume is removed by haemorrhage there may not be adequate replacement for many hours.³ It is a common practice to limit the fluid intake by mouth in all cases of haematemesis, and this tends further to reduce the volume of reserve fluid available for replacement.

An estimation of the percentage of haemoglobin made during the phase of depleted blood volume is not a true index of the total amount of circulating haemoglobin; a true estimate of the loss cannot be made by haemoglobin estimation until the volume of blood has been fully restored. The importance of the reduction in blood volume during haemorrhage and its subsequent replacement is fully discussed in the reports of the Medical Research Council on transfusion during the great war.³ The blood plasma normally contains over 6 per cent. of

protein, so during haemorrhage there is a loss of protein as well as of haemoglobin. Whipple and his co-workers⁴ have shown that so long as there has been no preliminary starvation a reserve of protein is available in the body which can be called upon to restore the plasma proteins.

During rapid haemorrhage and immediately afterwards, there is a profound fall in blood pressure, owing to the depleted blood volume and a lag in the vaso-constrictor response. The pressure then rises to within 15 to 30 mm. of the level usual for the individual. This rise is due to widespread vasomotor constriction, in which the carotid sinus and aortic reflexes play a major part; the blood volume is still reduced and the percentage haemoglobin relatively high. Finally the blood volume is restored by the tissue fluids, the blood pressure still remaining at a moderately high level.⁵ This sustained blood pressure was a consistent finding in the majority of cases in the present series. During rapid blood loss a systolic blood pressure of about 100 mm. was usually maintained until the patient was *in extremis*. The various reserve mechanisms then failed. The blood pressure is thus a poor guide to the degree of blood loss. The pulse rate is elevated in every phase, being rapid and "thready" in the initial stage of rapid blood loss and, later, with the adjustments in blood pressure, the volume improves, but the rate remains between 110 and 140; with further haemorrhage the rate increases before there is a pronounced fall in the blood pressure. The pulse is therefore more informative than the blood pressure when assessing the degree of haemorrhage. However, an accurate estimation of the haemoglobin percentage is by far the most reliable single clinical test, and it should be frequently repeated if a concealed haemorrhage is suspected. To determine the blood volume as well as the haemoglobin would be ideal, but this is not practicable as a routine in clinical work.

Terminal Stages

As a result of the low haemoglobin and impaired circulation every organ is inadequately supplied with oxygen, with consequent impairment of functional efficiency, unless the reserve forces within the body are sufficient for readjustment. If the demand on the reserves is too great they will fail, and unless therapeutic aid is provided death will ensue. In the final stages, the failure of the circulation, the poor oxygen-carrying capacity of the blood and probably the depletion of plasma proteins contribute to a widespread oedema. Evidence of this is frequently seen

in the lungs—crepitations, dullness, and poor air entry at the bases being a common finding. Moreover, it is of interest to note that it has been shown by many observers that quite extensive oedema of the tissues must be present before it can be recognized clinically. Wright-Smith stated in a personal communication that oedema of the soft tissues, especially of the lung, is invariably a prominent finding at necropsy after death from haemorrhage.

Changes in Blood Urea

In studying the effects of these widespread changes special attention has been paid to the level of the blood urea during the various phases of exsanguination and repair. Although similar studies on the level of the blood urea have been previously recorded by Christianson,⁵ Ingegno,⁶ and others, there are no available reports on the effects of severe haematemesis and the administration of large volumes of blood. The rise in blood urea following haematemesis is illustrated by the following case history.

CASE I

A male aged 54 was admitted to the wards of the Royal Melbourne Hospital on October 7th, 1935. Ten years before he had vomited several pints of blood. Since then he had suffered from intermittent attacks of indigestion, epigastric pain occurring half an hour after meals; later the pain was noticed in the back and was more continuous. Twenty-four hours prior to admission he vomited about a litre of dark blood and passed three tarry motions.

Examination showed a pale, thin man. The pulse rate was 92, systolic blood pressure 80 mm., diastolic 60 mm., haemoglobin 34 per cent., and blood urea 88 mg. per 100 c.cm.

He was given saline per rectum, and only ice was allowed to be taken by mouth. Morphine was given in liberal quantities. The following day (October 8th) he appeared to be worse. He was paler, the systolic blood pressure was 90 mm., the pulse rate ranged from 100 to 120, the haemoglobin had fallen to 30 per cent., and the blood urea had risen to 123 mg. per 100 c.cm.: 600 c.cm. (one pint) of citrated blood was given without producing any definite improvement in his condition. On the third day after admission (October 9th) he passed a large quantity of blood per rectum, and gradually sank into a moribund state. The pulse rate rose to 140 and then became imperceptible, the systolic blood pressure fell to 55 mm., the blood urea rose to 145 mg. per 100 c.cm., and the haemoglobin was 28 per cent. Several hours later he died.

FIG. 1.—Case I. A rapid haematemesis. 600 c.cm. of citrated blood was given.

Post-mortem examination showed a chronic ulcer on the posterior surface of the stomach, opposite the centre of the lesser curvature. An eroded artery with thickened walls protruded from the ulcer bed. There was widespread oedema of both lungs and general pallor of the organs. The kidneys appeared normal to macroscopic examination.

Hypothesis of Renal Mechanism

In the present series it was found that the blood urea rose progressively with rapid severe exsanguination and slowly fell when the anoxaemia, impaired circulation, and other stresses on the kidney were relieved. The exact mechanism of normal kidney function is not fully understood, but the following hypothesis, based on the views

of van Slyke and his co-workers, affords an explanation of the changes which have been observed in the present series. When the initial fall in blood pressure occurs, both during and immediately after the sudden blood loss, the renal excretion is reduced. Complete anuria appears when the blood pressure falls below 40 mm., because a positive pressure is necessary to force fluid from the glomerular tufts into Bowman's capsule against the osmotic pressure of the plasma proteins. After a few hours the stage of profound circulatory failure passes and the blood pressure rises, owing to a widespread vasomotor spasm. Should the glomerular capillaries or the afferent arteries and arterioles participate excessively in this constriction the excreting field of the kidney is depleted and the blood urea rises. A third and more lasting factor is the effect of anoxaemia on the kidney cells. Although 97 per cent. of the water excreted by the glomerulus is reabsorbed in the tubule, only about 30 per cent. of the urea is returned to the circulation. If the function of the cells in the tubules is depressed by a deficiency in the oxygen supply it is likely that their power of selective reabsorption may be correspondingly impaired and that an excessive amount of urea may pass back into the blood stream.

All the patients who were admitted within thirty-six hours of a severe haemorrhage had high levels of blood urea, figures ranging from 60 to 140 mg. per 100 c.cm. being common; and, moreover, a sudden rise in the level has drawn attention to a further haemorrhage (see Case V). The blood urea usually remained high for one to three days and then slowly fell to 10 to 30 mg. per 100 c.cm. above normal, provided the haemorrhage had ceased and the critical level for recovery was not exceeded. Often the normal urea level was not reached till the haemoglobin had risen above 50 per cent., but occasionally this occurred at a lower haemoglobin percentage.

Stages of Exsanguination

In assessing the value of treatment it is important to consider the extent and rapidity of the haemorrhage, and also the prognosis should no treatment be instituted. The effects of slow exsanguination extending over a period of several weeks will be discussed separately at the end of this paper. With rapid blood loss three stages may be arbitrarily defined:

Stage 1: Moderate Exsanguination.—Haemoglobin 50 to 60 per cent. Slow but progressive recovery occurs.

Stage 2: Critical Exsanguination.—Haemoglobin 35 to 50 per cent. Severe disorder of bodily function is present, but there is a tendency to spontaneous recovery after a period of readjustment if there is no further haemorrhage.

Stage 3: Terminal Stage of Exsanguination.—Haemoglobin under 35 per cent. The power to readjust usually fails, and death ensues unless therapeutic aid is rapidly applied. This will fail if the damage is severe.

The duration of these stages, especially stages 2 and 3, is important. Prolonged anoxaemia probably causes irreparable damage to vitally important cells. It should be our object to restore blood volume and haemoglobin to a normal level as rapidly as possible, so as to allow the debilitated tissues to recover. It is important to realize that if there are other factors present which have reduced the functional efficiency of the various organs then the prognosis in each stage becomes correspondingly worse. Thus haematemesis often arises in subjects who cannot respond in a normal manner to the emergency of a severe haemorrhage owing to vascular degeneration or chronic kidney disease. A further danger lies in the susceptibility of the oedematous tissues to bacterial invasion, the lungs being especially vulnerable. When the bleeding vessel is not ligated a further haemorrhage may

occur which readily precipitates a patient from Stage 1 or 2 to Stage 3. If the haemoglobin is in the region of 35 per cent. the patient is in grave danger, for a further haemorrhage (not a rare event) will usually be fatal. Irreparable damage will be done to the vital centres before a blood transfusion can be administered. Should the untransfused patient recover the rise in haemoglobin is slow, many weeks elapsing before a high level is regained.

Glucose-Saline Transfusion

At the beginning of the present investigation all cases in Stage 2 or 3, and some in Stage 1, were given large volumes of glucose (2 to 5 per cent.) and normal saline intravenously by the continuous drip method. The fluid was allowed to run in at the rate of four to five ounces an hour, and the administration was continued for one to three days. A moderate improvement was noted in most of the cases. Thirst, which is always a distressing symptom, was greatly relieved, although it was not entirely appeased till fluids were given by mouth. The tongue became less parched and brown, and a smaller amount of sedative was necessary to quell the restlessness, which is always a marked feature in these cases. Besides affording an isotonic solution, the sodium chloride helped to restore the salt loss resulting from the haemorrhage, and the glucose assisted in tissue nutrition. It has been shown,⁸ however, that glucose and sodium chloride both diffuse rapidly through the capillary walls, and glucose-saline solutions are therefore very inferior to citrated blood in maintaining the restored blood volume after a large haemorrhage. If given in excess they would contribute to the oedema.

Following glucose-saline transfusion the blood pressure rose slightly and the pulse improved in volume, but there was only a slight fall in the rate. The patients in Stages 2 and 3 usually remained drowsy and stuporose, taking but little interest in anything except the urgent call for fluids to quench their thirst. The duration of the profound anoxaemia appeared to be important in determining the degree of mental confusion, for immediately after the haemorrhage the patient was often quite rational, whereas two hours later, when the circulation appeared to be partially recovered, there was great mental confusion and torpor. It is probable that cerebral oedema played a part in the mental changes, and it is interesting to note that it is a time-honoured custom in the Royal Melbourne Hospital, and probably in many other institutions, to nurse all cases of severe haematemesis flat in bed, with the foot of the bed elevated in order to "help the cerebral circulation." The head thus becomes the most dependent part, and impairment of the venous return may contribute to the development of cerebral oedema. It was found that these patients were far happier if their heads were elevated with one or two pillows. The breath had a heavy abdominal or even urinary odour, and there was a typical restlessness. Crepitations sometimes appeared in the lungs, and the respirations became rapid and laboured.

Haemoglobin Variations

As the transfusion of saline and glucose proceeded there was no rise, but occasionally a slight fall, in the percentage of haemoglobin, and it remained at a low level for a week or more. The return of the haemoglobin to normal, or to levels at which a further moderate haemorrhage would not be a serious risk, took at least two or three weeks. Occult blood often persisted in the stools, showing the presence of an ooze from the damaged vessel which retarded the rise in haemoglobin. The blood urea usually remained elevated for several days, and then, in cases which recovered, slowly fell. In the fatal cases a further rise took place prior to death. Finally, there was

a marked tendency for the haemorrhage to recur unless the vessel was ligated surgically. This may have been partly due to the poor state of oxygenation at the site of the bleeding and consequent delay in organization of the clot. It was two or three weeks before the patients in Stages 2 or 3 were in a suitable condition to undergo an extensive surgical operation such as is usually necessary for the surgical relief and cure of a peptic ulcer after haemorrhage.

CASE II

A male aged 58 was admitted to the Royal Melbourne Hospital on November 3rd, 1934, under the care of Dr. H. Turnbull. He had suffered from flatulent dyspepsia for many years, both the gall-bladder and the appendix having been removed without any benefit. Twelve hours before admission to hospital he fainted, remaining unconscious for several minutes, and later vomited over a litre of blood.

Examination showed a very pale, obese man; pulse rate 88 a minute; systolic blood pressure 120 mm.; haemoglobin 42 per cent.; blood urea 88 mg. per 100 c.cm. On the following day he appeared to be improving, but twenty-four hours later (November 5th) he grew paler, and vomited about half a litre of blood. The haemoglobin fell to 31 per cent. He complained of thirst, and was slightly restless. There was little mental confusion. A continuous drip transfusion

FIG. 2.—Case II. Effect of transfusion of 7,200 c.cm. glucose-saline and 900 c.cm. citrated blood after haematemesis due to cirrhosis of the liver.

of glucose-saline was begun, and during the following forty-eight hours 7,200 c.cm. (twelve pints) of glucose-saline were given. In addition, two small transfusions of citrated blood (total 900 c.cm.) were given at the twelfth and thirty-sixth hours. The blood urea fell progressively from 88 to 49 mg. per 100 c.cm. during this period, but the haemoglobin remained at 32 per cent. After the glucose-saline solution began to run into the veins the thirst was partly relieved, but after twelve hours the restlessness became marked, and he gradually drifted into a state of extreme mental confusion. The abdomen was distended and crepitations appeared at the bases of the lungs. The temperature was elevated to 102° F., and "bronchopneumonia" was diagnosed by the house-physician. However, a rapid resolution of all the distressing symptoms began as soon as the administration of the glucose-saline solution was discontinued. From then onwards he made a very slow but uninterrupted recovery. Fourteen days later the haemoglobin had only risen to 39 per cent. A diagnosis of cirrhosis of the liver was made, an x-ray examination showing no abnormality in the stomach or duodenum.

Massive Blood Transfusion

At this juncture the contribution of Marriott and Kekwich¹ appeared, and their methods were adopted, with minor modifications in the technique of administration. Citrated blood was given at the rate of 90 to 150 c.cm. (3 to 5 ounces) an hour and the haemoglobin was slowly elevated to over 70 per cent. in twelve to forty-eight hours. After the transfusion was finished the haemoglobin sometimes tended to fall slightly, but finally rose towards normal. The dramatic changes in patients during the continuous transfusion of blood have been described by Marriott and Kekwich, and the present study confirms their reports. Thus it was usual to find that a

patient who twenty-four hours previously had lain moribund and deathly pale became pink, and could no longer be distinguished by his colour from the other occupants of the ward. The drowsiness and mental confusion usually passed away, but occasionally persisted for two or three days. The tissues were recovering from the hours of stress. The pulse volume improved rapidly and the rate diminished, although it usually remained for several days in the neighbourhood of 90 to 110 a minute. The blood pressure rose, but for the first week it often remained about 20 mm. below the normal level for the individual. There was no tendency for oedema to appear, as was seen in the cases treated with saline and glucose intravenously, or where no intravenous therapy was given. The protein in the transfused blood draws fluid into the blood stream and thus aids the circulation.

A marked improvement was observed in the level of the blood urea. It usually began to fall in less than twelve hours after the blood began to run into the vein. Sometimes there was a lag of twelve to twenty-four hours before the fall appeared. In the limited number of cases in this series the rate, extent, and early onset of the fall contrasted favourably with cases treated with saline and glucose intravenously or where no intravenous injection was given.

It has often been stated that the transfusion of blood increases the liability to haemorrhage because it raises the blood pressure. While the author is not prepared to say that the rapid transfusion of blood may not open a vessel, slow transfusion by the continuous drip method does not appear to increase the liability to haemorrhage; in fact, the possibility of haemorrhage seems to be more remote. Gordon-Taylor⁹ has recently made a comprehensive survey of the mortality from haematemesis, and has drawn attention to the rapidity with which patients with haematemesis can be prepared for surgical operation by the slow transfusion of blood. In two or three days a moribund patient may be transformed into a good surgical risk; a gastro-enterostomy or gastrectomy can then be performed with comparative safety. Should no transfusion be given it would take at least a month for such a satisfactory stage to be reached. If an operation is performed immediately after the haemorrhage and before any blood transfusion has been given then the danger to life is grave. The patient is drowsy and stuporose, the haemoglobin is low, the blood urea is high, and he is therefore not a suitable subject for a major operation.

CASE III

A male aged 23 was admitted to the Royal Melbourne Hospital on November 15th, 1935, under the care of Dr. H. Turnbull. For five years prior to admission he had suffered from mid-epigastric pains at an indefinite time after meals, often awakening him from sleep in the early hours of the

FIG. 3.—Case III. Beneficial effect of massive blood transfusion (3,900 c.c.m. by continuous drip).

morning. Twenty-four hours prior to admission he had experienced a sensation of weakness and nausea, and he fainted on two occasions. Later he vomited 300 c.c.m. (half a pint) of blood and passed a tarry motion.

Examination showed a very pale, drowsy, and restless young man. His tongue was dry and furred, and his breath had an abdominal odour. The systolic blood pressure was 100 mm.; pulse rate 120; haemoglobin 40 per cent.; and blood urea 90 mg. per 100 c.c.m.; 3,900 c.c.m. (6½ pints) of citrated blood were given by continuous drip transfusion, and the haemoglobin was elevated to 68 per cent. in forty hours. During this period the blood urea fell slightly (90 to 86 mg. per 100 c.c.m.) but sixteen hours later it had fallen to 66 mg. per 100 c.c.m. It took five more days to return to the normal level of 40 mg. per 100 c.c.m. Laparotomy was performed under local anaesthesia by Mr. G. R. A. Syme thirty-six hours after the drip transfusion had begun. A chronic ulcer was found on the anterior surface of the duodenum, and this was infolded. He made a rapid and uninterrupted recovery, and was enjoying good health when seen five months later.

CASE IV

A female aged 73 was admitted to the Royal Melbourne Hospital on February 1st, 1936, under my care. Twelve years previously she had suffered from indigestion. Investigation then showed an ulcer situated in the centre of the lesser curvature of the stomach. A suitable diet and alkaline powders were ordered, and she enjoyed moderately good health with only an occasional attack of indigestion. However, six

FIG. 4.—Case IV. Effect on the blood urea of severe haematemesis and transfusion of 2,700 c.c.m. citrated blood and 3,000 c.c.m. glucose-saline. The second rise in the blood urea is difficult to interpret.

weeks prior to her admission to hospital she began to experience a stabbing pain in the epigastrium about fifteen minutes after taking food. Twenty-four hours before admission she vomited about 600 c.c.m. (1 pint) of dark blood and fainted for about twenty minutes.

On examination when admitted to hospital on February 1st she was moderately pale, but not drowsy or suffering from mental confusion. The pulse rate was 120; systolic blood pressure 120 mm.; haemoglobin 52 per cent.; blood urea 85 mg. per 100 c.c.m. She was given saline, glucose, and alkalis by mouth, and morphine was administered. Two days later she had a severe haematemesis, more than two litres (over 3 pints) of dark blood being vomited within three hours. She then lost consciousness. Soon after this haemorrhage began the blood urea had risen to 127 mg. per 100 c.c.m., and the haemoglobin had fallen to 43 per cent., but probably the haemoglobin eventually reached a level of less than 35 per cent. A continuous drip transfusion of citrated blood was then begun and she slowly improved; 2,700 c.c.m. (4½ pints) of blood and 3,000 c.c.m. (5 pints) of glucose-saline solution were given in thirty-six hours. The haemoglobin rose to 72 per cent. and the urea fell to 73 mg. per 100 c.c.m. but four hours later the urea had risen to 150 mg. per 100 c.c.m. At this stage (February 4th) laparotomy was performed under local anaesthesia by Mr. G. R. A. Syme, who found a chronic ulcer on the anterior wall of the duodenum. After excising

the ulcer he did a posterior gastro-enterostomy. The following day the blood urea had fallen to 93 mg. per 100 c.cm.; two days later it was 103 mg.; a week later 41 mg. per 100 c.cm., where it remained for two months. She left hospital five weeks after the initial haemorrhage, having fully recovered. When seen two months later she was very well. The haemoglobin was 85 per cent. and the blood urea 32 mg. per 100 c.cm.

CASE V

A male aged 53 was admitted to the Royal Melbourne Hospital on December 29th, 1935, under the care of Dr. W. W. S. Johnston. For years he had suffered from epigastric discomfort after meals and often vomited. Two hours before admission he vomited two pints of bright red blood. His motions had not been dark in colour.

Examination showed a pale man lying comfortably in bed and complaining of thirst. His mental condition was perfect; lungs clear; blood pressure, systolic 68 mm., diastolic 50 mm.; pulse rate 84; respirations 20 per minute; haemoglobin 70 per cent. He was given ice to suck, water per rectum, and morphine in liberal quantities. On the following day he appeared to be improved, and citrated milk was given. The

systolic blood pressure rose slowly to 160 mm. Four days after admission he became paler and vomited more than half a litre of blood. The haemoglobin fell to 55 per cent. and the systolic blood pressure to 110 mm.; the blood urea was 76 mg. per 100 c.cm. A continuous transfusion was begun, and 1,800 c.cm. (three pints) of citrated blood were given in twelve hours. The haemoglobin rose to 68 per cent., and he showed great improvement. The urea fell slowly to 40 mg. per 100 c.cm. Twelve hours after the termination of the transfusion a laparotomy was performed under local anaesthesia and a chronic prepyloric ulcer was excised by means of a V-shaped incision. He appeared to be making a perfect recovery when eleven days after operation the blood urea was found to be rising, an increase from 53 to 100 mg. per 100 c.cm. occurring in the next three days. It was then discovered that the haemoglobin had fallen to 30 per cent. He was pale, restless, and semi-comatose. Later he vomited a litre of blood. A drip transfusion was begun, and 3,300 c.cm. (5½ pints) of blood were given in forty-eight hours. The haemoglobin rose to 51 per cent. and the blood urea fell to 51 mg. per 100 c.cm. His colour was greatly improved, and he was less confused mentally. On the following day he vomited over a litre of blood and sank into a typhoidal state—he was drowsy, restless, and delirious. Although the haemoglobin was maintained at a moderately high level by a further transfusion of 1,200 c.cm. the urea rose to 129 mg. per 100 c.cm. After existing in a moribund state for four days he died.

A necropsy was carried out by R. J. Wright-Smith, who reported that in the stomach there was wide acute ulceration around the suture line left after excision of the chronic ulcer. Three eroded arteries were seen in the floor of the ulcer. The pylorus was hypertrophied and the pyloric orifice was narrowed and admitted a finger with difficulty. The kidneys were of normal size; the capsules were slightly adherent, tearing the surface a little on stripping. The right kidney showed an old depressed scar at its upper pole; on section, the cortex and medulla were wide; the cortex was slightly swollen, the substance was pale red. The outlines of the pyramids were prominent, and the vessels congested in them. The large vessels were thickened and firm. The

pelvic fat and mucosa appeared normal. The ureters were small, thin-walled, and clear. Microscopically the glomeruli showed some patchy swelling with cellular increase, and here and there some vacuolation, but no gross disturbance was apparent. There was some tubular spoiling and degeneration.

Subacute Haemorrhage

During the present study it was observed that a separate group of patients could be identified who were suffering from subacute haemorrhage. The rate, as well as the magnitude of the fall in the circulating haemoglobin appears to be of great importance in determining the degree both of prostration and of mental confusion, as well as the level of the blood urea. It has already been shown that a severe haemorrhage leading to a fall in haemoglobin from 80 to 30 per cent. in twenty-four hours produces a profound effect. On the other hand, a slow haemorrhage which lasts a week or more but eventually causes the same fall in the haemoglobin produces quite a different picture. The patient is deathly pale, but

apart from breathlessness on exertion and a sense of muscular weakness he experiences but little discomfort. A haematemesis finally causes the physician to be summoned to his bedside and his immediate removal to hospital is ordered. There it will be found that the patient's mental condition is perfect and he merely protests that he is feeling quite well and

FIG. 5.—Case V. Effect of repeated haematemeses. Two continuous drip transfusions of citrated blood (total of 6,300 c.cm.) were given with 1,800 c.cm. of glucose-saline.

hopes to be back at work in a day or so. He may state that he has become progressively paler for the past few weeks. His tongue is moist and slightly coated and his thirst is easily appeased. The blood pressure is within normal limits and the pulse rate varies from 80 to 110 a minute. The blood urea is usually slightly elevated or may even be normal, but fluctuations are recorded from day to day, usually between the levels of 35 and 60 mg. per 100 c.cm.; readings of over 75 mg. are rare. If the bleeding has been present for many weeks there will probably be depletion of the iron stores and a typical microcytic hypochromic anaemia will be present.

It is probable that only a small vessel has been eroded and the danger of a large haemorrhage is remote. In the majority of these patients the blood picture is restored to normal after many weeks of careful dieting, alkalis, and adequate iron medication. However, to allow these patients to remain for many days in a state of extreme exsanguination involves great risks. The tissues are inadequately supplied with oxygen, the resistance to infection is diminished, the kidneys show signs of stress as indicated by the moderate rise in blood urea, and, finally, if a sudden haemorrhage of two pints or more does occur, it will probably prove rapidly fatal. It appears rational to elevate the haemoglobin by slow transfusion unless sufficient evidence is forthcoming that such a procedure definitely increases the liability to a further haemorrhage. In my limited experience continuous blood transfusion does not promote haemorrhage, and the haemoglobin is

rapidly elevated to normal levels. There is then a wide margin of safety should another haemorrhage occur, or should a surgical operation be necessary, this can be safely performed within a week of the transfusion.

CASE VI

A female aged 43 was admitted to the Royal Melbourne Hospital on February 1st, 1936, under the care of Dr. H. Turnbull. Nineteen years previously she had had a haematemesis, and since then she had suffered from intermittent attacks of indigestion, epigastric pain coming on two hours after taking food. She had had two subsequent haemorrhages, the last one being three years before, after which an operation was performed for excision of a chronic duodenal ulcer and a gastro-duodenostomy. Since then she experienced pain under the left costal margin with no definite relationship to food, but relieved by alkalis. During the three months prior to admission she became noticeably paler, and this had frequently been discussed by her friends. Muscular weakness, breathlessness, and a fear of fainting soon prevented her from carrying out her household duties. Her motions were often black in colour. Twenty-four hours before admission she vomited about half a litre of coffee grounds, and although she was greatly alarmed she experienced but little discomfort. A physician was summoned, and he advised her removal to hospital.

Examination showed an extremely pale woman of middle age. She was mentally alert, and gave an accurate account of her former illnesses. Her tongue was clean and moist, and her breath was not offensive. The lungs were clear. The systolic blood pressure was 105 mm., diastolic 80 mm.; pulse rate 90; haemoglobin 35 per cent.; blood urea 42 mg. per 100 c.cm. Unfortunately the red cells were not counted at this stage. A serious view of the case was taken, and a continuous drip transfusion of blood was immediately begun; 2,400 c.cm. (four pints) of citrated blood, followed by 1,200

c.cm. (two pints) of glucose-saline solution were given in thirty-six hours. The haemoglobin rose to 72 per cent. The red cells were 3,680,000 and the colour index was 1.0. A week later the red cells had increased to over 5,370,000 and

FIG. 6.—Case VI. Small haematemesis following chronic gastric haemorrhage causing only a slight rise in blood urea. Continuous drip transfusion of 2,400 c.cm. citrated blood and 1,200 c.cm. glucose-saline produced a rapid rise in haemoglobin.

the haemoglobin to 85 per cent. Iron was given (60 grains of iron and ammonium citrate daily) for three days, and was then withheld to see if the colour index would drop. One month later the red cells were 5,180,000 and the haemoglobin 71 per cent., giving a colour index of 0.7, and at the end of the following month the red cell count had fallen to 4,750,000 per c.mm. and the haemoglobin to 63 per cent., giving a colour index of 0.6.

CASE NO.	SEX	AGE	DIAGNOSIS	TREATMENT	RESULT	DAYS AFTER INITIAL HAEMORRHAGE											
						1	2	3	4	5	7	10	14	21			
7	M	82	CHRONIC GASTRIC ULCER	GLUCOSE-SALINE 2,400 C.C.M.	Death	Haemoglobin 30	30										
						Urea 150											
8	M	37	ACUTE GASTRIC ULCER	GLUCOSE-SALINE 3,000 C.C.M.	Relief	Haemoglobin 70	68			70			79				
						Urea 33	25			30							
9	M	64	ACUTE GASTRIC ULCER	GLUCOSE-SALINE 4,800 C.C.M.	Relief	Haemoglobin 62	46	48	44	43	48	40	42	42	48		
						Urea 57	63	34		31							
10	M	40	CHRONIC DUODENAL ULCER	GLUCOSE-SALINE 3,000 C.C.M. BLOOD 1,800 C.C.M.	Relief	Haemoglobin 50	43		47								
						Urea 38	67	67	35								
11	M	50	ACUTE GASTRIC ULCER	GLUCOSE-SALINE 5,000 C.C.M. BLOOD 1,800 C.C.M.	Death	Haemoglobin 69	47	50	66	51	52	35					
						Urea 72	66	59	45	59		110					
12	M	42	CHRONIC GASTRIC ULCER	GLUCOSE-SALINE 3,600 C.C.M. BLOOD 2,100 C.C.M.	Death	Haemoglobin 70	70		38	35							
						Urea 94	82		82	193	112						
13	M	55	GASTRIC CARCINOMA SECONDARY GROWTHS	BLOOD 2,100 C.C.M. GLUCOSE-SALINE 5,600 C.C.M. LAPAROTOMY	Relief	Haemoglobin 38		55	52			60	60	60	55		
						Urea 61	57	49	67	51		50					
14	M	52	ACUTE DUODENAL ULCER	BLOOD 2,100 C.C.M. GLUCOSE-SALINE 600 C.C.M.	Relief	Haemoglobin 59	35	51	77	69	64	59	67	67	74	45	
						Urea 119	137	132	125	87		54	67	50			
15	F	53	ACUTE GASTRIC ULCER	BLOOD 2,000 C.C.M. LAPAROTOMY (NIL FOUND)	Relief	Haemoglobin 55	62	70	72	68		70	75	80			
						Urea 127	160	87	82	39	41	31					
16	M	50	CHRONIC DUODENAL ULCER	BLOOD 2,100 C.C.M. GLUCOSE-SALINE 800 C.C.M. ULCER INFOLDED	Relief	Haemoglobin 68	50	58		65		42			66		
						Urea 72	53	45		41		47	41	39			
17	M	42	CHRONIC DUODENAL ULCER	IRON (PALLOR ONE MONTH BEFORE HAEMATEMESIS)	Relief	Haemoglobin 36	37	34	32			33	33	32	38	41	
						Urea 46	48	37	33	45		52	50		45		

Table showing the effect of the transfusion of glucose-saline (shaded area) and citrated blood (black area) on the haemoglobin percentage and blood urea in eleven additional cases of haematemesis. The blood urea is expressed in mg. per 100 c.cm. "H" denotes haemorrhage.

Discussion

Experience with the various methods of transfusion in cases of severe haemorrhage has shown that massive transfusion of blood by the continuous drip method gives by far the best results. There is not as yet sufficient evidence to indicate the place of surgery in the treatment of haematemesis from peptic ulcer, but should surgical intervention be decided upon it is essential that the patient be adequately prepared for the operation. The surgeon should be able to carry out a gastrectomy or gastro-enterostomy without endangering the patient's life. This point has recently been emphasized by Gordon-Taylor,⁹ who insists that operation should not be attempted until the haemoglobin has been restored to 80 per cent.

A study of the changes in the level of the blood urea has proved of value in assessing the extent and rapidity of the haemorrhage. Results also suggest that it is desirable to postpone operation for twelve to twenty-four hours after the haemoglobin has been restored by the transfusion. Whether a high blood urea is a contra-indication to operation is a question to be answered by further investigation, but it is a reasonable assumption that if the blood urea is relatively low and falling the results will be better than when it is high and rising. When viewed in conjunction with the general appearance of the patient and the height of the haemoglobin, the blood urea has proved in some cases to be a useful guide both to treatment and to prognosis.

Summary

1. The effects and dangers of severe haemorrhage are discussed with special reference to the haemoglobin percentage, blood pressure, and pulse rate, and to the reduction and replacement of blood volume.

2. The effects of anoxaemia on the tissues are described, and a study is made of the disorders of kidney function and elevation in blood urea.

3. The advantages of massive blood transfusion by the continuous drip method of Marriott and Kekwick in the treatment of severe haematemesis are compared and contrasted with glucose-saline transfusion. A series of cases are described illustrating changes in mental state, haemoglobin percentage, and blood urea. The production of oedema and the effects of repeated haemorrhages are also discussed.

4. The effects of chronic haemorrhage are described with special reference to the relatively small elevation in blood urea.

5. It is suggested that the changes in height of the blood urea indicate the extent and rapidity of gastrointestinal haemorrhage, and aid in prognosis and treatment.

Technique.—The haemoglobin estimations were carried out by Sahli's method, 14 grams of haemoglobin representing 100 per cent. McLean's modification of van Slyke's method was used for the blood urea estimations.

Acknowledgement.—I wish to express my sincere thanks to Dr. Charles Kellaway, director of the Walter and Eliza Hall Institute, and Dr. Hume Turnbull for their help and encouragement during this investigation. I am deeply indebted to Miss Beryl Splatt, biochemist of the Royal Melbourne Hospital, for carrying out all the biochemical tests, and to the members of the honorary medical and surgical staffs of the hospital for placing their records at my disposal. The untiring service of the resident medical staff has made this work possible.

REFERENCES

- ¹ Marriott, H. L., and Kekwick, A.: *Lancet*, 1935, i, 977.
- ² Idem: *Proc. Roy. Soc. Med.*, 1936, xxix, 337.
- ³ Robertson, O. H., and Bock, A. V.: Medical Research Committee, Special Report Series, No. 25, p. 213.
- ⁴ Holman, R. L., Mahoney, E., and Whipple, G. H.: *Journ. Exper. Med.*, 1934, lxx, 269.
- ⁵ Christianson, T.: *Acta Med. Scand.*, 1935, lxxxv, 333.
- ⁶ Ingegnio, A. P.: *Journ. Amer. Med. Sci.*, 1935, cxc, 770.
- ⁷ Van Slyke, D. D., Rhodes, C. P., Hiller, A., Alving, A. S.: *Amer. Journ. Physiol.*, 1934, cix, 336.
- ⁸ Bayliss, W. M.: Medical Research Committee, Special Report Series, No. 25, p. 11.
- ⁹ Gordon-Taylor, G.: *Lancet*, 1935, ii, 811.

GONOCOCCAL ARTHRITIS IN THE MOTHER AND NEWBORN INFANT

BY

JEAN M. MACLENNAN, M.B., CH.B.

ASSISTANT MEDICAL OFFICER, SWANSEA CORPORATION

Gonococcal arthritis in pregnancy and the puerperium is a commonly recognized condition. Gonococcal arthritis in the newborn infant is less common. The coincidence of the two appears to be rare.

With regard to the incidence of arthritis complicating gonorrhoeal ophthalmia neonatorum Altland¹ (1902), who collected nineteen cases, including only one of his own, referred to the "extraordinarily rare occurrence" of the condition. Brehmer,² in 1905, was able to find accounts of twenty-five cases. Still³ mentions "various causes of swelling about or in the joints in infancy" (including gonorrhoeal arthritis), "not because they are common, for they are all very rare," but Holt and McIntosh⁴ state in connexion with gonococcal ophthalmia that "many cases of acute polyarthritis have been observed." Thomson⁵ says that the condition "has been described," and gives a number of references, while several paediatricians define it as "occasional" or "rare." In spite of its comparative rarity this form of arthritis has been very completely investigated, particularly by Continental workers, gonococci having been demonstrated in the affected joints by Deutschmann,⁶ Höck,⁷ Altland,¹ Brehmer,² and others.

The swelling of the joint or joints may occur from the fifth day to the fifth week after infection, but it appears usually about the second week. The joints most frequently involved are the knee, wrist, and ankle; more rarely the hip, elbow, and metacarpo-phalangeal joints are concerned. In some records it has been noted that the ophthalmia has been of a particularly virulent nature, several of the cases having terminated fatally, but in others the rapid improvement under treatment has been striking, and the prognosis regarding the joint is generally good.

The simultaneous occurrence of gonococcal arthritis in mother and newborn infant has been described by Brehmer² and Schiller⁸ (four cases). A hasty search has not unearthed details of any similar cases in the English literature, but no doubt some have been observed. The following case presents some interesting features.

Case Record

A primigravida, aged 27, was referred on May 29th, 1935, at the eighth month of pregnancy, to the ante-natal clinic by a midwife on account of breech presentation; she had then no complaints. External version was performed in hospital on the following day. She was seen again at the clinic on June 19th, complaining of pains in the left knee, both ankles, both elbows, and right wrist; the right wrist was swollen and stiff. She was sent home to bed, and aspirin and antiphlogistine were recommended. On the morning of June 21st the midwife reported that the membranes had ruptured but the foetal head was high; the wrist was still swollen. The patient was seen by her own doctor, who sent her into hospital at midday on June 22nd on account of delay in labour.

On admission the patient's temperature was 99.8° F. and the pulse rate 124; she was having slight pains. The foetus was presenting by the vertex in the L.O.A. position. There was a profuse vaginal discharge. At 8.15 a.m. on June 23rd the cervix was fully dilated. The liquor was noted as being "thick and green." At 11 a.m. the patient delivered herself of a baby girl, 5 lb. 5 oz. in weight. It was remarked at the time that the infant's eyes were open and "staring," and fouled with the thick, dirty liquor. The eyes were swabbed immediately, and 10 per cent. argyrol drops were instilled. After delivery the mother's temperature dropped to 97.6° and the pulse to 94. The mother made a good recovery, the