level. It was suggested that if the induction pipes on ground level were made to act as extraction pipes there would then be a uniformly downward system of ventilation. This suggestion was adopted, and when put in operation produced a most dramatic drop in the number of cases of keratitis, as is strikingly shown in the graph. Moreover, from this time keratitis became a rarity. A few outbreaks have taken place from time to time, but, with one very important exception, they were traced to the closing down for one reason or another of one or more of the extraction fans. This seems to be conclusive proof that the cause is gaseous, or at any rate an emanation from some volatile substance.

Thio-formaldehyde

The exception just mentioned was in December, 1935. There was an outbreak of keratitis which could not be attributed to any fault in the extraction system in the spinning room. On inquiry it was found that, owing to a coal shortage, the acid house evaporating temperature was very much lowered and the fans over the tanks were stopped. This at once suggested that normally the evaporation in the acid house served also to drive off excess of the causal substance which was produced in the spinning room. During the coal shortage this did not

take place, and so the acid returned to the spinning room already charged with the substance to such an extent as to render even the new ventilation insufficient. On asking whether there might be some volatile substance which failed to be driven off at this time, the reply was given that such a substance was present—namely, thio-formaldehyde, which is formed from CS_2 in the presence of nascent hydrogen. It is thought that this reaction takes place readily in the spinning bath. Thio-formaldehyde is a little-known substance, having a curious odour suggestive of stale cabbage. Might not this, plus H_2S , give rise to the "oniony" breath referred to early in this paper?

Experiments are being undertaken to ascertain whether this substance has any relation to keratitis, and, if so, in what way it acts. Should interesting results be obtained they will be published in due course.

In conclusion, I wish to thank those who have encouraged and helped me in the writing of this paper, and particularly Sir Stewart Duke-Elder, who kindly read the notes and advised publication, and Dr. Berkson, who translated the foreign writings.

REFERENCES

- ¹ *Schweiz. med. Woch.*, 1923, cxiii, 560.
- ² *Arch. d'Ophthalm.*, 1928.
- ³ *Ibid.*, 1929.
- ⁴ *Text Book of Ophthalmology*, vol. i, pp. 818, 817.

INFECTIVE HEPATITIS WITH A FATAL RECRUDESCENCE

BY

T. E. LOWE, M.D. MELB., M.R.C.P. LOND.
MEDICAL REGISTRAR, ST. MARY'S HOSPITAL, LONDON

(With Special Plate)

Clinically the syndromes of catarrhal jaundice and of toxic hepatitis merge into one another. Pathologically the underlying changes in catarrhal jaundice have not been clearly defined. Recent reviewers of the subject suggest that the clinical syndrome of catarrhal jaundice represents at least three different pathological states. An obstructive jaundice due to blockage of the common bile duct by duodenal catarrh, a scattered liver necrosis, and, thirdly, an obstructive jaundice from cholangitis have all been suggested by pathological examination.

In view of the lack of unanimity as to the pathological state of the liver in the clinical syndrome of catarrhal jaundice the following case is worthy of record. In brief, it is the history of a disease in a young girl, which began apparently as a catarrhal jaundice, and passed on to a condition of portal obstruction. After a period of 300 days she had recovered clinically save for the evidence of portal obstruction. Two and one-third years later a similar illness commenced and continued with jaundice and intestinal upsets for forty-seven days, at which time a laparotomy was performed. Subsequent to operation she died from septicaemia. The post-mortem examination revealed old cirrhosis of liver and also recent hepatitis.

Clinical Record

A young girl, a jam packer by trade, was first seen at the age of 12 as a medical out-patient on March 22nd, 1932, when she gave the following account of her illness. Two weeks previously she had begun to get attacks of epigastric pain a short time after meals. These attacks lasted for seven days, when there developed a slight yellowness of her skin. The jaundice rapidly became worse. Associated with its onset there was one attack of vomiting and some nausea. She was admitted to the medical ward on the same day (fifteenth day of illness) and was described as a cheerful, well-nourished, very jaundiced girl. Her temperature was

99.4° F., and the pulse rate was 100. There were no abnormal physical signs in the heart, lungs, or nervous system. The liver edge was felt $1\frac{1}{2}$ inches below the right costal margin in the mid-clavicular line. The urine was acid in reaction, with a specific gravity of 1030; it contained no albumin or sugar, but both bile salts and pigments were present in considerable quantities. She was treated with a fat-free diet and with small doses of calomel and magnesium sulphate by mouth. The patient remained in hospital until April 12th, during which time the following notes were made.

March 29th (22nd day).—A bacteriological examination of the faeces showed the presence of non-haemolytic and *viridans* streptococci in moderate numbers. No non-lactose fermenting organisms were isolated.

April 2nd (26th day).—A fat analysis of the stools showed: neutral fats, 0.4 per cent.; fatty acids, 1.8 per cent.; and soaps, 8.3 per cent. The jaundice was now much less intense, and the stools were dark in colour.

November 16th (254th day).—She was again seen in the out-patient department, when she was slightly jaundiced. The liver edge was still easily palpable and the spleen easily felt for the first time. From November, 1932, until April, 1935, she was in good health, and stated that her jaundice had completely cleared by the end of 1932 (300th day).

April 21st, 1935.—The jaundice returned with malaise but no abdominal pain. There was a little diarrhoea, the stools were pale, and the urine was very dark in colour. She was readmitted to the wards on April 29th with a temperature of 99.4°, a pulse rate of 120, and a respiratory rate of 24. She was markedly jaundiced, but well nourished, and she had no itching of the skin. There were no abnormal physical signs in the heart, lungs, or nervous system, but there was slight furring of the tongue. The spleen now reached the umbilicus, and the liver edge was felt two inches below the costal margin in the mid-clavicular line. There was some fullness in the left loin. The urine was alkaline in reaction, the specific gravity being 1015; it contained no sugar, a trace of albumin, and some bile salts, but no bile pigments. She was again treated with a fat-free diet, glucose drinks, and calomel and magnesium sulphate by mouth. The progress of her second illness is shown by the following notes.

May 4th (14th day).—The blood examination showed: haemoglobin, 85 per cent.; erythrocyte diameter, 7.4μ ; leucocytes, 6,000—polymorph cells, 70 per cent.; lymphocytes, 22 per cent.; large mononuclears, 7 per cent.; basophil cells, 1 per cent.; and reticulocytes, less than 0.5 per cent.

May 8th (18th day).—Fat analysis of the faeces showed: neutral fats, 4.2 per cent.; fatty acids, 2.4 per cent.; soaps, 5 per cent. The red cell fragility was within normal

limits. (Complete haemolysis at 0.365 per cent.) A bacteriological examination of the faeces resulted in a heavy growth of *Streptococcus viridans*.

May 14th (24th day).—Her jaundice was appreciably deeper, and fluctuating from day to day. The spleen edge reached one inch below the umbilicus.

May 15th (25th day).—A blood examination showed: haemoglobin, 65 per cent.; erythrocytes, 4,600,000 cells per c.mm.; leucocytes, 10,000 cells per c.mm., consisting of polymorphs 77 per cent., lymphocytes 22 per cent., and large mononuclears 1 per cent.

May 23rd (33rd day).—The patient was becoming very wasted and felt depressed. After this the spleen became smaller, and the fullness in the loin disappeared.

May 30th (40th day).—She complained of severe upper abdominal pain, and vomited several times. Temperature 100° F., pulse rate 106. The leucocytes numbered 6,000, with 52 per cent. of lymphocytes. On the following day her temperature rose to 101°, the jaundice increased, there was slight rigidity in the left upper abdomen, and the left loin was again bulging. The abdominal pain disappeared in the ensuing five days, but she developed a swinging temperature up to a maximum of 101.5°.

June 6th (47th day).—After pre-operative medication with calcium chloride a laparotomy was performed. Free fluid, dark yellow in colour, was found in the abdominal cavity. The liver was firm, knobby, and generally shrunken; it was described as a liver typical of late cirrhosis. The spleen was enlarged, and enlarged abdominal lymph glands were felt. After operation the patient was comfortable and free from pain. She remained so until June 14th (fifty-fifth day), when she rapidly became worse, and died on June 16th (fifty-seventh day).

Pathological Notes

At the post-mortem examination the following significant changes were found. A recent operation wound in the epigastrium was partially broken down, and there was over a litre of dark, semi-purulent fluid in the abdominal cavity.

The liver (1,400 grams) was tough and flabby, greatly deformed, and cirrhotic. The original phrenic surface was contracted and deeply grooved, whilst the inferior surface was increased and very convex. Both the face and cut surfaces were studded with tawny-red regeneration nodules. On microscopical examination numerous bile ducts, veins, arteries, and lymphatics were seen in the scar tissue. The course taken by these was made tortuous by the liver nodules. Aggregations of lymphocytes and plasma cells were present, with some lipid-laden histiocytes and an occasional eosinophil cell. The scar tissue was not dense, and there was moderate elastic tissue formation. The nodules presented a distorted lobular pattern, most of them being drained at several points by bile ducts, which penetrated or lay alongside them. Localized about the "junction zones" there was a mild inflammatory process in which lymphocytes, eosinophil, neutrophil, polymorphonuclear, and some plasma cells took part. The epithelial cells in these areas showed severe degenerative changes. In some tubules (both initial bile ducts and adjacent liver cell columns) the epithelial lining was missing, and in a few of these there were "casts" formed by the agglutination of desquamated cells. In most of the tubules the lumen was narrowed by great swelling of the cells. Away from the junction zone the liver cells were hypertrophied, and showed some fatty changes. Numerous bile plugs were present, especially about the efferent veins. There were also scattered focal necroses, and at other points post-mortem colonies of micrococci.

The gall-bladder contained concentrated bile, and between it and the duodenum there were old adhesions which probably dated back to the original illness. There was marked portal obstruction, with very striking anastomoses between the veins of the portal system and those of the anterior and posterior abdominal walls, diaphragm, and oesophagus. Near their junction the circumferences of the superior mesenteric, splenic, and portal veins were respectively 2.95, 3.1, and 3.5 cm., compared with 2.25, 1.7, and 2.4 cm. of a normal control.

The gastro-intestinal mucosa was oedematous, plum coloured, and showed mucous catarrh.

The spleen (1,000 grams) with a spleniculus (3 cm. in diameter) was greatly enlarged. The cut surface was dark,

swollen, and mushy, the Malpighian bodies being small. Polymorphonuclear leucocytes were numerous in the congested pulp, and around the Malpighian bodies there was a zone of large pulp cells.

The abdominal lymph glands were also enlarged, and some were dark red owing to the presence of blood and inflammatory products in the sinuses.

Throughout all organs there were petechiae and focal necroses due to the terminal septicaemia.

Commentary

The correlation between the clinical history and the subsequent pathological findings in this case is very close. The initial illness in March, 1932, commenced as a typical attack of catarrhal jaundice, with symptoms suggesting a mild gastric inflammation, followed after a period of seven days by jaundice, liver enlargement, and slight pyrexia. Its subsequent course, however, did not follow that of jaundice due to gastro-duodenal catarrh, for the jaundice did not disappear until the 300th day; moreover, the enlargement of the spleen was not noticed until the 254th day, when the liver was still palpable. This agreed with the finding of old adhesions in the duodenal region representing the initial gastro-duodenal lesion; the marked cirrhosis also explains the late splenic enlargement.

After an interval of two years and four months an illness began which again clinically suggested a catarrhal jaundice, but in contrast with the previous illness was associated with an enlarged spleen. Until the fortieth day of this illness the outstanding features were jaundice of varying intensity and the splenic enlargement. On the fortieth day a fresh set of symptoms developed: abdominal pain, vomiting, muscular rigidity in the left upper abdomen, and pyrexia—suggesting a fresh gastric inflammation. After the laparotomy on the forty-seventh day the jaundice remained stationary, and the intestinal symptoms disappeared.

Pathologically this second illness was associated with inflammation in the "junction zone" and destruction of liver cells. The appearances suggest that the jaundice was due to a mechanical effect from pressure on the initial bile ducts, and to the hepatic cell destruction. Possibly desquamation and obstruction by debris played a part, but this was difficult to demonstrate. As one would expect from this picture the jaundice was variable. The post-mortem examination gives the clue to the patient's terminal condition. It was due to a peritonitis and an associated septicaemia.

Considering the pathogenesis of the condition, there appear to be two types of lesion which might give rise to the microscopical appearances present: either a toxic destruction of liver cells with secondary inflammatory changes about the initial bile passages, or an actual infection of the liver tissue. As regards toxic factors, we must think of extraneous poisons, such as the benzene derivatives, which might have been introduced during her occupation. This was considered at the time of her illness, but no evidence of their presence was found. Blood-borne bacterial toxins give a second group; in this case the gastro-intestinal canal would appear to be the likely place of origin. If the lesion is primarily a destruction of liver cells, the inflammatory reaction about the initial bile passages must be secondary, and due to the drainage of the products of cell destruction to the "junction zone" of the lobules along the liver lymph spaces.

There are two types of liver infections to be considered: (a) bacterial invasion, or (b) virus infection. Since there is no evidence of damage to the larger bile passages, it would appear that the infection must have been blood-borne from the gastro-intestinal tract, where it seems likely to have made its entry. A virus infection would cause cellular destruction of the actual liver cells, and

the drainage of the products of disintegration to the "junction zone" would give rise to the inflammatory lesion seen there. Since little is known about any virus infections of the liver, this is simply conjecture along the analogy of the lesions in the central nervous system produced by the virus of poliomyelitis, where there is first a cellular destruction followed by a secondary inflammatory reaction in the most affected areas.

An important question in this case is whether the two illnesses were of the same type, or were entirely unrelated except in so far as the liver was badly damaged by the first. It is also possible that the whole picture was merely one condition, with an apparent remission of two years in the middle of its course. It is tempting to link the two states, and consider them either as one infection or a reinfection with the same causal organism. It is possible that there was a persisting bowel infection throughout, and that owing to some lowering of the patient's resistance a fresh invasion of the liver occurred. If the condition was a virus infection it might have been lying latent in the liver cells during the remission, to become active again with alteration in the host's resistance. Beyond saying that the mode of entry of the causal agent appears to have been via the alimentary tract, the foregoing discussion is conjecture, and the condition could equally well fall into the group of unexplained liver cirrhoses.

Summary

A disease is described in a growing girl in which a prolonged attack of jaundice was preceded by symptoms of gastro-intestinal infection and followed by hepatic cirrhosis. After a latent period characterized by increasing portal stasis with splenomegaly the jaundice returned. Death resulted from post-operative peritonitis and septicaemia.

It is suggested that, except for the terminal events, the whole course was due to a persisting bacterial or virus infection.

I wish to thank Dr. Reginald Miller, under whose care the patient was admitted, for permission to report this case. I also wish to express my indebtedness to the pathological institute of St. Mary's Hospital for the pathological notes and photographs, especially Dr. D. M. Pryce for helpful suggestions concerning the pathogenesis of the condition.

In a recent publication (*Public Health Reports*, 1936, li, 363) emanating from the School of Hygiene and Public Health of Johns Hopkins University, Dr. A. W. Hedrich discusses the changes in the incidence and fatality of small-pox in recent decades. The highest rates appear to be found in India, China, and Mexico. Intermediate rates ranging from 5 to 50 per 100,000 occurred during 1921-30 mainly in England, the United States, and Canada, but since 1920 the incidence in England has fallen to a very low level. The lowest rates are also found mainly in Continental Europe, Australia, and Japan. International reports indicate that well-vaccinated countries tend to have low attacked rates, and vice versa. Low case fatality has been prevalent in recent decades in Anglo-Saxon countries, South Africa, and Spain, while small-pox with a high fatality is the predominating form in India, Central Europe, Japan, and Mexico. In the United States the case rates are highest in the north-west and lowest in the north-east regions, the rates in the north-west during 1921-30 averaging well over 100 per 100,000 as compared with a mean rate of 2.6 for the New England and Middle Atlantic States. As regards case fatality an important part is played by vagrants, migratory labourers, and the like in importing malignant small-pox from Mexico and Asia, where the type of disease is endemic. In conclusion Hedrich maintains that small-pox deaths or case fatality rates at present probably represent the best available index of malignant small-pox.

TREATMENT OF RECENT DISLOCATION OF THE CARPAL SEMILUNAR BONE*

BY

A. F. GOODE, M.B., CH. B., F.R.C.S.

ASSISTANT IN THE SURGICAL UNIT, WELSH NATIONAL SCHOOL OF MEDICINE, THE ROYAL INFIRMARY, CARDIFF

(With Special Plate)

This article has been written for two reasons. First, dislocation of the carpal semilunar bone is not an uncommon condition, and during the last twelve months of general surgical practice I have seen six cases. Secondly, it would appear that insufficient recognition has been given to L. Böhler's methods of reduction.¹

In order to appreciate the methods of reduction the mechanism of dislocation must be described. Rather than a dislocation of the semilunar the lesion at first is really a dorsal dislocation of the hand on the semilunar bone and radius. The dislocation is produced by a hyperextension injury of the wrist—usually by a fall or a blow on the outstretched hand. In an x-ray photograph of a normal wrist from the side it will be seen that the radius, semilunar, and os magnum are in one straight line (Fig. i). For descriptive purposes dislocation of the semilunar may be divided into three degrees:

1. *First Degree*.—Neither anterior nor posterior ligaments attaching the semilunar to the radius are torn; the hand dislocates backwards, and there is little or no upward displacement of the hand (Fig. ii).

2. *Second Degree* (Fig. iii).—Here the hand has been displaced upwards as well as backwards, and as it tends

to spring back into its normal position it pushes the semilunar from behind, the relatively weak posterior ligament gives way, and a certain amount of forward rotation occurs.

3. *Third Degree*.—Here the semilunar has been rotated 90 degrees—it may be more—and the os magnum is occupying its place, articulating with the distal end of the radius.

If the scaphoid is also fractured the inner fragment usually follows the semilunar.

Diagnosis and Method of Reduction

A dislocation may be inferred by the presence of a swelling over the anterior part of the wrist and the limitation of movements of the wrist, particularly flexion. In addition, there is frequently tingling along the distribution of the median nerve in the hand. Proof of dislocation can only be given by x-ray examination, a lateral one being of greatest value.

I have always used general anaesthesia, but local anaesthesia undoubtedly would be better in early cases. First degree dislocation is very easy to reduce. All that is necessary is hyperextension of the wrist and the application of pressure over the anterior swelling. The dislocation can be felt to snap back into its correct position.

* From the Surgical Unit, the Welsh National School of Medicine, the Royal Infirmary, Cardiff.

T. E. LOWE: INFECTIVE HEPATITIS WITH A FATAL RECRUDESCENCE

FIG. 1.—Low-power view, showing extensive subcapsular destruction of liver tissue, lobulation of the liver, and gross fibrosis. (van Gieson stain.)

FIG. 2.—High-power view of the junction zone, showing recent damage to liver cells, and cellular infiltration of the fibrous tissue and desquamation of bile duct epithelium. (H. and E. stain.)

A. F. GOODE: TREATMENT OF RECENT DISLOCATION OF THE CARPAL SEMILUNAR BONE

FIG. 1.—Skiagram of early third degree dislocation.

FIG. 2.—Same after ten minutes' traction. Note the wide gap between os magnum and radius.

FIG. 3.—Dislocation reduced after open operation.