PARTIAL THORACIC STOMACH

WITH REPORT OF A CASE

BY

G. RONALD ELLIS, M.A., M.B.CANTAB., M.R.C.P.

HONORARY ASSISTANT PHYSICIAN, BIRKENHEAD GENERAL HOSPITAL
AND ROYAL INFIRMARY, WIGAN; SENIOR MEDICAL TUTOR,
DAVID LEWIS NORTHERN HOSPITAL, LIVERPOOL

(With Special Plate)

The condition of partial thoracic stomach is by no means a rarity, but comparatively few cases have as yet been recorded in the literature. The case to be described also presents several features which have not always been stressed in the recorded examples of the abnormality.

Case Report

A woman aged 50, in domestic service, was referred to the Birkenhead General Hospital by Dr. Dow. She had been complaining of lassitude and bodily weakness for the past six months. In addition, she complained of slight swelling of the ankles and pains in the head and legs. She stated that she had always been rather short-winded. There had been no previous illnesses of any importance.

On examination she was a well-built woman, but rather pale. There was slight pretibial oedema, and a cardiovascular system clinically normal; the blood pressure was 130/90. The chest was markedly "pigeon-breasted" and suggested emphysema, though the expansion was one and a half inches. The urine showed a moderate degree of albumin. The outpatient diagnosis seemed to lie between chronic renal disease and a hypochromic anaemia; the latter was supported by a history, gained by direct questioning, of brittle nails, falling hair, and a flatulent dyspepsia.

The patient was further questioned about her dyspnoea, and stated that she had never, even as a child, been able to run about as well as her sister. She dismissed this shortness of breath as unimportant, however, with the somewhat surprising statement that many years ago she had been told she "had only one lung." There had at no time been an x-ray examination of the chest. Apart from the "pigeon-breast" deformity there were no physical signs in the chest. A chest film, which was now taken (Fig. 1 on Plate), revealed a diaphragmatic hernia—as it was at first considered to be.

Clinical Investigations

The patient was admitted to the ward for fuller investigation. By means of a barium meal it was shown (Fig. 2 on Plate) that two-thirds of the stomach was above the diaphragm, posterior to the heart. No other contents of the abdomen were detected in the thorax. As well as could be ascertained by screening, the oesophagus appeared to enter at the summit of the thoracic pear-shaped sac, conforming to the type associated with a congenitally short oesophagus rather than to a true herniation of the diaphragm. The thoracic loculus emptied readily into the abdominal portion of the stomach, especially in the upright posture. I endeavoured to take a fractional test meal from each sac, but it proved impossible to extract any secretion from the upper loculus. The secretory curve obtained with the tube passed to the normal distance revealed that there was some delay in the disappearance of starch and the entry of bile.

The results of other investigations were as follows:—urine: albumin and pus cells; blood non-protein nitrogen, 37 mg. per cent.; urea clearance, 56 per cent. of normal. The Wassermann reaction was negative, and the blood count was: red blood cells, 4,710,000 per c.mm.; white blood cells, 14,680 per c.mm.; haemoglobin, 60 per cent.; colour index, 0.78. While in the ward the patient steadily lost weight. The tests revealed a definite degree of renal inadequacy, and this was held responsible for the anaemia. The latter, however, failed completely to respond to iron in massive doses, no impression being made on the blood picture.

Comment

Most of the recorded cases of true thoracic stomach, either partial or complete, have been in children. Adults have sought advice for severe dyspeptic symptoms, and

have usually been very ill patients. It is seldom that the condition has been discovered accidentally. The interest of the present example lies in that it was a chest symptom—namely, dyspnoea—which led to the discovery of the condition. The vague dyspepsia complained of by the patient, only admitted on direct questioning, was never such as to warrant an x-ray examination.

In a recent review of diaphragmatic herniae Sir Thomas Dunhill¹ mentions fourteen cases of thoracic stomach, and states that it is astonishing how frequently the condition causes symptoms. The cause of the condition is a failure of the oesophagus adequately to elongate, the stomach being partially arrested in the posterior mediastinum. The stomach comes to be hour-glass shaped, the cardiac loculus being within the thorax and the pyloric within the abdomen. According to Sir Thomas Dunhill dysphagia is common and usually the most disabling symptom, and dyspnoea may occur through pressure on the pericardium. He also states that the symptoms of the condition are such as strongly to suggest their origin. This is true of most of the recorded cases, and is the reason for their recognition, but it would appear highly probable that there are quite a number of individuals who carry the condition unrecognized to their graves, either through the total absence of symptoms or the failure to associate existing symptoms with the possibility of the condition.

The importance of dyspnoea as a symptom is stressed by Goodall and Hoyt,² who record five cases of thoracic stomach, in each of which the symptom calling for treatment was dyspnoea. The writers, having made the diagnosis in one case, were able to recollect two examples of dyspnoea where the cause had never been explained; on subsequent recalling these patients they were able to demonstrate thoracic stomachs. They divide symptoms into two groups, due to distinct mechanisms: (1) interference with normal peristalsis of the gastrointestinal tract; (2) embarrassment of the thoracic organs.

It would seem that the larger the thoracic loculus the more extreme the dyspnoea. The latter is said to be more marked after a meal, but in the present case the patient would not admit to this. The explanation of this breathlessness is not obvious. It would appear greater than can be explained by a simple reduction of pulmonary ventilating surface due to the bulk of the thoracic loculus in the chest. There seems little evidence to support the suggestion that interference with the mechanism of the diaphragm plays any part. The barrel shape of the chest in the case here recorded suggested emphysema as a possible explanation, but a chest expansion of one and a half inches is against this. It is difficult to disregard this striking chest deformity, and take the view that it is a coincidence. Might not the extreme prominence of the sternum be explained as compensatory to the greater number of structures which have to be accommodated between it and the vertebral column? Cardiac embarrassment due to pressure by the sac is not a convincing explanation of the dyspnoea; there was no cardiac hypertrophy detected in the present case such as would be expected in long-continued cardiac embarrassment. The precise cause of the dyspnoea has still to be explained.

Goodall and Hoyt state that dysphagia, often a distressing symptom in true diaphragmatic hernia, is not a symptom of thoracic stomach; the patient in the present case did not show dysphagia. Anaemia has been described, associated usually with peptic ulceration, but it would be a mistake to draw conclusions from the anaemia in this case.

REFERENCES

- ¹ Dunhill, Sir Thomas: *Brit. Journ. Surg.*, 1935, 478.
- ² Goodall, W. H., and Hoyt, L. H.: *Arch. Int. Med.*, 1934, 595.

G. RONALD ELLIS: PARTIAL THORACIC STOMACH

FIG. 1.—A direct x-ray of the chest, revealing the thoracic sac.

FIG. 2.—Demonstrates thoracic and abdominal sacs, after a barium meal. The upper sac is partly, the lower completely, filled with barium.

W. McN. NIBLOCK: PSEUDO-PANCREATIC CYST OCCURRING AFTER CHOLECYSTECTOMY

FIG. 1.—Skiagram taken before operation.

FIG. 2.—After operation. The arrow indicates lipiodol lying in the sinus.