TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some of the renal diseases met with in general practice. The first group of articles on Treatment in General Practice, thirty-five in number, was republished in book form last March by H. K. Lewis & Co., and a second volume will appear in the autumn.

THE TREATMENT OF NEPHROSIS

BY

ROBERT PLATT, M.D., F.R.C.P.

The type of renal disease generally known as nephrosis, or lipoid nephrosis, is characterized by its insidious onset and chronic course, and by the presence of copious albuminuria and generalized oedema, which is often extreme. Some hold that it is of a different nature from other types of renal disease; others that it is merely a variety of subacute nephritis, which it closely resembles. The controversy cannot be pursued here, but whatever view is held the differentiation of nephrosis, whether as a distinct entity or a variety of renal disease, is of some practical value, because in this type of case recovery may be complete (though it rarely is) even after months or years of illness, and in the fatal cases death is far more likely to supervene from intercurrent infection than from renal failure (uraemia). It is, in fact, typical of nephrosis that throughout its chronic course no failure of the excretory power of the kidney usually develops. The treatment of nephrosis, therefore, differs in some important respects from that of nephritis, and it is essential at the onset that a correct clinical diagnosis should be made.

This is done by a general survey of the history and the signs and symptoms. The onset, as already stated, is insidious. If there has been a previous attack of acute nephritis, if there is or has been any haematuria, or if the blood pressure is raised, the patient should on no account be treated as a case of nephrosis, but as one of active or unhealed nephritis. Should haematuria or hypertension develop during the further course of the disease the diagnosis should be revised, and the case treated as one of nephritis. Blood tests are of little value in the differentiation of nephrosis, because the characteristic high cholesterol and diminished protein content are also found in subacute nephritis. As there is no renal insufficiency the blood urea is not increased.

Two other conditions may closely resemble lipoid nephrosis and must further be distinguished because of their special treatment. (1) Syphilis may give rise to a similar clinical picture; if it is suspected and a positive Wassermann reaction is discovered the usual anti-syphilitic remedies should be employed. (2) Amyloid nephrosis is distinguished by the coexistence of tuberculosis or chronic suppurative disease and by the other signs of amyloidosis—namely, enlargement of the liver and spleen, and diarrhoea. The prompt treatment of the underlying infection represents the only hope of recovery.

Lipoid nephrosis occurs in infants, children, and young adults.

General Management

The diagnosis having been made, the first duty of the practitioner is to see that competent nursing can be obtained for the patient and continued if necessary for a very long time. Records will have to be kept of the quantity of urine passed, of the amount of albumin therein, and of the pulse and temperature. Diets must be accurately prepared, and the response to changes in treatment carefully noted. It is a great advantage if the patient can be weighed regularly, this being the most reliable guide to changes in the amount of oedema. Admission to a nursing home or hospital for several weeks

is therefore almost an essential, and during this time the patient's relatives, if intelligent, may be trained to continue the treatment later at home. In nearly every instance the patient should be confined to bed throughout the illness. This rule should only be relaxed if practically all the oedema has disappeared and the only remaining evidence of disease is albuminuria which every possible effort has failed to clear up.

Owing to the danger of infections of all kinds the greatest possible care should be taken to guard against exposure to cold and contact with persons who may be carrying infection, and scrupulous attention must be paid by the nurse to the cleanliness of the skin and mouth. If there is reddening of the oedematous skin over the pressure points, a water bed should be obtained and a cradle to keep the weight of the bedclothes from the patient's legs and feet. Oedematous patients are usually more comfortable when supported by a bed rest, and this has the advantage of encouraging the oedema to collect in the more dependent parts.

Dietetic Treatment

The principles of dietetic treatment differ in some important respects from those applicable to other types of renal disease. The constant loss of considerable quantities of protein in the urine leads to a depletion of the protein of the blood plasma. This, as Epstein has shown, is an important factor in the causation of the oedema, and an adequate supply of protein must therefore be ensured in the diet. In nephrosis this is not contraindicated by any consideration of renal inadequacy, because, as already stated, the renal efficiency is maintained. Epstein introduced in 1917 the high-protein, low-fat diet for nephrosis, and although it has not produced the uniformly successful results which were hoped for, it will provoke diuresis and reduce oedema in a fair number of cases, and is worth trying at intervals during the treatment.

Whatever the cause of the oedema it represents a retention in the body of salt and water which the kidney is failing to excrete. Salt and water must therefore be given very sparingly, as any excessive ingestion of these only increases the amount of oedema present.

The following diet fulfils the desired features of restriction of salt and water, with liberal protein and minimal fat.

Breakfast.—5 oz. tea or coffee with skimmed milk. 4 oz. fish. Bread. Marmalade.

Mid-morning.—One small orange or apple.

Lunch.—Oysters if procurable. 4 oz. lean meat or chicken. Peas, beans, or lentils. Milk pudding. 3 oz. fluid.

Tea.—5 oz. tea with skimmed milk. 2 oz. fish or lean meat. 6 oz. salad or fruit.

Supper.—4 oz. fish or lean meat. Vegetables, salad, or mushrooms. 5 oz. tea or coffee with skimmed milk.

No salt to be cooked or eaten with the food. No extra fluid.

This diet may be taken for a month or two at a time, provided the appetite of the patient is good. In the intervals an ordinary diet may be ordered containing as much protein as the patient wishes, but with the fluid content restricted to about one pint per day for an adult. In no case will salt or salted foods be allowed.

For a child the fluid allowance will be restricted in proportion to the average weight for his age, the actual weight being unreliable on account of oedema. To allow fish and to exclude meat as a source of protein has no

scientific advantage, and merely reduces the variety and palatability of the diet. The records of the case will show whether the high protein diet is causing an increased diuresis and reduction of the body weight (oedema).

Drug Treatment

There being no drug known to influence the course of the pathological process in the kidney, the drug treatment of nephrosis consists essentially in a further attack on the oedema. Of the many diuretics which have been used in this disease two groups only need be discussed in detail.

1. Urea is an efficient diuretic if given in sufficient quantity. At least 15 grams, dissolved in 100 c.cm. of water and flavoured with tincture of orange, should be administered three times daily. Diuresis will be noticeable within three or four days if the treatment is successful, and the drug may be given for several weeks at a time, though it sometimes has to be discontinued on account of thirst or nausea.

2. In obstinate cases of nephrotic oedema, when other measures have failed, the mercurial diuretics may be employed, always provided that the essential criteria of the diagnosis have been fulfilled—absence of haematuria, high blood pressure, or nitrogen retention. This group of drugs acts more efficiently if the patient has received some days beforehand one of the acid-producing salts. One may therefore prescribe:

R. Ammon. chlorid.	...	...	...	gr. xxx
Ext. glycyrrhiz. liq.	...	...	...	℥ xxx
Aq.	...	...	...	ad. 3 j

3 j t.d.s.

On the second or third day the mercurial drug is given. This may be either salyrgan, 1/2 to 1 c.cm. intravenously, or neptal or salyrgan, 1 c.cm. intramuscularly, or a novurit suppository per rectum. The ammonium chloride can be continued and the diuretic drug repeated every third or fourth day for a dozen or more injections. If the intravenous route is chosen extreme caution must be used to prevent even the smallest drop of salyrgan escaping into the subcutaneous tissues as the drug is highly irritating and may cause intense swelling, pain, and redness at the site of the injection. It is even as well, after taking the salyrgan into the syringe, to draw in a drop of sterile water, so that if there is any leakage from the needle as it pierces the skin only water will be introduced. A little blood is withdrawn to show that the needle is in the vein, the tourniquet is loosened without a jerk, and the injection is administered slowly. If the needle is in position it will be quite painless, and the patient will "taste" the drug within a few seconds. Before the needle is removed, another drop or two of blood is withdrawn into the syringe to rid the needle of any trace of salyrgan. If the intramuscular route is chosen a needle at least 1½ inches in length is inserted perpendicularly into the muscle of the upper and outer quadrant of the buttock, or into the deltoid muscle if this is sufficiently thick. If intramuscular injections are painful a little novocain may be given mixed with the salyrgan or neptal. The suppositories are less efficient but of course easier to administer. They occasionally give rise to diarrhoea. Diuretics of the caffeine group, such as diuretin and euphyllin, are of less use in nephrosis, but may be given if urea is not tolerated and injections not desirable, for instance in children.

Apart from diuretics, the following medicaments have some place in the treatment of nephrosis. Iron should be prescribed if anaemia is present. This should be judged by an actual haemoglobin estimation, as the characteristic pallor of the disease is due not to anaemia but to oedema. Thyroid extract or thyroxine is extremely well

tolerated in nephrosis, and appears to improve the general condition and encourage diuresis in some cases: 5 to 10 grains of the extract, or 1 mg. or even more of thyroxine, per day may be safely prescribed if the pulse is watched for any undue increase in rate. Finally, constipation must be avoided, and if an aperient is necessary magnesium sulphate is the most suitable, as it promotes the withdrawal of water without irritating either colon or kidney.

Mechanical Treatment of Oedema

If the pleural cavities are involved the fluid should always be removed, because to do so relieves respiratory distress, and an unexpected empyema may occasionally be found. Ascites should only be tapped if very large and giving rise to much discomfort. Subcutaneous oedema should never be relieved by Southey's tubes or by incisions unless it is very distressing to the patient and diuretics have failed to reduce the oedema sufficiently. The reason for this advice is the danger of a spreading cellulitis resulting in the oedematous tissues. An exception may have to be made if swelling of the genitals renders it difficult for the patient to pass urine. In that case a number of pricks can be made in the penis with a sterile sharp surgical cutting-needle.

Removal of fluid may be attempted via the sweat glands by either hot packs or radiant heat baths or an electric blanket. It is difficult to produce diaphoresis in grossly oedematous patients, and the strictest precautions must be taken to avoid burning or blistering the skin. For these reasons the methods are not of much practical value.

Treatment of Infections

The only established fact in the aetiology of nephrosis is that it is a disease which has some association with infective conditions in the body. Thus it may develop in connexion with infections of the antra and sinuses, mastoiditis, otitis media, inflammatory conditions of the skin, or more deep-seated diseases such as appendicitis or cholecystitis. Its association with tonsillitis is less obvious than is the case in nephritis. The frequency of septic complications has already been mentioned, and the likelihood of the pneumococcus being the infective agent is striking. Thus pneumococcal empyema, peritonitis, pericarditis, pneumonia, and meningitis are the common causes of a fatal issue. Parotitis is another complication which may prove fatal. These are treated on the principles applicable to the particular case, and though such infections may prove to be terminal the recovery from a pyrexial complication has been known to be closely followed by complete recovery from the renal disease as well.

Because of this association with infections it is extremely important that a careful search should be made early in the disease for any active focus of sepsis, and if such be found a suitable operation should be performed. Thus an antrum must be drained, a mastoid operation performed, and even the appendix removed if the indications are sufficiently definite. Although such operations are not to be lightly undertaken, they represent the only possibility of curative therapy—that is, of the removal of the cause of the disease—and it is extraordinary how well an oedematous patient will often come through operative procedures of this kind.

Finally, a few words may be said on the general outlook. Nephrosis is usually a fatal, and always a serious, disease. It is, however, subject to spontaneous remissions, often sudden ones at unexpected times, and in some cases these may become complete, the patient recovering entirely after many weeks or months of illness. However unfavourable the position may appear, therefore, it is always worth continuing the therapeutic attack, and thus increasing the likelihood that the patient will survive long enough for a natural remission to appear.