BRITISH MEDICAL JOURNAL

LONDON: SATURDAY, OCTOBER 3rd, 1936

PATHOLOGICAL CHANGES IN BONES AND JOINTS INDUCED BY INJURY*

BY

JAMES F. BRAILSFORD, M.D., PH.D., M.R.C.P.

HONORARY RADIOLOGIST, QUEEN'S HOSPITAL, BIRMINGHAM ; RADIOLOGICAL DEMONSTRATOR IN LIVING ANATOMY,
UNIVERSITY OF BIRMINGHAM

(With Special Plate)

Fractures

The most obvious lesion resulting from injury to bone is a fracture. The extent of the fracture will depend upon the degree of trauma, the angle at which it is applied, the weight or stresses which the fractured bone is supporting at the time, and the fragility of the bones. While in the majority of cases the fragments are separated so much that a radiograph taken in any of the usual planes will provide definite evidence of the fracture, in some cases little or no separation results, and radiographs in one or more planes fail to reveal any sign of the lesion. It is therefore essential that all patients x-rayed for evidence of fracture should have radiographs taken in two or more planes. Where possible, one plane should be at right angles to another. Further radiographs should be taken immediately after the injured limb has been immobilized, as the good position shown on the primary radiographs may have been lost during the process of immobilization.

Value of Serial Radiographs

The stage of healing and some idea of the time when the limb can be subjected to any of its normal stresses and strains can be shown by radiographs taken at intervals of one to two months. Serial radiographs show that gradually the fractured ends lose their sharp contours with some diminution of their density, while bridging the gap between the fragments faint shadows of callus appear which slowly increase in density. The fractured ends become rounded off, the callus sweeps over them, more and more calcium is deposited until the whole assumes the density and trabeculation of normal bone, and ultimately the site of the fracture has the appearance of a plumber's wiped joint, in which all detail of the broken ends is completely obliterated. Such is the sequence in the uncomplicated case which can be radiographically recorded in two to four months.

Sometimes, notably in fractures of the neck of the femur, the carpal scaphoid, the shaft of the humerus, and at the junction of the middle and lower thirds of the tibia and fibula no evidence of union may be detected. Much callus may be shown surrounding the fractured surfaces of the humerus, tibia, or fibula, but the gap is not bridged and no trabeculae can be seen passing from one extremity to the other. The callus on each extremity may be splayed out and present an irregular sclerosed border to its fellow. Such an appearance may persist for years and a false joint may even develop at the site which permits of movement and angulation of the fragments. Atrophy and absorption

may occur in any fragment of bone, especially the head of the femur, which has been deprived of its physiological function as a result of non-union. As long as it is possible to detect the extremities of the fragments and impossible to see evidence of bone trabeculae bridging the gap one must conclude that consolidation of the fragments has not taken place.

Fate of Bone Grafts

Attempts to unite the fragments may be made by fixing a bone graft across the gap. Serial radiographs will show the ultimate fate of the graft. It may be:—(a) vascularized, and become an integral part of the bone; a successful graft gradually becomes decalcified to such a degree that it assumes the density of the fractured bone, and after fusion has taken place and the graft is endowed with physiological functions it gradually increases in density by the storage of calcium and assumes the structure and outline of the normal segment: (b) it may be gradually absorbed, and then the radiograph will indicate a gradual erosion and disappearance of the graft; the explanation for this phenomenon has not been definitely established: (c) the graft may retain its calcium, which will be inferred from the fact that serial radiographs show its density and sharpness of outline to remain unchanged; such a graft is ultimately discharged through a sinus or has to be removed surgically.

In those cases of compound fracture complicated by sepsis the radiograph will indicate extensive osteoporosis of the living bone unless its vascularity has been impaired, but any fragment of bone which has been deprived of its blood supply will retain its density and sharp outline. These characters enable us to detect the sequestrum from the living fragments which show the same diminution of density and rounding off of their surfaces as is exhibited by the main fragments. Such are the radiographic findings in the majority of fractures. There remain a minority which show extraordinary changes.

Kuemmell's Disease

In my paper* on "Deformities of the Lumbo-sacral Region of the Spine" I pointed out that in some cases radiographs of patients who had received injuries to the spine did not show any detectable damage if the radiographs were taken within a day or so of the injury, but if radiographs were taken two to four weeks after the injury had been sustained, evidence of damage to one or more vertebral bodies might then be present. If the injured spine has not been relieved of pressure or strain a degree of collapse of the affected vertebral body may occur. It is

* Read in opening a discussion in the Section of Radiology at the Annual Meeting of the British Medical Association, Oxford, 1936.

* *Brit. Journ. Surg.*, 1928-9, xvi, 562.

probable that Kuemmel's disease, which is described as a non-tuberculous rarefying osteitis, has its origin in this way. The explanation of the changes in the radiographic appearances may be that the injury caused damage to the cancellous tissue which could not be detected until absorption of the damaged trabeculae had permitted collapse and produced an alteration in the outlines of the vertebral body. Consolidation of the fragments may proceed normally, but in some cases there is considerable delay and increasing deformity.

Somewhat similar changes are to be seen in the carpal scaphoid. In this bone we have the advantage of closer apposition to the radiographic film, consequently the detail of the cancellous structure is more clearly defined and fine fissures may be detected on careful inspection. But even here damage may not be revealed until the damaged trabeculae have been absorbed and a cyst-like area has been formed at the site. At this stage a fissure through the wall of the "cyst" may be shown.

Another site in which abnormal changes follow trauma is the wall of the acetabulum. Though radiographs taken soon after the injury may show no definite change, a progressive protrusion of the acetabular roof into the pelvis develops as the result of pressure on the damaged bone, which now appears to have become somewhat plastic. This is one of the causes of the lesion referred to as "protrusio acetabuli."

Fractures of the femoral neck fail to unite in a relatively high percentage of cases. The bone in the region of the fracture undergoes gradual absorption leaving a gap between the fragment bearing the femoral head and the lateral extremity of the neck. More recently an added percentage of successes has been recorded following the fixation of the fragments by the Smith-Petersen pin and other devices.

Wiring and Plating

Serial radiographs of bones which, following fracture, have been held in alignment by metal plates, wires, etc., show that good apposition of the fragments and often union may be secured thereby, but that as long as the metal is retained the bone will not completely recover its normal strength and elasticity. The appearances suggest that a chronic inflammatory process is induced by the foreign body which leads to plasticity and bending of the bone in spite of the metal attachments, which may either be torn out or bent with the bones. Such radiographic findings, which have been seen to arise in bones several years after the surgical introduction of the metal, suggest that as soon as radiographic evidence of union between the fragments can be obtained, the metal attachments ought to be removed, if ultimate effective consolidation and healing is to take place.

The junction of the middle and lower thirds of the tibia and fibula when fractured in some infants appears to be very liable to non-union even after repeated bone grafts, and a pseudo-arthritis develops at the site of which the lower fragments are angulated backwards at about 90 deg. Compensatory changes occur in the bones of the foot as a result of the abnormal positions which they are thus forced to attain. Radiographs suggest that the lesion is due to a congenital defect in ossification at the site of fracture, for in some cases cyst-like areas with defects in the cancellous structure may be detected in this neighbourhood before actual fracture has occurred.

Osteochondritis

Until the beginning of the present century all localized rarefying conditions of bone were regarded as due to tuberculous infections. Many of them were undoubtedly of this nature, but it was noted that in a considerable per-

centage of such cases the clinical course of the condition was not such as would be expected if it actually was due to tuberculous infections. With the routine use of x-ray investigation it is now possible to follow these cases and to study the living pathology. Gradually, doubt has given way to certainty that many of the localized rarefying changes in bone, affecting the primary and secondary centres of ossification, are definitely non-tuberculous in origin. Such localized rarefying bone changes have now been described in many centres, for example:

Kuemmel (1895) has written on the spine; Osgood (1903) and Schlatter (1908) on the tubercle of the tibia; Köhler (1908) on the tarsal scaphoid and patella; Legg, Calvé, Perthes and Waldenstrom (1910) on the epiphysis of the head of the femur; Kienbock (1910) on the semilunar; Preiser (1910) on the carpal scaphoid; Sever (1912) on the os calcis; Freiberg (1914) on the head of the second metatarsal; Scheuermann (1921) on the spine; Renander (1924) on the medial sesamoid of the big toe; Van Neck (1924) on the pubis; Buchman (1925) on the iliac crests; Voltancoli (1925) on the ischio-pubic junction; Calvé (1925) on the spine; Friedrich (1924) on the clavicle; Lewin (1927) on the shoulder; Mauclore (1927) on the heads of the metacarpal bones; Mouchet (1928) on the astragalus; Wagner (1930) on the epiphysis of the first metatarsal; and Burns (1931) on the lower ulnar epiphysis. I have recorded these changes in the pedicles of the fifth lumbar vertebra, the adult scaphoid, the femoral neck, the accessory scaphoid, and the roof of the acetabulum.

The number of sites in which bony changes of this type are recorded is added to every year, and it seems to be generally accepted that the underlying changes are of the same nature, that is, an osteochondritis. Though we have extended our knowledge as to the sites in which these changes may be found, agreement has not been reached as to the pathology of these bone lesions. After carefully weighing the evidence obtained from over 100 cases personally observed, I am of the opinion that trauma is the exciting cause of these conditions, but in view of the fact that it is the minority only of the cases of trauma in which the sequence of characteristic changes develop, I feel that there must be some other contributory factor. Stunted development following changes resembling those of osteochondritis are often seen to follow manipulative reduction of congenital dislocation of the hip-joint. Such changes have never been seen in the dislocated femoral head where reduction has not been attempted. It is impossible in this paper to define the serial radiographic changes found in osteochondritis; I must refer you to a previous paper.

Ossification of Haematomata

Injury to the surface of a bone may be associated with the formation of a haematoma which will not be revealed by a radiograph in the early stages. Some haematomata are prone to calcification in which subsequent ossification may occur. In the early stages of calcium deposition radiographs will show a cloudy shadow which may somewhat blur the outline of the involved bone, which at this stage often shows a localized osteoporosis. As these radiographic appearances are associated with a tumour devoid of the clinical signs of inflammation in a patient who may not remember the injury which caused it, it is not surprising that the suspicion of sarcoma has been aroused and amputation performed before the error was discovered. Such radiographic appearances should indicate that a further x-ray examination ought to be made after an interval of one or two months. It will then be seen that the abnormal shadow has lost its cloudy appearance and is taking on a fairly well-defined outline. Ultimately the appearance of new bone will be shown. This new bone

may appear as (1) an irregular boss on the periphery of the shaft; the lateral surface of the middle third of the femur and the upper third of the humerus being the most common sites: (2) a fusiform deposit around the shaft of a bone such as is seen in the march fracture of the second metatarsal: (3) irregular spurs in the neighbourhood of a joint, particularly the elbow and other big joints—hip, shoulder, knee or ankle—which have been dislocated: or (4) bands of new bone stretching over and fixing neighbouring joints; the hip-joint area is the most common site for these changes.

In certain cases calcium is densely laid down in the haematoma and the latter then appears on the radiograph as a dense granular structure having well-defined rounded borders. If the trauma injures the periosteum and such blood as is extravasated is absorbed, subsequent radiographs may indicate the site of the trauma by a localized thickening or sclerosis of the cortex.

Cyst-like Changes after Trauma

Injury to cancellous bone which can be studied in radiographs of the hand and wrist may not be sufficient to cause a breach of continuity of the surface of the bone, but may yet lead to a localized absorption of the damaged trabeculae and produce a cyst-like appearance. I have found such appearances in the heads of the metacarpals and in all the carpal bones of men who had worked for ten years or so with compressed air drills (see Plate, Fig. 1). But the commonest site for this development to be seen is in the carpal scaphoid to which reference has been made in a previous paragraph. The larger areas of cancellous destruction with cyst-like development in the head and neck of the humerus, which are sometimes referred to as localized osteitis fibrosa cystica, are considered by some authorities to be due to trauma, for on histological examination the contents often resemble the structure of old blood clots. The simple cysts, which are most commonly found in the shaft of the humerus and in the neighbourhood of the trochanteric region of the femur, are also thought to be due to trauma. Such lesions are usually brought to light by so-called spontaneous fracture—that is, fracture due to trauma of relatively trifling degree.

Changes in the Joint Space

Many injuries to the articular cartilage must occur which do not produce any appreciable alteration in the radiographic appearance of the bones entering into the joint. In some cases in which the injury has been followed by effusion into the joint it may be possible by comparative examination of radiographs of the joints on both sides to recognize an increase in the joint space on the affected side. For the purposes of description the gap between the surfaces of the sub-articular bone is usually referred to as the joint space, but it is recognized that in the normal joint this gap represents the thickness of the opposing articular cartilages which are in contact with one another. With effusion into the joint the surfaces of the articular cartilages are separated and the gap between the bones is accordingly increased. On the other hand, in those joints in which atrophy or destruction of the articular cartilage has occurred the gap between the sub-articular bony surfaces is reduced or obliterated. Such reduction in the joint space is rarely seen to follow injury until several weeks or months have elapsed if the joint is not infected, even though movements of the joint may be painful for a month or more after the injury.

Thus it is unusual to be able to detect radiographic changes in the joints of active footballers though these may have been subjected to repeated trauma of a severe degree. The physically fit man seems to be able to endure

such injuries without his bones and joints registering any appreciable radiographic change. Nevertheless it would appear from radiographic examinations that certain joints which have been so injured yet given insufficient rest to completely recover not infrequently develop degenerative changes during middle life. These take the form of diminution of the joint spaces indicating atrophy of the articular cartilage, particularly where this is subjected to the greatest strain.

In more severe degrees, such appearances as pointing of the articular margins, and osteophytic development around them, the formation of loose bodies within the joint, eburnation, and erosion of the bony surfaces which have been brought into contact may be detected—in fact, all the degenerative changes which we associate with the term osteoarthritis. Such a sequence of changes does not always take place, and this suggests that some other factor must be present.

Aetiology of Degenerative Changes

In a previous paper I suggested that these degenerative changes were probably due to the products of faulty metabolism or to the effect of bacterial toxins acting over a long period on joint surfaces which had been injured, for osteoarthritis is only manifested in those joints which have been subjected to abnormal stress or strain in a person who also exhibits elsewhere the signs and symptoms referred to as rheumatism, fibrositis, etc. Such findings suggest that the persistence of pain in an injured joint which does not yet show any radiographic departure from the normal indicates faulty metabolism or absorption of toxins: the discovery, correction, or eradication of the latter has resulted in the disappearance of the symptoms and the prevention of those pathological changes in the joint to which reference has been made.

Once these degenerative changes have been indicated in a joint, serial radiographic examinations will show a slow progression of the lesion with gradual limitation of the range of movement due to destruction of the articular surfaces: but bony ankylosis never occurs unless induced by surgical resection of the articular surface and close approximation of the bared bony surfaces. Rest appears to be the only factor which checks the progress of the lesion or leads to its repair. If the injured joint is devoid of normal trophic influences, as in syringomyelia, tabes dorsalis, spina bifida, etc., normal regeneration appears to be checked or abolished and the joint surfaces with the underlying bones are slowly and painlessly worn away, the liberated calcium often appearing as an amorphous deposit in the neighbourhood. These changes are frequently associated with effusion into the joint and signs of oedema in the neighbouring tissues. Such changes are not confined to the anatomical joints. They have been seen to develop at the site of fractures of the shafts of the long bones. It is the development of changes of this nature in such pseudo-arthroses and in joints other than the hip, shoulder, knee or elbow which may give rise to difficulties in diagnosis. Such changes have been mistaken for those of secondary neoplasm.

Subluxation and Dislocation

Subluxation or dislocation of any joint may be produced by trauma. In some cases the ligamentous and muscular control may permit of easy displacement of the joint surfaces and in these cases recurrent dislocations are common. Easy displacement usually permits of easy reduction and therefore secondary changes due to trauma do not readily appear. Normally developed joints usually require severe trauma before they can be dislocated, and in some cases it would seem that reduction is accomplished by the exercise of force which subjects the part to trauma of

even greater severity. In some of these cases radiographs show that the rim of the joint has been fractured in the procedure. In others the radiograph of the reduced joint gives no indication that the part has been subjected to trauma, all the joint surfaces and surrounding tissues appearing quite normal. But it is unwise to base an opinion of the severity of an injury on the radiographic appearance of the part soon after reduction. Serial radiographs taken at three-monthly intervals will demonstrate the truth of this. Depending on the severity of the injury, changes will gradually be shown on the radiographs.

The first sign may be a relative osteoporosis of the ends of the bones entering into the formation of the joint. The damaged articular surfaces may become blurred in outline, and hazy ill-defined amorphous deposits of calcium may be seen in the neighbourhood. Later radiographs will show organization of this calcium in the form of irregular strands or spurs of new bone, some of which may appear to have firm attachment to the bony surface, others lying in the intermuscular planes. The joint may ultimately be bridged by a band of new bone which effectually prevents all movement. In other cases the lesion shows the gradual development of the typical appearance of an osteoarthritic joint. It may take several years before we can estimate the extent of the damage which the joint has sustained (see Figs. 3-5).

The Intervertebral Disks

Injury to an intervertebral disk may not be apparent for several months after the trauma. It may then be indicated by diminution of an intervertebral space and lipping of the adjacent margins of the vertebral bodies, and perhaps partial localized ossification of the anterior common ligament. If, in addition, the articular facets have been involved, some degree of curvature may arise, and, after a few years, the articular processes may show irregular sclerosis, most marked at the site of the injury, but also indicated in the bodies on the concave aspect of the compensating curvatures, above or below the site of the injury. Damage to the cartilaginous plates may result in herniation of the nucleus pulposus into the affected vertebral body.

The disk may be injured by the needle used in lumbar puncture, performed for diagnosis, treatment, or anaesthesia. Following such a procedure the patient complains of a persistent backache, and serial radiographs will show a progressive reduction in the disk space with some osteoporosis of the adjacent vertebral bodies. Ultimately localized ossification of the anterior common ligament occurs with sclerosis of the adjacent surfaces of the vertebral bodies. R. W. Billington, reporting on spondylitis following cerebro-spinal meningitis, stated that twelve of fourteen cases he studied showed pathological changes in the lumbar region. Because of this he conjectured that the vertebrae were affected by trauma caused by the needle and by direct inoculation with the meningococcus from the passage of the needle through the infected spinal fluid.

B. J. Alpers has recorded the clinical and radiological findings occasioned by chondromata of the intervertebral disks, cartilaginous extensions of the intervertebral cartilages into the spinal canal. He states that they are commonly to one or other side of the mid-line, usually the left, firmly anchored to the disk, and causing pressure on the anterior aspect of the cord. They have been found in the lumbar and cervical regions; they appear to be related to trauma. The symptoms are slowly progressive and recovery follows removal. No indication of their presence may be given on the straight x-ray, but following lipiodol injection an obstruction will be indicated by the hold-up of the opaque injection.

Post-traumatic Osteoporosis

The bones of a limb which is immobilized following trauma or a localized surgical operation show a gradual and somewhat irregular decalcification even of those bones which were not involved by the injury. Serial radiographs of the bones often show very marked changes not only in the calcium content but in the cancellous and compact structure of the bone. The cancellous extremities of the long bones, particularly in the neighbourhood of the epiphyseal line, though the epiphysis may have been fused with the diaphysis for years, show a marked decalcification and consequent radio-transparency. With the same exposure as for a normal limb the radiograph will appear to be considerably over-exposed. With the time reduced sufficiently to give a correct exposure for the decalcified bone the radiograph will show a compact cortex to the shaft which gives the false impression of increased density and calcium content. An examination of the structure of the bone in a serial collection of radiographs of an immobilized limb will reveal not only irregularity in density of the compact tissue in which there may be cortical islands which are very radio-transparent, but very marked changes in the cancellous structure. Coarse and thickened trabeculae enclosing larger meshes appear to replace the fine and exquisite reticulation of the normal cancellous tissue. These appearances may persist for a year or more after immobilization of the limb has terminated, but eventually the delicate architecture of the normal is restored (see Special Plate, Fig. 6). There is a class of case, however, in which these retrograde changes show a progressive development in association with certain well-marked clinical signs and symptoms.

Following trauma, particularly in the neighbourhood of the multi-articular joints of the wrist and foot, but also in the neighbourhood of the larger joints such as the knee and shoulder, a sequence of symptoms develop in some patients which may give rise to the diagnosis of tumour or tuberculosis. The trauma may in itself not occasion any great pain or disability: even that which is present soon after the injury may gradually disappear. Following a period of quiescence in the affected part or in the extremity distal to the injury a painful and disabling condition may develop which is not relieved by rest and often resists all electro-medical applications. The affected part appears turgid, its skin thin, shiny, and red or bluish, and often moist. Movement of the part is painful. Serial radiographs taken during the development of the condition reveal a progressive decalcification of the cancellous tissue, at first as small circumscribed foci in the cancellous extremities of long bones, or in the short tarsal or carpal bones. The compact tissue of the shaft may appear to contrast with the decalcified extremities, but ultimately this, too, may show a progressive decalcification, at first rather irregular in its distribution, giving the bone a mottled appearance.

This condition, which was first described by Sudeck, is often referred to as "acute bone atrophy," a name which unfortunately does not readily occur to observers of this type of lesion, for it may persist for a number of years. As the condition develops in but a small proportion of patients who have suffered traumata, some other factor such as focal sepsis or intoxication may be the sensitizing agent. The radiographic appearances during the progress of the lesion suggest that the bone changes are due to some disturbance in the trophic regulation of the part, and the reaction towards improvement, at any rate for a short time, which seems to follow sympathectomy, appears to support this.

Where such changes follow traumata which have necessitated immobilization, the radiographic appearances will be regarded as being due chiefly to disuse of the part and consequently will not occasion any serious reconsideration of the diagnosis, but in those cases in which the initial traumata have been given little or no prominence in the clinical history the progressive clinical signs and abnormal radiographic appearances may give rise to very serious errors in diagnosis unless the observers are familiar with the condition. In some cases these clinical and radiographic signs have led to such grave diagnoses as bone tuberculosis, sarcomata, or other malignant changes, and even amputation has been advised, yet in other cases the slow progression of the lesion, the vasomotor disturbance, and the appearance of disuse atrophy which accompany them have been regarded as evidence of malignancy. Judging from the radiographs which have been referred to me for an opinion, it is particularly in the region of the larger joints, the knee, shoulder, and elbow, that these changes in the bone have been most confusing. A number of these cases have been observed over several years during which little or no improvement has been recorded on the radiograph and the affected limb has become more and more disabled, particularly in those patients who have lost faith in the benefits of physical medication. Turner claims to have obtained good results by iodine injections, while Middleton and Bruce found favour with acetylcholine. To instance the consequences of misinterpretation of such lesions the following brief account of a case submitted to me for an opinion is given:

CASE REPORT

The patient developed a painful elbow and was sent for a radiographic examination. The radiographs revealed the changes seen in traumatic arthritis, but these were attributed to tuberculosis and other more serious conditions for which he received, over a period of three years at the hands of various men in this country and many places on the Continent, courses of ultra-violet irradiation, Hertzian irradiation, deep x-ray therapy, injections of tuberculin, iodine, and mercury, and an exploratory operation. In spite of all these interfering irritations the lesion presented surprisingly little change in the radiographic appearances and with a complete rest from all treatment I suggested that it might resolve.

Trauma and Focal Inflammatory Changes

It has been established experimentally that localized trauma to an animal with pathogenic organisms circulating in its blood can initiate the development of a bone abscess at the site of the trauma. Corroboration of this experimental evidence is abundantly obtained from examination of the histories of patients exhibiting localized acute or chronic inflammatory lesions. Many cases of acute septic abscess, tuberculosis, or gummata of bone present clinical evidence of a primary septic, tuberculous, or syphilitic focus apart from the bone lesion which often appears to have originated at the site of a definite injury. Because localized injury dominates the clinical picture in the early stages of the lesion, the superadded pathological changes may escape discovery for weeks or months. Indeed, in some cases, though the nature of the lesion may now be obvious, it is not recognized except by a fresh observer who is not obsessed with the traumatic evidence. Serial radiographic examinations of injuries to bone will give the earliest indication whether healing is proceeding with normal regularity, or whether it is complicated by some underlying pathological process. In some cases the nature of the process will be suggested by the radiographic appearances.

Trauma and Malignant Disease

It has long been thought that malignant disease could be initiated by injury, but until we know the other essential factors concerned in the metaplasia which occurs it will be impossible to prove the theory definitely. We know that there must be other essential factors besides trauma because it is only a very small proportion of injuries which are followed by malignant metaplasia. The statistics of the German Army and Navy show that 1 in 15,000 injuries is followed by malignant changes. Many cases of malignant disease seen in practice appear to have a definite association with previous injuries, therefore it is not surprising that many medical men hold that there is a distinct relationship. The belief appears to have an even wider following amongst the general public, in whom the question of injury is immediately raised when a diagnosis of malignant disease of bone is made. Cases that I have seen have usually shown at the first radiographic examination evidence of definite changes in the bone of the injured limb.

When taking the history, the patient may state that the part was severely injured a few weeks, months, or years previously and there may be a scar to indicate the site of the injury. In two instances which I observed, the patient had had a gun-shot wound with fracture at the site of the malignant tumour of bone; in another case, an osteotomy had been performed. Such histories seem to lend strong evidence to the theory. Three cases of endothelial myeloma of bone in young people who subsequently died of metastases in other bones and viscera gave definite histories of previous injuries.

CASE REPORT

A youth, aged 15 years 9 months, told me that at Easter, 1934, a heavy weight which he was supporting fell on his upper arm. He was taken to hospital but was not x-rayed for three weeks. The film evidently did not arouse the suspicion of malignancy, for after the taking of the radiograph (April 24th, 1934) he was given electrical treatment and massage for several weeks, until a further radiograph (June 14th) revealed additional changes in the shaft of the humerus which were now regarded as evidence of malignancy and for which amputation was advised but refused. It was at this stage that I was asked to report on a further radiograph, and the diagnosis of endothelial myeloma was made. We administered x-radiation, to which the tumour readily responded; so well in fact that the patient did not come to hospital for further treatment for several months, when the arm had again increased in size. Further radiation was given with equal response, after which nothing further was heard of the patient until he was admitted into another hospital, in which he died in February, 1936.

In this case the history of trauma was very definite, but unfortunately no radiograph was taken until three weeks after the injury, and this radiograph, which I have since been able to see, showed a fracture of the humeral shaft with very definite evidence of malignant changes in the bone.

But it is not only in primary malignant disease of bone that trauma appears to play a part. The histories of such cases as the following suggest that trauma may determine the site of metastases.

CASE REPORT

A patient sustained a severe injury to the buttock when knocked down by a car. When called to x-ray him the next day with a portable unit, I found that he had a very large bruise over the buttock, but the radiograph did not reveal any changes in the bones or injury to the pelvis. The patient apparently recovered from the injury, but six months after this he consulted me on account of a pulsating tumour in the buttock at the site of the contusion sustained at the

accident. He said that he had first noticed the pulsation during the previous week when he had been sitting for a long period on a hard seat, but he regarded it as an aneurysm of the gluteal artery and sought radiographic confirmation. The radiographs showed an area of cancellous destruction with an ill-defined margin. At the post-mortem a few months after, a secondary hypernephroma was found at the site of the tumour.

Further instances of this nature tend to encourage a belief in the association of trauma and malignancy.

Review of the Literature

In the literature there is much to support the belief. Second in 1907 read a paper before the French Congress of Surgeons in which he put forward certain conditions which he considered should be fulfilled in any case of malignant disease deemed to be due to trauma. They include:

- (1) Definite evidence of severe trauma within a reasonable time to a part which previously had appeared to be quite sound.
- (2) The growth of the tumour at the site of the previous injury.
- (3) The establishment of the diagnosis of malignancy by clinical, radiological, and histological examination.

W. B. Coley presented a paper before the New York Medical Society in 1897 in which he indicated that forty-six out of 170 cases of sarcoma gave a definite history of trauma. This worker and his associate, N. L. Higginbotham, presented a paper before the American Surgical Association, May 10th, 1933, in which they put forward a very strong case in support of the traumatic theory of malignancy. They claim that no fewer than 280 cases of sarcoma which they had personally investigated were associated with antecedent local trauma. An account of forty-six cases of sarcoma following war fractures and details of individual cases of malignant disease with antecedent severe injuries was also given. They further stated that

"the interval of time elapsing between the injury and the appearance of the tumour is often much shorter than is recognized by most writers. In the majority of cases the tumour develops within the first month or six weeks of the injury, but in a considerable number of cases it may develop within one or two weeks. The latter cases justify the classification of acute traumatic malignancy originally suggested by the English surgeons."

Coley considers that trauma creates a "favourable soil" for the development of a ubiquitous cancer organism. He states that "the micro-organism finds a suitable soil in the damaged cell, forming a symbiosis with the cell and causing a proliferation and multiplication resulting in a malignant tumour." But if such organisms exist everywhere it is very surprising that so few injuries are followed by malignant changes. These two factors alone cannot decide such developments.

Ewing is of the opinion that "mechanical trauma is an important factor in the causation of tumours." Its effects are: (1) solution of continuity, minute and gross; (2) separation of cell groups and tissue masses, as of skin, glands, bone; (3) necrosis of tissue; (4) confined haemorrhage requiring absorption or encapsulation; (5) accelerated regenerative processes with hyperaemia and new growth of specific cells, blood vessels, and supporting tissues; (6) cicatrization.

The percentages of malignancies with histories of antecedent injuries which are recorded by different observers vary considerably. The enthusiast for the theory is satisfied with evidence of trauma which would be rejected or ignored by others. Until we can produce at will the other essential conditions we shall not be able to determine the exact significance of trauma in malignancy.

Occupational Injuries

Those joints which, as the result of particular occupations, are subjected to abnormal stresses and strains show chronic arthritic changes. The latter gradually reduce the range of movement of the affected joint, and the workman is forced to acquire the ability of manipulating his tools with restricted joint movements. Ultimately the chronic arthritic changes produced by constant trauma may lead to incapacity, but this eventuality may be produced earlier in a joint already exhibiting evidence of traumatic arthritis by trauma of a degree unlikely to affect seriously a normal joint (see Figs. 2 and 7). Damage to bone and joints caused by occupational strain has not received its due acknowledgement under the Workmen's Compensation Act, and with the present law it sometimes becomes very difficult to assess the amount of responsibility when a workman, whose joints show evidence of severe occupational strain, claims that he has been incapacitated by a specific injury. By a law introduced in Germany in 1929 pathological changes in joints of workers using drills, hammers, etc., driven by compressed air, have been classified as an occupational disease. Following on the trauma any of the conditions which have been described in the preceding paragraphs may develop.

I would like to take this opportunity of stressing again the importance of an x-ray examination, as soon as possible after the accident, of any part which has been injured. Particularly does this apply to persons who are covered by the Workmen's Compensation Act. Such a radiographic record may prove invaluable, for not only will it register the condition of the bones and joints immediately after the accident and be a guide to treatment, but at a later date it may be used to compare with radiographs taken subsequently. A comparative examination of this nature will enable the radiologist to determine the condition of the bones and joints at the time of the injury and the extent of the damage produced in them by the accident.

In some cases this evidence is of vital importance in assessing the responsibility. Too often the first radiographic record is one made several weeks after the injury, generally because the workman on being put back to his duties insists that he is now unable to do them owing to the injuries received. A radiographic examination at this time may show serious disease of the bones or joints and it may be impossible to estimate the condition of the area at the time of the accident. If we compare the radiographs of a healthy joint which has been dislocated and reduced by manipulation we see that those taken immediately after the injury show the bones and joint surfaces to be indistinguishable from normal, yet radiographs taken one, two, or more years after may show very extensive or progressive changes which are associated with limitation of joint movements. On the other hand, the first radiographs may show that pathological changes are already well established and in some cases they indicate that compensation is being sought after by the workman who alleges that a relatively trifling trauma has produced the lesion which must previously have given him definite signs and symptoms. The following cases illustrate the value of a comparative radiographic examination:

CASE I

A patient was sent to me for an x-ray examination of the spine, which the patient alleged had been injured three years previously. The documents accompanying the request for the examination contained the previous medical and lay reports, from which I learned that after the accident the patient had been taken to hospital, x-rayed, and on the radiographic evidence had been fixed into a plaster case for nine months; subsequently he had been supplied with block leather supports, but had not done any work.

The radiographs revealed evidence of an old lesion of the first and second lumbar vertebrae which had resulted in fusion of the two deformed bones. The appearances indicated a very severe trauma which must have necessitated a long period of immobilization, but the impression conveyed to me was that the lesion had been produced more than three years previously. On questioning the patient as to the nature of the injury I learnt that while loading a railway truck a large piece of coke had fallen on his back. Such a trauma I felt could not have produced the severe lesion indicated by the radiograph, for I found on weighing the largest pieces of coke that these did not weigh more than about one pound each. These facts were embodied in the report to the company concerned, with the suggestion that inquiries should be made into the past medical history of the patient and that the radiographs taken immediately after the injury should be submitted for examination. In spite of careful inquiries no evidence of a previous illness could be obtained. The man had worked well until the date of the alleged accident. Examination of the first radiographs in comparison with those I had taken enabled me to say that no definite change had occurred in the spine during the interval, and therefore the lesion was an old one and not caused by the accident. Therefore on the radiographic evidence it was decided to put the case before the courts. At the court the confidence of the company's counsel apparently caused the man's solicitor to more closely cross-question his client, as a result of which a small settlement was agreed upon before the court opened. After the decision had been made, in answer to a request from the workman's solicitor he was supplied with the nature of the radiographic evidence and he then informed us that as the result of the examination which he made of his client just before the court opened, he had learned that the man had had a severe injury to the spine some seven years previously for which he had received a large sum as compensation.

CASE II

In another case the patient was referred for a radiograph of the foot to ascertain if there was any evidence of fracture of the base of the metatarsals. The patient, who was claiming compensation, gave a history of a fall from a ladder a few months previously which had incapacitated him. The radiographs revealed the relatively rare condition of osteochondritis of the adult scaphoid, and the appearances were such that in the report I stated that the lesion was older than that suggested by the date of the alleged accident. Some months after, when the case came for revision, I discovered in my collection radiographs bearing a different name, taken at one of the hospitals two years previously, which duplicated all the appearances registered on the man's radiographs, even to the detail of the cancellous architecture. The patient at a subsequent examination denied any previous injury to the foot or radiographic examination, but he failed to pursue his quest for compensation.

In this connexion it should be remembered that certain developmental abnormalities may be detected on the radiographs taken immediately after an injury which may be misinterpreted as evidence of bone injury. Such conditions as: (a) epiphyses developing from multiple or irregular osseous nuclei—for example, the epiphysis at the base of the fifth metatarsal; (b) ununited epiphyses or ossicles—for example, the epiphyses of the vertebral bodies and processes, the os acetabuli, the accessory tarsal scaphoid, etc.; (c) developmental defects—for example, persistent notochordal remnants in the vertebral bodies, the central venous channels in the vertebral bodies, defects in the vertebrae at the junctions of the different regions, particularly the lumbo-sacral. Instances could be given in which the radiographic discovery of such lesions led to misinterpretation and claims for compensation.

Summary

1. It is essential for the study of the effects of trauma on bones and joints that a careful radiographic record be made of the part injured within twenty-four hours.

2. Such a record will register whether or not the bones or joints presented normal appearances at that time.

3. The effects of trauma can be determined only by serial radiographic examinations over a long period—in some cases several years.

4. Trauma appears to play a definite part in determining the focus of settlement of pathogenic organisms and malignant cells, but the evidence is not conclusive, and will not be until we have better radiographic records made immediately after the trauma has been applied.

5. Disabling arthritis may be produced by trauma in modern industry (that is, occupational strain) which is not catered for by the Workmen's Compensation Acts.

6. The importance of an immediate radiographic record after an injury to any bone or joint of a workman covered by the Workmen's Compensation Acts cannot be overestimated.

BIBLIOGRAPHY

- Alpers, B. J., Grant, F. C., and Yaskin, J. C.: *Ann. of Surg.*, 1933, xcvi, 10.
 Billington, R. W.: *Journ. Amer. Med. Assoc.*, 1924, lxxxiii, 683.
 Brailsford, James F.: *The Radiology of Bones and Joints*. J. and A. Churchill, London, 1935.
 Coley, W. B., and Higginbotham, N. L.: *Ann. of Surg.*, 1933, xcvi, 991.
 Middleton, D. S., and Bruce, J.: *Edinburgh Med. Journ.*, 1934, xli, Trans. Med.-Chir. Soc. Edin., May, p. 49.
 Pease, Chas. N.: *Amer. Journ. Dis. Child.*, 1935, xlix, 849.
 Sudeck, P.: *Arch. f. Klin. Chir.*, 1900, lxii, 147.

SUPRACONDYLAR FRACTURE OF THE ELBOW*

BY

NAUGHTON DUNN, M.A., M.B., Ch.B.,

LECTURER ON ORTHOPAEDIC SURGERY, UNIVERSITY OF BIRMINGHAM

Through the courtesy and help of several surgical colleagues I have had the opportunity of reviewing a series of cases of injury of the elbow attending the casualty department of a general hospital. We find:

1. That 50 per cent. of injuries to the elbow are supracondylar fractures.

2. That supracondylar fracture is rare after the age of 14. It occurs most commonly in children between the ages of 8 and 12.

Anatomy

Before considering the questions of cause, treatment, and possible complications and sequelae of this injury, it would be well to draw attention to certain points in the anatomy of the elbow-joint.

1. The lower end of the humerus is flattened, and curves forward in its relation to the shaft. Its lower articular surface is oblique, the outer condyle coming forward with the epiphysis, the inner condyle, lower than the outer, being directed slightly backwards.

2. The outer portion of the articular surface is domed to allow rotatory movement of the radial head, the inner portion is grooved and accurately adapted to the coronoid and olecranon processes of the ulna. The ulna is the direct continuation of the humerus, and owing to the obliquity of the articular surface of the latter, the forearm deviates outwards on extension. This establishes the normal carrying angle of the arm.

3. Three depressions are present on the lower end of the humerus: a deep fossa behind to accommodate the olecranon process of the ulna in extension, and two in front, one occupied by the coronoid process of the ulna, and a smaller one by the head of the radius in the position of flexion.

4. A knowledge of the ossification of the lower end of the humerus is important for the correct interpretation of radiographs. In relation to supracondylar fractures the important points to remember are that no centre of ossification is present in the lower end of the humerus until the second year, when that for the capitellum appears. No other centre

* Read in the Section of Orthopaedics at the Annual Meeting of the British Medical Association, Oxford, 1936.

JAMES F. BRAILSFORD: PATHOLOGICAL CHANGES IN BONES AND JOINTS INDUCED
BY INJURY

FIG. 1.—Radiograph of the hands of a man who had worked with a compressed air drill for ten years. Note the rounded areas of cancellous destruction in the heads of the third metacarpal of the left hand, the heads of the second and third metacarpals of the right hand, and the carpal bones of the right hand. Similar changes were seen in the capitellum and the head of the radius.

FIG. 2.—Radiograph of the wrists and hands of a foundry-worker, showing very severe arthritic changes in the first metacarpal, carpal, and inferior radio-ulnar joints on both sides.

JAMES F. BRAILSFORD: PATHOLOGICAL CHANGES IN BONES AND JOINTS INDUCED
BY INJURY

FIG. 3.—Radiograph of the left hip-joint of a man aged 27 (22.3.27), showing a dislocation due to a fall.

FIG. 4.—Radiograph of the same hip-joint as Fig. 2 (17.8.27), showing rarefaction of the upper part of the femoral head and new bone formation in the soft tissue around.

FIG. 5.—Radiograph of the same hip-joint as Figs. 2 and 3 (14.12.31), showing consolidation of new bone causing marked restriction of movement. The superior lip of the acetabulum is sclerosed.

FIG. 6.—Serial radiographs of a proximal phalanx before, and two, four, six, and ten months after surgical treatment of one of the metacarpals. Note the changes in the cancellous structure induced by immobilization of the hand.

FIG. 7.—Radiographs of the elbow-joints of a foundry-worker. The elbow-joints were continually subjected to minor traumata and gradually marked limitation of movement developed. Such disablement when so produced does not entitle the man to compensation under the Workmen's Compensation Act.