this position should be maintained uninterruptedly for a period of three to four weeks. In the case in which no complications have arisen, the collar and cuff may then be loosened so as to allow 30 degrees of active movement of the joint, which active movement should be encouraged.

If it is found that the patient can voluntarily flex the joint to the position of previous fixation after three days, gradual relaxation is allowed, and usually at the end of four or six weeks movement from below a right angle to flexion is strongly controlled, and free use of the arm without any fixation may be permitted. It may be two or six months before extension is complete, but if reduction is complete and treatment is carried out on these lines there need be no anxiety as to the final result.

Forward displacement of the lower fragment may occur as a result, usually, of a fall on the flexed elbow. In these cases reduction will be established by traction and direct backward pressure transmitted to the lower fragment through the flexed forearm. In all recent cases of supracondylar fracture manipulative reduction should be possible. It must be remembered that because of the rapid formation of new bone late operation to restore alignment is difficult, and the results are seldom satisfactory.

Complications

The serious complications of this fracture which may occur are: (1) nerve injuries; (2) Volkmann's ischaemic contracture; (3) myositis ossificans.

(1) *Nerve Injuries*.—In supracondylar fractures the median is the nerve most commonly injured. The sharp lower anterior margin of the shaft of the humerus may cut it during the original displacement, or during reduction if the arm is hyperextended. In this series I have not seen a case of immediate division of the nerve, but three cases on which I have operated have shown the nerve frayed, adherent to, and stretched over, the sharp margin of the upper fragment. In some cases a paralysis of the ulnar or musculo-spiral nerve has appeared. In all of these natural recovery has taken place.

(2) *Volkmann's Ischaemic Contracture*.—The possibility of this, fortunately rare, complication of injuries to the elbow must always be remembered. This is less likely to occur when early reduction of the fragments is established, but the case must be under constant supervision for the first forty-eight hours in order to be sure that there is no interference with peripheral circulation. Undue congestion leads to rapid and irreparable destruction of muscle, so that its early recognition is of paramount importance. The early symptoms begin in a few hours; the hand is painful, the fingers are discoloured and numb, and voluntary power is diminished. This is the prelude to subsequent contractures, the treatment of which is not within the scope of this paper. Efforts to reduce the congestion should be directed to relaxation of flexion at the elbow, and, if necessary, reduction of tension in the antecubital fossa by incision.

(3) *Myositis Ossificans*.—This complication is to be suspected if it is found, after relaxation of the position of flexion at the end of the third week, that movement is not naturally restored. An x-ray may then reveal a shadow in front of the elbow. Passive movements, always contraindicated in injuries to the elbow, will increase the formation of new bone. The essential treatment is to continue rest of the joint in the position of flexion until either the bony deposit has reabsorbed or has completed its ossification. Early operation is wrong, as recurrence will surely follow. If operation is necessary it should be undertaken only when consolidation is complete and if the new bone is obstructing movement.

I am indebted to Dr. J. F. Brailsford for the line drawings which accompany this communication.

FRACTURES OF THE HEAD OF THE RADIUS AND CAPITELLUM INCLUDING EXTERNAL CONDYLAR FRACTURES OF CHILDHOOD*

BY

ST. J. D. BUXTON, F.R.C.S.

SENIOR ORTHOPAEDIC SURGEON, KING'S COLLEGE HOSPITAL

My task in opening this discussion on fractures on the outer side of the elbow-joint is difficult, as brevity is essential, and as I am particularly interested in these lesions. Controversy is more likely to be centred on the treatment than on the method of production of the injuries; hence it is my intention to point out the relative frequency of these fractures and the ages at which they are prone to occur and then to proceed to the methods of treatment suitable for each type of fracture. The end-results can only be studied by examination of the limbs and, as this is not possible at the meeting, I refrain from the useless classification into good, moderate, and poor results, but I am particularly anxious to invite discussion on the difficulties. In view of the classification which has been employed in our clinic for some years, it is necessary to divert somewhat from the title rather than substance of this discussion.

During the three-year period (1933, 1934, 1935) of about 3,000 ambulatory fractures 206 were fractures of the outer side of the elbow-joint. In private practice, and at country hospitals, one sees only the severe cases and often those only requiring operative treatment, so all statistics are purposely limited to one clinic.

The table shows the classification and some points of interest with regard to the age of those injured.

Analysis of 206 Cases

		Total Cases	Under Age 14	Youngest Age	Oldest Age	Average Age
Radius	Head ...	124	6	10	68	22½
	Neck ...	37	17	7 (23)	65	14
Humerus	Capitellum*	38	19	6	27	15
	Ext. Epicondyle	7	2	13	40	22
Total ...		206	44	—	—	—

* Usually flake of diaphysis and capitellar epiphysis.

Fractures of Head of Radius

These occur most frequently between the ages of 25 and 35, and are of three types:

- (1) A vertical fissure, descriptively termed a "chisel fracture";
- (2) A fracture with one loose fragment;
- (3) Comminution of the head, including the mushroom fracture.

(1) and (2). When there is little or no displacement the treatment consists in placing the forearm in full flexion and supination. If the limb cannot be brought into this position when the patient is first seen, the elbow is flexed as far as is comfortable by a collar and cuff or special sling, and daily adjustments will soon enable the desired position to be obtained. This position is retained for four weeks, then for one week the elbow rests at a right angle in a sling. After that, flexion and extension exercises are commenced. Although most patients have good function in two months, full function may not be obtained for four months.

(2) When one loose fragment is displaced, this should be removed by operation, preferably within one week of the accident, through an incision on the outer side of the joint.

* Read in opening a discussion in the Section of Orthopaedics at the Annual Meeting of the British Medical Association, Oxford, 1936.

In some of the cases the head is tilted owing to a further fissured fracture. In most cases it is advisable to limit the operation to removal of the loose fragment.

(3) In comminuted fractures the total circumference of the head is increased, by separation of the fragments. The operation of removal of the head and part of neck of the radius is necessary. My experience is that the best results are obtained by immobilization of the elbow for six weeks and then to operate.

If the arm is left still, ossification will not take place in the capsule during the waiting period, nor after the operation. As movement has not been allowed, there is no reason to suppose osteoarthritis will be more frequent in consequence of the delay, although it is probably inevitable in a few cases following any operation involving opening a joint. After this operation a full range of movement has been obtained, but usually extension is permanently limited by twenty degrees and other movements are full.

Fractures of the Neck of the Radius

These are usually transverse. A wedge-shaped piece of bone is often torn off the shaft from one side and is adherent to the head. Before the epiphysis has joined, the fracture is just distal to the cartilaginous plate, but the diaphyseal portion broken off is usually less than half the circumference of the neck.

(1) As a rule there is little displacement, or this may be a rotatory one. With little rotation or displacement, the treatment is similar to that of a fissured fracture of the head.

(2) In one-third of these fractures there was displacement, that is—a tilt to a considerable degree or displacement of the head right off the shaft, for it may be alongside the shaft, or be displaced to the inner side of the forearm. Operative treatment is indicated. Should the head be displaced as a whole, but also be broken into a number of fragments, it ought to be removed. In other cases reposition is possible and definitely indicated in patients under forty. In children it is very desirable, as removal of the head leads to cubitus valgus of extreme degree. This may be complicated by limited wrist-joint movement, radial deviation of the hand and prominence of the head of the ulna. The operation should be carried out within a week of the injury. If the head will stay in place after reposition, no fixation is necessary. If it will not remain in position it is often possible to fix it by a stitch of catgut through a tag of periosteum adherent to the neck or by passing a tiny needle through two drill holes distal to articular cartilage. In children the head may not stay in place without further fixation. I have reported a method of fixation by driving a bone peg through the articular surface of the head and up the shaft, so that at the end of the operation the peg was extra-articular and there was a minimum of damage to articular cartilage. Although the operation is difficult at times, I have continued to employ it with satisfactory results. I have seen on two occasions manipulative reposition of the head of the radius in children, when it had been displaced right off the shaft.

Fractures on Outer Side of Lower End of Humerus

In discussing fractures on the outer side of the lower end of the humerus, I am not including any which extend to the inner side, nor the comminuted fractures seen in adults associated with dislocation and fracture of the olecranon. Intra-articular fractures of the capitellum only appear to be very rare.

(1) In adults the line of fracture may be oblique from above the external epicondyle into the joint medial to the capitellum. More frequently the capitellum, part of the trochlea, and a piece of non-articular bone above and behind are fractured from the shaft. If there is no displacement, the elbow is put at rest at a right angle in a posterior

plaster trough and x-rayed in the plaster to see that the position is still satisfactory. When there is displacement, open operation is indicated. Reposition of the fragments is not difficult. In most cases fixation is not required, but if the fragments will not remain in place, a stitch passed through a drill hole made in the diaphysis will often hold the fragments, or a small bone peg an eighth of an inch across may be driven transversely through the fragment to hold it to the trochlea.

(2) In children the fractured piece consists of the capitellar epiphysis (that is, the capitellum and half the trochlea) and a flake of the diaphysis, usually off the back of the lower end of the shaft. When there is no displacement the fracture is treated after the manner of a supracondylar fracture with no displacement. When there is displacement, an open reposition is employed or non-union with cubitus valgus will occur. Formerly one used some form of peg or nail fixation, but this is seldom necessary. A catgut stitch through the diaphyseal flake to periosteum or muscle on the shaft will hold it in place. Some fixation is necessary or else the fragment will displace laterally. Normal elbows are obtained after efficient reposition. The cases of fracture of this area in childhood and young adults which have been neglected not only lead to a weak elbow, but sometimes ulnar nerve neuritis or paralysis.

(3) Fractures of the external epicondyle are isolated and uncommon injuries. Relaxation of the muscles by flexing the elbow until union has occurred is sufficient treatment.

Of the last 100 consecutive cases of fracture on the outer side of the elbow-joint, open operations have been performed on:

Head of Radius (8 cases):

1. For single loose fragments, 3 cases (fragment removed).
2. For comminuted fracture, 5 cases (head removed in 3 cases, pieces in 2).

Neck of Radius (6 cases):

1. Removal of head, 1 case.
2. Reposition of head with fixation: (a) by catgut or none, 3 cases; (b) by bone peg, 2 cases.

Humerus (5 cases):

1. Capitellum, 3 cases.
2. External condyle in adult, 1 case.
3. One operation for ulnar nerve paralysis.

FRACTURES OF THE INTERNAL EPICONDYLE OF THE HUMERUS*

BY

S. L. HIGGS, M.B., B.Ch.CANTAB., F.R.C.S.

SURGEON, ROYAL NATIONAL ORTHOPAEDIC HOSPITAL; ASSISTANT SURGEON, ORTHOPAEDIC DEPARTMENT, ST. BARTHOLOMEW'S HOSPITAL

(With Special Plate)

In 100 consecutive cases of recent fractures of the elbow-joint treated at St. Bartholomew's Hospital sixteen involved the internal epicondyle of the humerus, and this represents about the average recorded in other statistics. The average age was 12 years. It is therefore not a rare injury. From the number of patients who seek advice on account of late complications it is clear that recognition and treatment of the original condition are often at fault.

General Considerations

The anatomical peculiarities of the internal epicondyle make it prone to injury in childhood. It has its own centre of ossification, which appears at the fifth year, and does not fuse until about the age of 16. During this period separation of the epicondyle may occur through the relatively weak epiphyseal cartilage. To the epicondyle are attached the internal lateral ligament and the common origin of the flexor group of forearm muscles. Thus a

* Read in opening a discussion in the Section of Orthopaedics at the Annual Meeting of the British Medical Association, Oxford, 1936.