Conclusion

And now, having come to the end of my very summary account of these researches, there are a few points which appear of sufficient importance to deserve emphasis.

Experimental work to elucidate the matter of human infection has usually been done on the dog. Here in the diseases brought to your notice the affected animal is the sheep. The study of these conditions is of value because (1) they are natural infections, (2) the numbers concerned are large, (3) experiments can be controlled in many ways by normal sheep available at every abattoir, and (4) the experimental animal for a crucial test of the reproduction of the disease or its prevention is the same animal as is naturally attacked.

A point of great interest has been to discover the conditions allowing of infection. Williams wisely pointed out in his paper that it requires very particular conditions for the establishment of *B. welchii* infection in living human tissues, and our survey has provided ample justification for that remark, with a more general application.

The avoidance of any complication from post-mortem invading organisms has undoubtedly facilitated these researches. It is easy in the sheep, it is very difficult in man, and the confusion which may follow must be reckoned with and avoided if possible. The precise identification of an organism and the determination of a particular type of strain with its own specific toxin is an essential feature of any successful investigation. As with the pneumococcus so with *B. welchii*—typing is essential if serum therapy or prophylaxis is to be of use.

Lastly, the presence—potential or actual—of latent spores must be considered. Let us fully understand what are the possibilities of infection, and the conditions which make it actual, and we shall be well on the way to the discovery of means of prevention.

BIBLIOGRAPHY

- Albiston: *Aust. Journ. Exper. Biol. and Med. Sci.*, 1927, iv, 113.
 Bennetts: Commonwealth Council for Scientific and Industrial Research, *Bulletin*, 1932, 57.
 Cleland: New South Wales Microbiological Laboratory, fourth report, 1913.
 Dalling: *Journ. Comp. Path. and Ther.*, 1926, xxxix, 148.
 Dodd: *Ibid.*, 1921, xxxi, 1.
 Gilruth: *Veterinary Journal*, 1910, lxvi, 254.
 Holt, R. L.: *Lancet*, 1934, i, 724.
 Oxer: Commonwealth Council for Scientific and Industrial Research, 1932, Pamphlet 35.
 Turner: Commonwealth Council for Scientific and Industrial Research, *Bulletin*, 1930, 46.
 Turner and Davesne: *Ann. de l'Inst. Pasteur*, 1927, xli, 1078.
 Williams, B. W.: *Brit. Journ. Surg.*, 1926, xiv, 54.
 Wilsdon: University of Cambridge Institute for Animal Pathology, second and third reports, 1931 and 1933.

The Minister of Health has addressed to the London County Council a letter arising out of his consideration of the report made by his officers following a survey of the L.C.C. public health services—a survey which is referred to at length in the annual report of the Ministry of Health published in August last. The Minister states that he has observed with appreciation the great work of organization and development of the medical and institutional services for the sick which has been carried out by the Council since the Act of 1929 came into operation. He draws attention to three points which afford him special satisfaction: the improvements in structure, equipment, and staffing of the hospitals which the Council inherited from the guardians; the separation from the Poor Law of services which could lawfully be provided under other powers; and the close interrelationship of the special services, such as those for tuberculosis and venereal diseases, for which the Council was responsible before the Act of 1929, the general hospital service, and the special hospital service. The Minister has also observed a number of interesting points of contact between the Council's services and those of the voluntary hospitals of London, especially the facilities afforded at Council hospitals for training medical students in obstetrics. He congratulates the Council both on its enlightened policy and on its efficient organization.

CHRONIC RHEUMATISM*

BY

R. FORTESCUE FOX, M.D., F.R.C.P.

PRESIDENT OF THE INTERNATIONAL LEAGUE AGAINST RHEUMATISM

After a long period of stalemate a concerted effort against rheumatism is making good progress. Like the campaign against tropical diseases, it has widespread social, economic, and industrial purposes to serve in addition to its primary medical aim of prevention and healing. There are moments when those who are engaged in a long effort against disease should take stock of their position, of success and failure, of the ground won, of immediate objectives and those which are more remote, and of the best measures by which those objectives may be attained. It is ten years since the International League against Rheumatism had its small beginning. Two national organizations now represent the Anglo-Saxon countries: the Committee on Chronic Rheumatism of the Royal College of Physicians of London, which has just issued its first report, and the American Association for the Study and Control of Rheumatic Diseases. At the latter's recent annual meeting in Philadelphia much suggestive new matter was presented by good workers in rheumatology in the United States.

Causation

At such a moment it is well to consider, even at the risk of dogmatism, what new facts have been clearly ascertained, and, in default of facts, what guiding hypotheses or stepping-stones to facts may have emerged. It is the more desirable to arrange new facts and postulates in a country where rheumatism is so prevalent, for it is only as guiding principles emerge that it is possible to grapple with the great problem of rheumatism in its various aspects.

The reference of the College of Physicians committee was the chronic rheumatic diseases, and especially arthritis. It was agreed that chronic rheumatism forms a series of definite clinical pictures. Whatever anatomical or aetiological links may subsist between chronic rheumatism and "acute rheumatism," these two conditions differ widely in their evolution and clinical course, and present in practice two entirely separate problems. Incidentally, the acute disease, to which the name of cardiac fever might well be applied, appears to be diminishing in our country, at all events in the form in which we knew it forty years ago; this can hardly be said of the chronic and disabling maladies now under review.

There is good histological evidence that the connective tissues are the seat of chronic rheumatism. What are regarded as the typical abnormalities of the rheumatic state are found in these tissues of the mesenchyma. It is also generally accepted that the causation of rheumatism, like that of other chronic diseases, is complex. Some years ago the International League formulated four possible causal factors, internal or external to the body. It is impossible to state them in order of importance, since sometimes one, sometimes another, seems to be dominant. They are: (1) constitutional anomaly; (2) abnormalities in the circulation of the skin; (3) external factors, such as cold, damp, trauma, and occupation; and (4) focal or other infection.

A diagnosis of causes may well be made in the first instance. This will discover, so far as is possible, what in each case is the dominating, and what the contributory or accessory, causal factors. Only then can a scheme of treatment be framed which will counteract the combined forces of causation. The dominating factor may well be

* A British Medical Association Lecture (abridged) delivered to the South Suffolk Division at Ipswich, June 21st, 1935.

an infective invasion, or an inadequate resistance to infection; or an abnormality of the skin and circulation, a mere deficiency of body heat; or some continuous or recurring injurious external influence; or a constitutional abnormality preventing recovery from disease. Any one of the causes of a chronic disease may be dominant, in the sense that the patient cannot be cured without removing or counteracting it. Nearly all countries have long had spas or institutions for the treatment of chronic rheumatism, and the League has endeavoured to pool their varied experience. The first and entirely rational objective in the treatment of rheumatism is the alleviation of pain. The injurious effects of pain are evidenced by the fact that its alleviation by physiological methods is often directly curative. Unquestionable experience shows that systematic physical treatment at an early stage of the disease directed to the skin and circulation in many cases prevents more serious developments. It is clear that every stage in the evolution of rheumatism requires its own systematic care, and that much can be done to combat rheumatism in all its stages. Even in very advanced cases, when the attempt might seem to be of little use, periodical treatment may give admirable results in preventing further damage and in maintaining functional capacity. The importance of this fact from the occupational point of view needs no comment.

Desiderata

Certain desiderata become obvious where numbers of cases are under treatment together. It is necessary, for instance, to frame comparable case records on an agreed basis, so that the practice and results of treatment at the great spa hospitals and elsewhere in this and other countries may be compared, and the present ambiguities of nomenclature minimized, if not prevented. Other problems relate to the measurement of functional capacity, and therefore the degree of industrial invalidity. Some countries, like Sweden, have made special progress in this direction. In the case of insured patients precise measurements have been made at the beginning of a course of treatment, at its termination, and again at intervals of three, six, and twelve months. It is known that the effects of physical treatment cannot be accurately stated until a period of months has elapsed. Comparisons and measurements of this kind are obviously necessary, both for medical and for industrial purposes.

Again, the field of biophysical research is as yet but little cultivated. This may be held to include the study of functional disturbances in rheumatic subjects, such as abnormalities of body heat—low temperatures (deep or superficial), abnormal reactions to external cold—as well as the equally important study of the effect of physical impressions, first on the skin and circulation, then on nerve centres, metabolism, and nutrition. It is clear that the effects produced by heat and cold in rheumatic persons are not the same as those produced in health. The reactions to physical impressions are very variable, even in the same individual. It is vain to attempt to define these reactions in rheumatism by experimental studies of healthy subjects and animals: results so obtained have often been misleading. It is, however, true that without precise biophysical data derived from the study of rheumatic patients the best attempts at physical treatment will be a groping in the dark.

Another fruitful field of investigation is the causation of rheumatic disease (in its different forms) as affected by locality and climate, by occupational strain or bad posture, by work in the open air, factories, or workshops, as well as by housing, age, and sex. Here comprehensive statistical records on an agreed basis are much needed, especially in regions and occupations where rheumatic disease is prevalent.

Centres for Treatment

The incidence of rheumatism is so great in this country, and its nature and causation still so complex and obscure, that there is good reason for setting up *ad hoc* centres where every possible causal factor can be investigated, and where large numbers of cases can be carefully studied at every stage of the disease. In several countries the medical profession has been feeling its way towards the ideal centre for the investigation and treatment of rheumatism. In England the first demonstration clinic for rheumatism to be established in a city was opened in London by the British Red Cross Society in 1930. The spas are also experimenting in this direction. The long-established care of rheumatism at Buxton has been supplemented by the newly opened residential clinic. At Droitwich the Royal Brine Baths Clinic has been opened in 1931. In Sweden rheumatism departments have been organized in several of the chief hospitals, also in Czechoslovakia and Hungary. In Germany, Holland, and Belgium there are ambulatory clinics for private and insured patients, and in the Soviet Republic the numerous spas and treatment institutes for rheumatism in the towns form an important part of the public health service.

Types of Arthritis

There are three main forms of arthritis, which present different clinical pictures, different prognoses, and different therapeutic indications; they are also related to different periods of life. They are: (1) the rheumatoid disease of early adult life, or, as the Americans call it, atrophic arthritis; (2) the osteo-arthritis of age, which the Americans call hypertrophic arthritis; and (3) the arthritis of middle life, including what has been called in several countries climacteric arthritis. This third type has apparently a more complex causation than the others, and certainly shows very variable clinical characters. It is not surprising that metabolic, nervous, circulatory, and endocrine changes in middle life should modify its clinical features, or, as many now believe, bring to the surface latent hereditary characters.

Rheumatoid Arthritis

This type (seven out of ten cases are in women) presents many of the features of a specific disease with profound constitutional disturbance. In the worst cases the arthritis becomes generalized with characteristic deformities, but occasionally, after active tonic treatment at an early stage, the disease is arrested, at all events for some years, leaving a few spindle-shaped enlargements in the first row of the interphalangeal joints. The terminal joints often undergo primary atrophy and ankylosis in hyperextension.

The appearance of a rheumatoid patient is suggestive of infection; the rate of sedimentation of blood cells is increased, as it is in all inflammatory states. Failing specific treatment, it is of great importance to restore the circulation and to increase the general resistance of the body. Cod-liver oil, iron, and arsenic are of undoubted benefit, also light and air baths, with frictions and generous feeding—measures to increase metabolism and body heat. Unfavourable occupational or housing conditions should of course be rectified; a change of climate should be considered. For girls with cold skin, subnormal temperature, anaemia, and amenorrhoea hyperthermal baths have been used, the temperature of the bath rising day by day from 105° to 115° or over for one to three minutes. If there are signs of focal infection in the pelvic organs diathermy or zinc ionization should be prescribed.

Degenerative Arthritis or Osteo-arthritis

The general health of patients with this type is but little affected: signs of infection are uncommon, and the sedimentation rate is usually normal. The traumatic or

postural factor in causation suggests mechanical measures of relief to the joint. In many painful cases deep x rays have been used in recent years with advantage. Occasional blistering or Paquelin cauterization is also beneficial where one or a few joints only are affected. Applications of heat relieve pain and sometimes seem to stabilize the disease, probably through modification of the circulation and nutrition. Thus good results often follow from hyperthermal packs of estuary mud or fango. In Russia the use of muds (peloids), followed by movement as the pain subsides, now takes a leading place in treatment.

The muscular spasm so generally associated with osteoarthritis, especially in the hip, may be counteracted by prolonged pool baths. The muscles are gradually relaxed, and both active and passive movements can often be obtained without pain. Adductor contraction, in particular, can often be overcome in the pool by movements and manipulation under water. Applications of luminous heat to the hip, prolonged for one hour, are praised by some rheumatologists. At a later stage, when movement can be effected without pain, mechanotherapy, especially cycle exercises, may be used. For robust and stout patients thyroid extract may be given, and carbohydrates excluded from the diet. An occasional fast is often beneficial.

"*Senile rheumatism*" (the "rheumatics" of the rustic) is a benign osteo-arthritis common in old people who have worked hard or been much exposed to the weather. It begins in the phalangeal joints, and leads to slow and often painless bony enlargements. Since in these cases there is usually a feeble and failing circulation frequent warm baths and protective clothing, such as a chamois jacket and woollen mittens, should be used to increase and conserve the body heat. Sulphur and guaiacum are still popular remedies.

Arthritis of Middle Life

The arthritis of middle life is one of the commonest forms of breakdown at this period, and in certain families either arthritis or "gouty manifestations" are almost usual at the climacteric in women. A toxic or metabolic factor can generally be traced in all the chronic illnesses of middle life, due, it may be, to the formation of abnormal substances or to simple failures of excretion. It may be remembered that gout also is modified by age and sex. In middle life pre-arthritis symptoms are usually met with, such as excess of uric acid, dyspepsia, chilliness, disturbances of circulation, heat or cold in the extremities, disturbances of the mucous surfaces or skin (catarrh, eczema). The average age of the arthritic onset is 48. It may come on suddenly, usually first in the knees, and the evolution is centrifugal from the large joints to the small, that of rheumatoid disease being centripetal. It is of a mixed character, and peri-arthritis and fibrositis are common. In advanced stages the characteristic changes of osteo-arthritis supervene. Interruptions in the development of the disease are not uncommon, followed in favourable cases by comparative quiescence after the climacteric is passed, with permanent changes in one or a few joints. It is thus a critical malady, and less generalized than the arthritis of young subjects, with a more favourable prognosis. As in all forms of arthritis, depressing influences such as overwork, anxiety, parturition, and influenza may be dominant causal factors. An endocrine factor, and exposure, occupation, and trauma must be considered. Infective causation will be noted immediately.

The treatment of climacteric arthritis will naturally be directed not only to the affected joints but to the underlying state. The best remedies will be modifying or "alterant" in character, and affect the body as a whole, such as the "cure" at a spa or clinic prolonged for three weeks (seldom longer) and repeated for two or more years. In a large number of cases the periodical course

of waters and baths fulfils this indication better than any other form of treatment. The change of scene, habits, diet, and regulated rest and exercise do much, but an eliminative water in middle life, combined with external applications, is an effective prescription. Pure sulphuretted waters also have a favourable influence in middle life. Appropriate baths or "peloid" packs continued for a limited "cure" period appear to produce a cumulative and semi-permanent effect on the skin and peripheral circulation and on the absorption of deposits. Recent reports from Russia show that nearly 100,000 rheumatic workers were referred to the mud spas in 1933 and 44,000 to the sulphur spas. In the home, active treatment of the skin, with a course of alternating sulphur and iodine preparations, may be helpful, especially in the autumn or spring: in congestive subjects, mild mercurials and alkalis. Fasting and low calorie diet, rest in bed, and diminishing the strain upon arthritic knees when body weight has increased are especially applicable to stout middle-aged women, overworked housewives, and charwomen.

The Infective Factor

Very diverse views are still expressed as to the part played by focal infection in arthritis. An increasing number of authorities accept the hypothesis that rheumatoid disease (the progressive deforming polyarthritis of young subjects) is a specific one, and that poor nutrition, exposure to cold and damp, anxiety, and previous illnesses are predisposing causes to a bacterial or toxic invasion. On the other hand, they hold that osteoarthritis is the result of failing nutrition in the joints in later life, long wear and tear, strains, traumatism, and hereditary tendency, without invoking an infective factor. That is not to say that in exceptional cases an accidental infection may not light up an osteo-arthritis in damaged tissues. A recent example in a middle-aged woman showed a subacute osteo-arthritis in one shoulder-joint, with pain, coarse crepitation, and great limitation of movement, which rapidly subsided when a septic tonsil of the same side was removed by zinc electrolysis. An active focus may be present in a proportion of middle-aged patients, but in a large number of cases the existence of such a focus cannot be demonstrated. Unless there is clear evidence of septic absorption from teeth or tonsils, as shown by constitutional disturbance and a high sedimentation rate, their removal can hardly be justified. It was commonly stated not long ago that septic foci ought always to be removed as a preliminary to treatment at a health resort. Against this practice may be cited the frequent disappointments attending surgical treatment, and the interesting fact that cases are commonly met with at the spas in which a suitable course of waters and baths has given good results, notwithstanding the presence of septic foci. If an active invasion is going on it is still possible for a "cure" to do essential service by increasing the resistance of the tissues. Less emphasis is now laid on the universality of infection as a factor in arthritis than was the case a few years ago, both in America and in England, and more importance is attached to restoring lost immunities to the ubiquitous micro-organism. It has been shown, for example, that infective foci may be as common in healthy persons as in the subject of arthritis.

This change of viewpoint is exemplified by two recent statements. "Few greater injustices," say Pemberton and Osgood,¹ "can be done to these unfortunate sufferers than to permit their complicated problem to be envisaged by the surgeon or others, be they never so skilful, who hang the whole syndrome upon a single aetiological hook." The same writers relate that at a certain State sanatorium in the United States the children led an open-air life. Large numbers of cases for tonsillectomy were received

when new buildings were under construction. The operation was delayed for several months while the throat department was being finished. When the surgical quarters became available only four out of forty waiting children required operation. Sutcliffe has observed similar improvement at the Sea Bathing Infirmary at Margate in the health of children waiting for operations which subsequently proved needless. Danischewsky, the honorary secretary of the Russian Committee on Rheumatism, also writes: "Infection [with us] is now considered as a function of three variables, the micro-organism, the macro-organism, and the environment." It is well known that many French authorities are inclined to dissociate from true rheumatism all acute infective conditions.

The use of vaccines must follow the accepted doctrine on infection. Treatment by preparations of gold is still in an experimental stage. Strasser of Vienna and others advocate a "simultaneous therapy" of arthritis by a combination of remedies. This practice is opposed by other authorities, because of the difficulty of interpreting the reactions which are produced. They hold that "in the use of physical remedies, especially baths, the daily reaction is a necessary guide to treatment. The simultaneous use of vaccines or of gold inevitably blurs the picture and deprives the practitioner of essential guidance in conducting the cure.

Conclusion

The knowledge of chronic rheumatism has been clarified and extended in recent years, and new suppositions are leading to new knowledge. More reliance is now placed on physical medicine, both for the prevention and for the relief of chronic rheumatism. Many observers have been impressed by the disturbances in the peripheral circulation, the systemic vaso-constriction and curtailment of blood flow in the arterioles and capillary bed, by the failure of the normal reaction to cold, especially when muscular activity is in abeyance, and by the consequent slow invasion of chill. What is the remedy of choice to deal with the invasion of cold on the surface of the body? Interesting new evidence of the effect of external heat in removing stiffness and pain and other effects of exposure to cold, fatigue, and occupational trauma is forthcoming from pithead baths.

The same processes seem to be at work in articular and in non-articular forms of chronic rheumatism. Consequently, the methods of treatment applicable to arthritis also apply to sciatica, lumbago, and fibrositis. To this entire category of chronic painful diseases the distinctive generic name of rheumatism has always been given, and this older and limited connotation of the name has been confirmed by recent knowledge.

The establishment in England of rheumatism centres for consultation and treatment³ for large numbers of cases, served by medical officers who devote their whole time and thought to the study of rheumatology and physical treatment, appears to offer advantages which it would be difficult, if not impossible, to obtain in any other way. The early physiological abnormalities of rheumatism could be properly investigated; light would be thrown on many obscurities of causation, and well-ordered schemes of preventive and curative treatment could be devised. They would go far to establish the methods of physical medicine on a scientific foundation, and would help to launch a medical-social campaign against rheumatism in England which might, in time, be carried to a point that few have dreamed of.

REFERENCES

- ¹ *Medical and Orthopaedic Management of Chronic Arthritis*, 1934.
- ² *La Balnéothérapie des Maladies Rhumatismales et ses Indications*, Moscow, 1934.
- ³ Fox, R. Fortescue, and van Breemen, J.: *Chronic Rheumatism: Causation and Treatment*, 1935.

MALIGNANT ENDOCARDITIS DUE TO BRUCELLA ABORTUS

BY

J. K. RENNIE, M.D., B.Sc., F.R.F.P.S.GLAS.

AND

C. J. YOUNG, M.B., D.P.H., D.T.M.

(From the Bradford Royal Infirmary)

The occurrence of death in cases of undulant fever is unusual, and a fatal termination as a result of endocarditis would appear to be very rare. This is true of infection both by *Brucella melitensis* and by *Brucella abortus*. The earliest record we can find of the former is that of Hughes,¹ who in 1897 described four cases of undulant fever in which the heart was affected and all of which terminated fatally. Three of the patients suffered from definite vegetative endocarditis, the fourth from pericarditis. Recently Casanova and d'Ignazio² have described a case of vegetative endocarditis of the aortic valve due to *B. melitensis*.

In infection by *B. abortus* cardiac involvement is equally uncommon. Kristensen³ has recorded one case in which the lighting-up of an old endocarditis caused death, the terminal event being probably an embolus of the mesenteric artery; but no post-mortem examination was held, so that the precise condition was not established. Scott and Saphir⁴ have recorded the case of a patient who had previously suffered from rheumatic fever with endocarditis, and who died as the result of infection of *B. abortus*. Post mortem, recent vegetations were found superimposed upon the old valve lesions, so that death was almost certainly due to this cause, but no histological examination of the vegetations was made. Profound myocardial damage occurred in two cases described by Attinger.⁵ In neither was any endocardial involvement recorded. Thus beyond the case of Scott and Saphir we have been unable to find a record of the condition established by post-mortem examination.

Case Report

The following case of a patient who died of endocarditis due to *B. abortus* is therefore worthy of record.

In March, 1935, the patient, a wool-packer aged 47, suffered from "influenza" followed by persistent diarrhoea, which necessitated his admission to hospital on April 27th, 1935. He was much emaciated. The abdomen was slightly distended, but neither liver nor spleen was palpable. The heart was not enlarged and no murmurs were detected. There was tenderness over both kidneys posteriorly, and albuminuria and haematuria were present. The Wassermann test was negative. The patient gave a history of occasional attacks of tonsillitis until tonsillectomy was performed four years previously. His work never brought him in contact with live animals, and he drank milk very sparingly. No other member of the family complained of illness.

The diarrhoea quickly cleared up, but pain persisted in the upper abdomen. There was pyrexia ranging up to 100° F. with a relatively higher pulse rate. The blood agglutinated *B. abortus* to a titre of 1 in 6,400, and blood culture, which was negative with routine media, showed *B. abortus* to be present in large numbers when suitable methods were employed.

After a slight improvement the patient's condition deteriorated steadily. A systolic murmur developed at the apex. The liver became enlarged, but the spleen could not be palpated. The fingers showed definite clubbing. There was progressive anaemia, and casts were found in the urine. On one occasion only a stool containing bright red blood was passed. The temperature was never undulating, and the leucocyte changes usual in undulant fever were absent. Later a presystolic murmur developed and the diarrhoea recurred. Oedema appeared, the pulse gradually failed, and the patient died on July 10th.