that we do not "expect" bruises in any case of violence. If they are present, and agree in age with the date of the violence, well and good; but even then the extent of the bruising may not have the slightest relation to the severity of violence. Extreme cases in this respect are instanced in the thrombopenics and the haemophiliacs.

Plate III: All the figures are multiplied 100 diameters. Photomicrographs by C. F. Hill and W. E. Cooke.

REFERENCES

- ¹ Osler, Sir W., and McCrae, T.: *The Principles and Practice of Medicine*, London, 1920, p. 79.
- ² Allbutt, C., and Rolleston, H. D.: *A System of Medicine*, London, 1909, v, 194.
- ³ Cooke, W. E.: *British Medical Journal*, 1934, ii, 624.
- ⁴ Giovanni, C.: *Annali Omodei*, 1861, clxxi, 14.
- ⁵ Moore, G.: *Lancet*, 1842, i, 205.
- ⁶ Smith, F. J.: *Taylor's Medical Jurisprudence*, 1920, p. 411.
- ⁷ Le Conte, R. G.: *Ann. of Surg.*, 1908, xlvii, 383.
- ⁸ Wideroe, S.: *Norsk Mag. f. Laegevid.*, 1913, v, 11, 1193.
- ⁹ Kerr, J.: *Medical News*, Philadelphia, 1894, lxiv, 214.
- ¹⁰ MacTaggart, D. D.: *Montreal Med. Journ.*, 1903, xxxii, 432.
- ¹¹ Spiers, H. H.: *Cincinnati Lancet Clinic*, 1900, xiv, 127.

DERMOID CYSTS AND TERATOMATA OF THE MEDIASTINUM

BY

P. J. MOIR, M.C., F.R.C.S.

SURGEON, GENERAL INFIRMARY AT LEEDS

(With Special Plate)

Mediastinal dermoids are more accurately called teratomata, since tissue derived from all the embryonic elements may be found in their composition. Their comparative rarity is borne out by the few cases recorded. Writing in 1929 Robertson and Brown¹ could only find records in the literature of 123 cases. Heuer² traced reports of 134 cases, to which he added four that had come under his own care. Since 1929 Lambert³ has reported two cases, and individual cases have been recorded by Smith,⁴ Robertson,⁵ Cochrane and Nowak,⁶ Crosby and Graham,⁷ and Eggers.⁸ Only two cases have been traced in the records at the Leeds Infirmary in the last ten years.

Case I

A girl aged 16 was admitted to the General Infirmary at Leeds under my care on October 15th, 1935. She complained of pain in her chest, which had been intermittently present for the last twelve months. The pain was felt in the front of the chest, and had also radiated into the right axilla and down the right arm. Occasionally she had difficulty in swallowing.

Previous History.—She had pneumonia as an infant and had suffered from rickets. She has had tibial osteotomies performed.

Physical Examination.—The patient was well nourished and lay comfortably in bed without distress. Temperature and pulse were normal. The right side of the chest appeared slightly prominent, but no swelling could be detected on palpation. Percussion revealed an area of dullness over the front of the right side of the chest, extending from the level of the second rib downwards to become continuous with the liver dullness below. Laterally the dullness extended outwards from the mid-line for about three inches. On auscultation some diminution of breath sounds was noted at the right base and in the axilla.

X-Ray Examination (See Plate).—An antero-posterior view demonstrated a well-defined shadow occupying the upper and inner part of the right lung field. A lateral x-ray showed the tumour to be the size of a Jaffa orange lying more anteriorly than posteriorly.

OPERATION

Avertin 0.08 gram per kilo of body weight was administered with atropine 1/100 grain. Ethylene-gas-and-oxygen anaesthesia under positive pressure was given by Dr. Stanley Sykes.

A transverse incision was made at the level of the fourth costal cartilage, and extended outwards from the sternum to the right side for four inches. The fibres of the pectoralis muscle were split, and the fourth and fifth costal cartilages cleaned and excised. A hard mass, the upper margin of which could not be reached, was felt extending into the right thorax. A vertical incision was continued up from the transverse one, and the third costal cartilage was split. This gave a better exposure, and separation of the tumour from the surrounding structures was begun. The tumour was adherent to the pleura, and this was opened in two places. It was realized that the size of the tumour would make delivery through the wound impossible. Accordingly aspiration was carried out, and 8 ounces of dark grey fluid containing sandy gritty material were evacuated. This considerably reduced the size of the tumour and, after further separation, allowed its delivery into the wound. It remained attached medially to the pericardium, where it proved very adherent. Separation here led to opening of the pericardium at one place. It was decided, therefore, to leave a small portion of the inner wall of the tumour attached, as it was feared that complete separation might lead to tearing of the great vessels and to further damage to the pericardium. The rest of the tumour was removed. The tears in the pleura and pericardium were repaired. This left a large space, but the wound was closed except for a small drainage tube at the lower angle. The operation took two hours and five minutes.

Post-operative Progress.—The patient left the theatre in good condition, and at no time during convalescence was any anxiety felt. Slight infection of the wound occurred and drainage continued for some weeks. She left hospital two months after the operation with her wound soundly healed.

PATHOLOGICAL REPORTS

1. *Fluid from Tumour.*—This contained a few red blood cells and epithelial cells, together with much amorphous debris and some dark pigment.

2. *Tumour.*—This consisted of a cyst showing two projecting nodules obviously covered by squamous epithelium, together with smooth nodules showing on section a variety of tissues. No obvious hairs were seen. Histologically it was a typical teratoma partly lined by squamous epithelium, partly by gastro-intestinal epithelium. Many other tissues were seen: cartilage, mucous glands, a large amount of pancreatic tissue, and some adipose tissue. There was no histological evidence of malignancy.

Case II

A woman aged 25 was admitted to the General Infirmary at Leeds on June 7th, 1928, under Dr. Maxwell Telling. She complained of cough and pain in her chest of twelve weeks' duration. The cough was somewhat productive and the pain, which was not severe, was in the right lower part of the chest and in front.

Previous History.—The present complaint started four months previously, when she had an influenzal cold. She soon recovered from this, but was left with the pain in her chest. Before this illness she had always been well.

Physical Examination.—The patient lay comfortably in bed, and there was no dyspnoea. Temperature and pulse were normal. There was slight prominence of the right chest at the level of the third rib. The movements of the left chest seemed better than the right, and this was confirmed by palpation. No swelling could be felt. Percussion revealed an area of dullness extending downwards from the second rib and merging with the liver dullness below. Laterally the dullness extended outwards to the mid-axillary line. On auscultation there were diminished breath sounds over the area of dullness.

X-Ray Examination.—The report was as follows. Antero-posterior view: There is a large collection of fluid in the right chest. Outlines of shadow are very well defined. Lateral view: The posterior mediastinum is obscured by a dense circular shadow extending behind the shadow of the heart.

Sputum examination was negative for tubercle. A diagnosis of encysted empyema was made and aspiration carried out. The fluid contained necrotic pus cells, but no organisms. Both aerobic and anaerobic culture produced no growth.

The diagnosis was altered to mediastinal dermoid, and the case was transferred to the late Mr. Dobson for operation.

OPERATION

The operation was performed on January 28th, 1928, under open ether anaesthesia. An incision was made internal to the right breast, and the breast was retracted laterally. About three inches of the fifth rib were removed, and the visceral pleura was stitched to the parietal. The skin incision was closed. Twelve days later, under open ether anaesthesia, the incision was reopened. A cyst about the size of a coconut was felt. This was opened and a quantity of yellow fluid evacuated, together with much sebaceous material. The inner wall was scraped out and the cyst was marsupialized and drained.

Post-operative Progress.—She left hospital on August 31st, 1928, with the wound discharging. The discharge continued, and in 1933 she returned to hospital, when the wound was reopened and the cyst was packed and drained. She left hospital with a sinus, which was still discharging when last heard of. The patient has since left the district, and it has not been possible to get her up for re-examination (October, 1935).

Commentary

Rarely teratomata of the mediastinum remain dormant, cause no symptoms, and are found at necropsy. Although congenital these tumours may cause symptoms at any age. Robertson records a case in a boy of 7, and of Heuer's four cases one was a man of 53, but usually symptoms arise in early adult life. Auroiseau⁹ gives a table of fifty-seven cases in which the ages are recorded. In 57 per cent. the age lay between 17 and 30. The sex of the patient does not seem to be a factor of importance.

Symptoms vary and are rarely typical. In uncomplicated cases—that is, those in which infection is absent or in which no large communication with important structures exists—the symptoms that arise are largely the result of pressure. Pain is common, and may be an ache or a feeling of weight situated behind the sternum, or it may radiate in the direction of an intercostal nerve or shoot down the arm. A large tumour may press to such an extent on mediastinal structures that engorgement of the vessels of the neck or even oedema of the face results. Some dyspnoea is present in many cases, and dysphagia has been complained of occasionally. Cough may be troublesome, and in those cases in which the teratoma communicates with a bronchus considerable sputum may be expected. The expectoration may contain greasy material, or hairs may be coughed up and lead to a definite diagnosis. In other patients bleeding into a bronchus might give rise to haemoptysis, and a wrong diagnosis of tuberculosis may be made. If infection occurs, fever, anorexia, and loss of strength result, and the infection may spread to the pleura and cause an empyema.

Diagnosis

These tumours must be considered in the differential diagnosis of other intrathoracic neoplasms and they must be differentiated from certain other pleural and pulmonary conditions. Examination may show some asymmetry of the thorax, or a swelling may be apparent. The swelling may present itself in the suprasternal notch or, more often, may bulge between the ribs, to one or other side of the sternum. In these cases a boggy or doughy consistency of the swelling has been described, suggesting a diagnosis of cold abscess. An area of dullness may be mapped out and auscultation reveal altered breath sounds, or diminished air entry to the affected part of the lung. When the tumour is very large the whole pleural cavity may be filled by it. Fluid aspirated from the swelling may yield sebaceous material, hair, or cholesterol, and these or similar findings will help to distinguish the condition from hydatid cyst of the lung. Exploratory

thoracotomy, should this be necessary to establish the diagnosis, is a wiser and safer procedure than aspiration.

An x-ray examination is always necessary, and both antero-posterior and lateral views are called for. The shadow cast by the tumour appears, as a rule, in the upper part of one or other lung field and the outline is well defined. The direction of the growth may be downwards, when the shadow will appear as resting on the diaphragm. An oblique view helps to determine the relations of the tumour. Areas of calcification may show in the radiograph, and occasionally the presence of imperfectly formed teeth may be noted. Infected cases will not give such clear x-ray findings, and the shadow may be mistaken for an encysted empyema.

Treatment

The aim in treatment should be complete removal. Drainage operations inevitably lead to infection of the cyst, and result in discharging sinuses; the same objection applies to marsupialization. It must be remembered that teratomata are potentially malignant, and this is another good reason for extirpation. If the teratoma is infected, however, preliminary drainage is advisable, as attempts at removal may lead to spread of the infection to the pleura and the mediastinum.

If the tumour is small, and presenting suprasternally, a low collar incision in the neck may give sufficient access. In addition, median sternotomy, with separation of the two halves of the sternum, can be carried out should more space be required. For the large tumours various approaches have been recommended and tried. These include an intercostal incision over the site of the tumour and wide separation of the ribs by a rib spreader, resection of ribs or costal cartilages with or without lateral sternotomy, and the trap-door approach by means of a flap of the chest wall together with rib resection. Cases in which the tumour or cyst is densely adherent to or has perforated important structures may make surgical removal impracticable. More conservative measures will then have to be adopted. Partial removal of the tumour may be possible, or the surgeon may have to content himself with scraping out the contents of a cyst and drainage.

Mortality

The mortality from mediastinal teratomata is high. The best results are obtained by excision in clean cases. Heuer,² in the article already referred to, has published the after-histories of 138 cases. At necropsy twelve were found; in eight details are unknown, forty-six died untreated, and seventy-two were operated upon. Of these seventy-two cases, one had simple drainage of the pleura, thirty-four had incision and drainage of the cyst, sometimes with marsupialization, and thirteen died. Eight recovered with the formation of fistulae, one recovered but was not improved. Five were cured for varying periods, five recovered but the end-result is not stated, and in two the outcome of the operation is not given. Of thirty-four cases, therefore, only five are known to have been completely cured. Thirteen patients had incomplete extirpation; two died and one was going rapidly downhill; nine recovered, but had discharging sinuses; one recovered, but the result was unknown. Twenty-four patients had complete extirpation of the tumour; three died; twenty-one recovered and were cured.

Summary

1. Two cases of mediastinal teratoma are reported.
2. Symptoms and diagnosis are discussed.
3. Treatment by complete removal offers the safest and best chance of cure.

REFERENCES

- ¹ Robertson, C., and Brown, R. E. B.: *Brit. Journ. Surg.*, 1929, xvii, 197.
- ² Heuer, G. J.: *Ann. of Surg.*, 1929, xc, 692.
- ³ Lambert, O.: *Bull. et Mém. Soc. Nat. de Chir.*, 1933, lix, 822.
- ⁴ Smith, R. E.: *Guy's Hosp. Reports*, 1930, lxxx, 466.
- ⁵ Robertson, D. E.: *Brit. Journ. Surg.*, 1931, xviii, 666.
- ⁶ Cochrane, R. C., and Nowak, S. J. G.: *New England Journ. Med.*, 1931, ccv, 1077.
- ⁷ Crosby, E. H., and Graham, A.: *Journ. Amer. Med. Assoc.*, 1932, xcvi, 1789.
- ⁸ Eggers, C.: *Ann. of Surg.*, 1933, xcvi, 135.
- ⁹ Auroousseau, L.: *Rev. de Chir.*, 1924, lxii, 553.

"NEO-NATAL DERMATITIS"

BY

H. CARTER, F.R.C.S., M.C.O.G.

MEDICAL SUPERINTENDENT, MILL ROAD INFIRMARY, LIVERPOOL
AND

H. A. OSBORN, D.M., D.P.H.

SOUTHERN GROUP PATHOLOGIST, CITY OF LIVERPOOL

(With Special Plate)

Dermatitis in various forms is a source of danger to the newborn infant, and its occurrence causes anxiety to those in charge of maternity units. Undoubtedly the type of dermatitis which most jeopardizes the neo-natal infant is that usually called "pemphigus neonatorum" but which we prefer to designate "subepithelial dermatitis." This disease has been surrounded by obscurity; the nomenclature is unscientific and misleading, the aetiology more than uncertain, the many advocated treatments often unsuccessful, and the mortality rate high.

There are no reliable statistics of the incidence of "pemphigus neonatorum." In the last century it was at times a scourge in some hospitals, but the use of antiseptics and an aseptic technique led to fewer outbreaks. From a study of the literature it is apparent that since 1917 there has been an increase in the number of epidemics arising in America, and an examination of the annual reports of maternity hospitals shows that the problem constantly recurs in this country. According to C. O. Stallybrass (1935) an average of sixty cases of infantile skin diseases, mainly "pemphigus," are notified each year by the midwives in Liverpool, but it is impossible to get reliable statistics for the whole country as the disease is not notifiable.

Classification and Diagnosis

The group of diseases comprised by the term "neo-natal dermatitis" includes several distinct entities:

1. Simple infantile dermatitis (Jacquet).
2. Syphilitic dermatitis.
3. Erysipelas neonatorum.
4. Bullous dermatitis due to mechanical or chemical irritants.
5. Pyodermitis: (a) folliculitis; (b) subepithelial dermatitis (pemphigus neonatorum); (c) deep cutaneous abscesses.

In the first three groups the pathology is clear, and there is little difficulty in diagnosis, while the fourth group is recognizable by the sterile fluid in the blisters. The last group is by far the most important. Some observers have found difficulty in distinguishing between folliculitis and subepithelial dermatitis. In most cases the lesion of the folliculitis is no bigger than a millet seed, whereas the blister of the subepithelial dermatitis is about the size of a threepenny-piece. When, however, a group of follicles becomes infected, and the neighbouring pustules coalesce to form a blister, it is not easy to make a diagnosis before it is evident that further spread will occur.

In considering the differential diagnosis, dermatitis herpetiformis, epidermolysis bullosa, and hydroa vaccini-forme must be mentioned; these diseases are indeed stressed in the textbooks, but are rarely met with in actual practice. The blister usually seen is either pyogenic or sterile: sterile blisters are due to mechanical or chemical irritants, or else are of toxic origin. In the leading article in the *Lancet* of June 8th, 1935 (p. 1333), it is stated that the contents of the bullae in pemphigus neonatorum are initially clear, but in contradistinction to this, pus and micrococci were found in every case examined at Mill Road Infirmary, and several hundred blisters were examined. These findings were of great value in early differential diagnosis.

Experience at Mill Road Infirmary

Subepithelial dermatitis had been present in the maternity unit since 1926, and in 1933 fifty-four cases of "pemphigus," and many cases of folliculitis, were recorded out of a total of 1,122 births. The problem needed thorough investigation, and had given rise to great concern previously. In contrast, puerperal pyrexia, fever cases, and maternal deaths were few, and in other respects the results were quite satisfactory in the unit.

The dotted line indicates the births in the Maternity Unit; the shaded area the "pemphigus" deaths; the broken line subepithelial dermatitis outbreaks due to *Staph. aureus*, except where *Staph. citreus* is indicated; and the continuous line total skin lesions (pyodermitis). A, Wards closed and cleaned. B, Silver nitrate treatment begun. C, Laundry inspection. D, Laundry changed.

Note: (1) Value of silver nitrate treatment (B). (2) Curve of cases of subepithelial dermatitis coincides with the number of births. (3) Temporary improvement after laundry inspection (C); cessation of epidemic following change of laundry (D). (4) Pyodermitis curve closely follows subepithelial dermatitis curve and also ceased with the laundry change (D).

In the first series of cases of "pemphigus neonatorum" examined from August to December, 1932, thirteen were investigated, and the cultures contained a growth of *Staphylococcus aureus*. Five patients died, giving a mortality rate of 38 per cent. One of the dead babies was premature, and another had hydrocephalus, so the corrected rate was probably 27 per cent.

The unit had a quiet period during December, 1932, and the first half of January, 1933, then the number of births increased and another case of "pemphigus" occurred. This led to a new outbreak of twenty-one cases in March, 1933, and the wards were closed. Bacteriologically this had particular interest, as *Staphylococcus citreus* was recovered from the blisters. There were at least two strains present: one fermented milk with the production of acid, and the other clotted milk. Further, these strains actively fermented the sugars and produced indole, in contrast to the common strains, which do not ferment the sugars. Clinically the blisters were similar to those previously seen, but in this outbreak there were only three deaths, giving a mortality rate of

P. J. MOIR: DERMOID CYSTS AND TERATOMATA OF THE MEDIASTINUM

FIG. 1.—Case I. Antero-posterior radiograph showing dense shadow in right lung field.

FIG. 2.—A lateral radiograph of Case I.

FIG. 3.—Radiograph of chest in Case I two months after operation. Lung field clear—some increased opacity at lung root.

FIG. 4. The tumour removed from Case I.