Of the four cases treated with Felix's Lister Institute serum two showed dramatic improvement. One further case showed lessening of toxæmia. The remaining case showed no change.

It is our impression that the new antityphoid serum is definitely beneficial when given to the typhoid patient during the first or second week of illness in decreasing the toxæmia and shortening the febrile course.

Acknowledgements

We desire to record our indebtedness to Drs. J. Edgar and R. H. Robertson of the Keylock and Pratt Infirmary for help in the preparation of serum, and to Drs. W. R. Johnston, A. J. Watson, S. N. Cheer, A. E. Towers, M. H. Huang, H. W. Tseng, R. Wong, W. S. Fu, H. T. Chen, and C. L. Kao of the various hospitals in Shanghai, Nanking, Wuchang, and Yunnanfu, who have been in charge of the patients treated.

We also desire to record our indebtedness to Dr. A. Felix of the Lister Institute, London, for much help and encouragement in the preparation of this paper.

BIBLIOGRAPHY

- Chantemesse, A.: *Sérolithérapie de la Fièvre Typhoïde*, Paris, 1907.
Felix, A.: *Lancet*, 1935, i, 799.
Felix, A., and Pitt, R. M.: *Journ. Path. and Bact.*, 1934, xxxviii, 409.
McSweeney, C. J.: *Lancet*, 1935, i, 1095.
Robertson and Yu: *Brit. Journ. Path. and Bact.*, 1936, xlii, 53;
ibid., 1936, xliii, 191.

DYSPHAGIA WITH ANAEMIA IN A MALE

BY

E. WATSON-WILLIAMS, M.C.

SURGEON IN CHARGE, EAR, NOSE, AND THROAT DEPARTMENT, BRISTOL
ROYAL INFIRMARY

The association of dysphagia with anaemia was noted over sixty years ago: more recently a third symptom, stomatitis and glossitis, has been added to complete the syndrome usually known as the "Plummer-Vinson syndrome"—no doubt because we owe our first description of the characteristic changes in the oesophagus to A. Brown Kelly and Dr. R. Paterson.¹ The condition is not of great rarity: I have observed twenty-odd cases in a series of just over 500 patients submitted to oesophagoscopy. But most observers agree in describing it as a disorder confined to women; the finding, therefore, of an apparently typical case in a male subject has the interest attaching to the unusual.

Case Report

A man aged 71 came to the out-patient department on August 6th, 1936, complaining of difficulty in swallowing during two months, with no appreciable loss of weight. I was not there then, but it is recorded that the hypopharynx was full of mucus, and there was no evidence of obstruction to a bolus viewed by *x* rays: a lozenge was prescribed. Six weeks later I first saw him; the symptoms were unchanged, but I noticed that he was very pale, with conjunctival pallor; he had never known he was anaemic. He had also glossitis, with loss of papillae except in the centre of the tongue; the insides of the cheeks were sore, and he had a linear scar from each angle of the mouth, and when questioned said that the corners of his mouth had been sore for several years. The blood showed: red blood cells, 5,000,000; haemoglobin, 60 per cent.; Dr. A. D. Fraser reports "anisocytosis fairly well marked, poikilocytosis moderate, polychromasia moderate, microcytosis fairly well marked, megalocytes and nucleated reds absent; erythrocytes show lack of haemoglobin; hypochromic anaemia." Van den Bergh negative, direct and indirect. Test meal: no free HCl in resting juice; one hour after meal combined HCl in 1 c.cm. neutralized by 0.2 c.cm. decinormal soda. Oesophagoscopy: at upper end of oesophagus a thin white demilune web round the posterior aspect reduces the lumen to 8 mm. across. It yielded readily to slight pressure from the endoscope, with just a trace of bleeding. Below this the oesophagus was normal except for

pallor of the mucosa; cardia at 37 cm. from teeth, normal. Immediately following the instrumentation swallowing has become normal. The patient has been sent home, taking 30 grains of iron and ammonium citrate thrice daily; no dysphagia, the soreness of mouth is less, pallor not materially changed.

Pathology

The aetiology is in dispute. Paterson and Brown Kelly² regard the dysphagia as primary, with anaemia due to interference with nutrition; thus following the older descriptions of anaemia occurring in "hysterical dysphagia."

"At a meeting [of the Laryngological Section of the Royal Society of Medicine] in 1919 Brown Kelly and I drew attention to a clinical syndrome in women, the essential feature of which is a gradually increasing inability to swallow, due to spasm of the pharyngo-oesophageal sphincter; often associated with changes in the buccal and pharyngeal mucosa and at a later stage with anaemia, which is to be regarded as a secondary manifestation of the dysphagia. Plummer, approaching it from the physician's point of view, called it hysterical dysphagia, and later Vinson laid stress on the anaemia and splenomegaly.⁵ The blood changes are those of chlorosis and secondary anaemia with considerable reduction in the haemoglobin (Munro Cameron)."⁶

The patients are women and come under observation in middle life with a long history of gradually increasing dysphagia, which eventually confines them to a fluid diet. Mild glossitis with atrophy may be early; later, the tongue is smooth and the buccal mucosa dry and pale.

On the other hand, it has been suggested that the changes in the mouth and pharynx are secondary to the anaemia. Dr. R. S. Johnson³ reports six cases in St. Bartholomew's Hospital. All were women: the anaemia was severe, of secondary type; haemoglobin 32 to 50 per cent.; colour index about 0.5; complexion striking, brownish-yellow, suggesting pernicious anaemia; glossitis and cracks at the angles of the mouth were noted in four; dysphagia in five; complete achlorhydria in all; slight enlargement of spleen in one only; fragility of red corpuscles increased. He considers the changes in the mucosa secondary to the anaemia. With this view I agree. A feature not included in the above accounts is the formation of a thin web at the oesophageal entrance, instead of the spasm mentioned above: in my own experience this is the more usual finding.

Treatment

The essential is the administration of iron in large doses—inorganic preparations are to be preferred; the daily dose of iron and ammonium citrate is 60 to 120 grains for eight weeks or more. It may be necessary to repeat treatment during several years. Where spasm or webbing is present treatment should begin with dilatation of the oesophagus; this rarely needs to be repeated. The prognosis is good, although it is said that later malignant changes in the oesophagus are relatively frequent in these patients.

REFERENCES

- ¹ *Proc. Roy. Soc. Med.* (Sect. Laryngology), 1918-19, xii, 235.
- ² *Ibid.*, 1930-1, xxiv, 1203.
- ³ *Ibid.*, 1930-1, xxiv, 1206.
- ⁴ *Ibid.* (Sect. Therapeutics), 1930-1, xxiv, 543.
- ⁵ *Minnesota Med.*, 1922, v, 107.
- ⁶ *Quart. Journ. Med.*, 1928, xxii, 43.
- ⁷ *Ibid.*, 1932, N.S., i, 41.

The issue of the *Bulletin de l'Office International d'Hygiène Publique* for October includes articles on cholera in India, the duration of immunity conferred by vaccination in relation to the type of vaccinal reaction, vaccination in Mexico, poliomyelitis in Spain, lymphogranulomatosis in the French Colonies and in Sweden, and the supervision of milk in Turkey.