and about one and a half inches long; it was covered with many thickened and dilated blood vessels.

Improvement was immediate. The tingling and stiffness of the fingers disappeared in a few days, and the patient soon found that he could execute fine movements with his fingers, which had been previously difficult or impossible. Convalescence was normal. He was given massage and sinusoidal stimulation to the muscles. There has been marked restoration of the affected muscles so that at this date, eight months after the operation, there is very slight difference between the two hands. Power has returned to the limb and restoration is complete.

Summary

1. Late ulnar nerve palsy develops many years after the original injury—from three to forty-eight, in this case fifty-eight, years.

2. The external condyle is usually fractured, and non-union may persist for many years.

3. The syndrome consists of the deformity of the external condyle, cubitus valgus, and ulnar nerve palsy.

4. The lesion is an intraneural fibrosis. The prognosis is good if the tension is relieved and the course of the nerve is shortened by anterior transposition.

BIBLIOGRAPHY

- Adson, A. W.: *Minnesota Med.*, 1918, i, 455.
 Bowlby, A.: *Injuries and Diseases of Nerves and Their Surgical Treatment*, 1890, p. 255.
 Broca and Mouchet: *Rev. de Chir.*, 1899, xix, 701.
 Buzzard: *Lancet*, 1922, i, 317.
 Davidson and Horwitz: *Late or Tardy Ulnar Paralysis*, 1935, 17, 844.
 Miller: *Surg., Gynecol. and Obstet.*, 1924, xxxviii, 37.
 Platt: *Brit. Journ. Surg.*, 1926, xiii, 409; *Journ. Bone and Joint Surg.*, 1928, x, 403; *The Robert Jones Birthday Volume*, 1928, p. 195.
 Sherren: *Lancet*, 1906, i, 809; *Edinburgh Med. Journ.*, 1908, xxiii, 500.

SEPARATION OF CONJOINED TWINS

BY

D. W. McLAREN, M.B., B.S.

MEDICAL OFFICER, COLONIAL MEDICAL SERVICE, NIGERIA

(With Special Plate)

The separation of conjoined twins by operation must be rarely practicable. In the case here described the result was successful.

The twins (female), born to a Hausa woman in Sokoto Emirate, were united at the epigastrium by a circular funnel of tissue, which at the operation measured about one and a half inches in width and two inches in diameter. The union was such that the children could be turned away from each other to the extent of about a right-angle only. The sternum formed a continuous U-shaped bar of cartilage about half an inch in width across the top of the isthmus. A small single umbilical scar was visible on the under surface, about its middle. The photograph (see Special Plate) shows well the site and degree of the junction.

The mother, a healthy, well-developed woman aged about 25, had had previously two normal births, the first six years ago, the child (a boy) dying at the age of 10 months, and the second (a girl) now alive and well at the age of 5 years. The twins were born on December 20th, 1935, the estimated time of gestation being about 250 days. Birth was apparently not difficult, though the mother admits that she was in labour for nearly twenty-four hours, which is a relatively long labour for a multiparous native woman. She states that the head of one and the feet of the other presented together, and that the cord was single the whole way from the juncture to the placenta, which came away very shortly after the birth of the infants.

The twins were first seen at the age of 2½ months, when they appeared to be rather emaciated and anaemic, though quite vigorous. It was then apparent that operation was quite feasible, for by compressing the isthmus between the finger and thumb the bowel of the one or the other could be easily reduced into the abdomen, leaving nothing but integument and the bar of sternal cartilage between the

fingers. Since the mother was rather reluctant to agree to operation, it was decided to leave them till the age of 6 months, when it was hoped that they would be more developed and better able to stand the operation. This hope was fulfilled, for on admission to hospital on May 27th, at the age of 5 months, they had grown into quite normal plump infants, though still anaemic. It was more evident now that one was a little more developed than the other.

The blood examination on admission showed a red cell count of 3,600,000 per c.mm. and a haemoglobin index (Tallqvist's scale) of 40 per cent. Two weeks on an iron mixture raised the count to 4,310,000 per c.mm., and the haemoglobin increased to 55 per cent. The mother's blood revealed only a slight degree of hypochromic anaemia (compared with the average standard of the normal native), with a count of 5,040,000 erythrocytes and a haemoglobin percentage of 70.

The Operation

On June 16th the operation was performed under chloroform anaesthesia, without any preliminary preparation apart from the oral administration of glucose. The stronger twin was anaesthetized first in the hope that the other would also go under, but after ten minutes of full anaesthetization no obvious effect was observed on the second twin, and she had to be separately anaesthetized. The operation presented no great difficulty. The incision was made vertically through the middle of the junction, and on opening the peritoneum it was observed that there was a vertically placed peritoneal septum in the upper half with a "foramen" at the lower part, through which bowel from one or the other could easily pass. The only structure divided, apart from the integument, peritoneum, and the sternal cartilage, was the ligamentum teres of the larger twin's liver, and this was found traversing the upper part of the isthmus extraperitoneally. Its connexion with the umbilicus was not traced.

As soon as separation had been effected the infants were placed in the supine position, and each wound was sutured up vertically in turn. No attempt was made to trim the projecting cut edges of the sternum or the skin, and only excess of parietal peritoneum was excised. The rectus aponeuroses were thin and attenuated, but the recti themselves seemed undeveloped in this area; after operation it was evident that this was so, since there was in each around the site of union an oval-shaped area, which was quite devoid of muscular tissue. The gap was about two and a half inches in its widest part, and there was no muscle attachment to the lower ends of the sternum. The cut ends of this have straightened out since the operation; this makes the bulge of the abdominal wall in this area less obvious than it might have been had the sternal ends been resected. There is still the probability, of course, that they will develop large umbilical hernias later.

Apart from a slight attack of enteritis in the weaker infant, and a stitch abscess in each wound, the post-operative period was uneventful. The mother was able to breast-feed the children from the first day. The weights just after the operation were 10 lb. and 12 lb. respectively, and now, six weeks later, 13 lb. each. The blood count has improved to 4,900,000, and the haemoglobin to 60 per cent.; they are now in every way normal and vigorous infants.

It would be interesting to know if this operation is unique. It was exceedingly fortunate that the twins were united in what might be termed "the least degree," each having separate organs.

I am indebted to the Director of Medical Services, Nigeria, for permission to publish this article.

W. R. H. Kranenburg and J. J. Wallien (*Nederl. Tijdschr. v. Geneesk.*, August 15th, 1936, p. 3715) state that the notification of forty-one cases of lead poisoning in cigar makers in a certain district of Holland led to an inquiry which proved the existence of the sources of intoxication to be the zinc cutting sheet and the drinking-water in the factories and in the workmen's homes. The authors are of opinion that the zinc cutting sheet was a possible source of intoxication, though only in a very slight degree, whereas they regard the water containing lead as a very important factor.

F. GLYN-HUGHES AND R. C. WEBSTER: JOINT DISEASE BEGINNING DURING ACTIVE
TREATMENT OF SYPHILIS

FIG. 1.—Case I. Skiagram showing Charcot's disease of the spine.

FIG. 3.—Case II. Skiagram showing Charcot's disease of hip-joint with posterior dislocation.

FIG. 2.—Case I. Skiagram showing Charcot's disease of left hip-joint, with pathological fracture.

D. W. McLAREN:
SEPARATION OF
CONJOINED TWINS

Photograph showing site and degree of the junction.