TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some diseases of the digestive system met with in general practice.

TREATMENT OF FUNCTIONAL DYSPESIA

BY

C. M. WILSON, M.D., F.R.C.P.

The treatment of functional dyspepsia has little to do with the prescription of particular drugs and special diets; stripped to its essentials it consists in the careful exclusion of organic disease and in the ability to convince the patient that this has been done. It includes, too, the search for a cause of the dyspepsia, in order to remove it, or, if that is vain, such reformation of his or her way of life as may be possible. It is customary, and far from irrelevant, to preface an article on the treatment of functional dyspepsia by underlining the difficulties in diagnosis, since the most effective psychotherapy is a good history and a thorough examination.

Elimination of Organic Disease

In the elimination of organic disease—a matter beset with many difficulties—an accurate history is almost everything. A physician of quality in his search for truth will contrive to ask one and the same question in three different ways. He is obedient to the laws of evidence: he asks questions, he does not suggest answers, and he begins at the beginning. A patient under suspicion of malignancy complains of epigastric pain since October, but with a little gentle jogging of his memory he will presently own that he had similar pain in October a year ago. A single question has cut out a gnawing anxiety. In this field to listen patiently is the beginning of wisdom. We want to hear not so much what patients eat and drink, but what they do in the day. Is the complaint of pain really pain, or only discomfort? If it is pain an organic origin is probable; if the pain is also felt in the back, if it wakes the patient and makes him afraid to eat, or if it is so severe that to escape from it he makes himself vomit, then an organic diagnosis may be provisionally made. There is a precision about pain in organic disease which is wanting in functional dyspepsia. Ulcer pain, for example, appears after food by the clock; it is present for a certain time, measured in days or weeks, and then vanishes completely for months. The seat of the pain is indicated economically by one finger, and not by a gesture which embraces a considerable portion of the abdomen.

There are certain organic diseases always in our mind when we are about to make a functional diagnosis; they must not be lightly dismissed. In particular, we are aware of the quiet approach of cancer of the stomach and the difficulty of an early diagnosis. When a man of middle age, or more than middle age, first complains of dyspepsia, accompanied by loss of weight, of strength, and of appetite, and when his symptoms do not quickly respond to treatment, then carcinoma of the stomach is probable and laparotomy justifiable. The precision of ulcer pain has already been contrasted with the lack of definition of functional discontents; it yields to alkalis and leaves the patient in peace for months at a time; such intervals

of complete freedom are seldom the lot of the victim of functional disorders. Local tenderness and some muscular rigidity may help, but if after a careful analysis of the character of the pain we have not made a provisional diagnosis probably none will be made until the radiologist repairs the omission, a sequence of events which soon becomes a habit.

Reflex Dyspepsia

Much has been written of appendicular dyspepsia, but for my part I do not care to make such a diagnosis without good evidence that the appendix has been acutely inflamed. It is, however, the common dyspepsia associated with disease of the gall-bladder which so closely resembles a functional disorder that the very elect may be taken in error, and it is an everyday experience to see gall-stones removed from the abdomen of a woman who for ten years had worn a functional label. Flatulence and distension after meals, punctuated by attacks of nausea, in an obese middle-aged woman whose eyes are sometimes tinged yellow and whose urine at such times contains a trace of bile, can often be attributed to the gall-bladder. If the inflamed organ is brought down by a deep breath to fingers pressed up under the costal margin a sharp stab of pain may confirm the diagnosis.

Valuable as the history is in making a provisional diagnosis it is wise to support it when we can by other investigations. I find I have recourse to the test meal as a diagnostic measure less frequently than I did some years ago. An excess of mucus may unmask a chronic gastritis which would otherwise have escaped detection. Foul-smelling stagnant fluid, siphoned from the fasting stomach, would be valuable evidence of carcinoma of that organ. The test meal is serviceable in other conditions, but I do not think in the judgement of those with most experience of dyspeptic disorders it has vindicated its claim as an essential part of routine diagnostic technique. The position of radiology, on the other hand, is not open to question, though the amateur in this field is nothing less than a dangerous nuisance.

Approach to the Patient

If it is important to exclude so far as we can the possibility of organic disease, it is as important to convince the patient that this has been done. To fail in this is to fail in everything. Yet failure is common, and the reasons are not far to seek. There are physicians who are handicapped by temperament in their approach to functional disorders. Why should they bother to search for a cause in the patient's mode of life when it is the way these weaklings are made, and not the stresses to which they have been subject, that is responsible for their dyspepsia; when, indeed, the demonstration of a neuro-pathic constitution is in their eyes essential to the diagnosis. It is an attitude as old as the Pharisees, and familiar enough to those who were concerned with shell shock in the war. Under the strain of active service an unstable man soon gave way, but if the test was sufficiently severe and sufficiently prolonged anyone, however sound his stock, might develop functional disorders.

In like manner the mind may be hurt in peace as in war, and functional dyspepsia may develop in anyone if

the wear and tear of life fastens on them and will not let go. It is our first concern to make that plain to the patient. He has been told that if his malady is not organic it is less than nothing, that he is a bundle of nerves, and that his troubles are imaginary. We have to convince him that we are not for one moment in doubt as to the reality of his sufferings, and are interested in them and in him. Then, when our sympathy and understanding are no longer in question, he will be prepared for a simple explanation of how an emotion such as fear can, through an endocrine disturbance, affect two factors of digestion—the movements of the stomach and the vascularity of the mucous membrane. In this way it is made clear to him that the cause of his dyspepsia is some mental process, which must be removed before he can get well. He begins to understand his own problem of health, and to see that it can never be solved by drugs or by dieting. He is ready now for some radical readjustment in his way of living. He will prosper according to our success in removing the cause of his dyspepsia, or if that is beyond our reach he may have to confess that he has not learnt to face adversity with equanimity, that he has failed to build up a working philosophy which would help him to make the best of things, as cheerfully as may be.

I have dwelt on our reward if the cause of the dyspepsia is detected and removed, but we shall be spared disappointment if we can recognize a group of dyspeptics for whom dieting, symptomatic treatment, and psychotherapy are equally futile. No disorder of gastric function can be demonstrated in these patients. They are suffering not from a phobia but from a delusion, and they can be known from their reaction to attempts to explain to them that their symptoms have a nervous origin. Such an explanation is not tolerated. They are convinced there is an organic basis, adhesions, or what not. They hanker after an operation. Reasoning with these people is seldom profitable. It would not be right to leave the question of causation without emphasizing the importance of extirpating from the patient's mind the fear of cancer. Hidden away, it must be routed out and dealt with firmly.

Food and Medicine

In the literature devoted to the nervous dyspeptic it has not been customary to set aside as little space for food and medicine as I have done. It is, of course, a matter of emphasis, a question of degree. Those most sceptical of drugs and diets are not independent of them. It may be profitable, therefore, to review their proper usage, and in so doing it is convenient to follow the common practice of distinguishing two groups of functional dyspepsia—the hypersthenic and asthenic. Hurst believes that a man's liability to indigestion depends upon his constitution, that it is a congenital weakness. The majority of people, which he puts at 80 per cent., escape this defect, and will probably grow old without ever suffering from chronic indigestion. Of the remaining 20 per cent. half, he believes, are born with the hypersthenic diathesis and half with the asthenic diathesis. (These percentages, I imagine, are meant for precision of speech, and are not intended to bear the scrutiny of pedantic statisticians.) If these 20 per cent. are obedient in their lives to the laws of health, if they are moderate in the use of alcohol and tobacco, if they eat slowly at regular intervals when not overtired, and remain free from sepsis, then, Hurst believes, they will not develop dyspepsia, and chronic indigestion will become a rare disease. But if they neglect these precautions they will suffer from chronic gastro-duodenitis.

In the war this doctrine was tested on a vast scale: all these laws were set at naught and the dentist was forgotten, but it was, I think, the common experience that

those who in city life had suffered from functional dyspepsia shed their symptoms in the open air and enjoyed a kind of gastric renaissance. Hurst is reluctant to make a diagnosis of nervous dyspepsia; he appears to be feeling his way towards a total renunciation of the view that disorders of the stomach, apart from hysterical vomiting and the like, exist without structural change. I believe that chronic gastritis is often present in patients who have been labelled functional dyspeptics, but a faith of that kind cannot for one moment blind one to the dominant part played by the mind in the disorders with which this article is concerned. I cannot credit that a gastric millennium will appear on the earth when men drink less, smoke less, and go more often to the dentist. It is more likely to come if and when England ceases to be an industrial state with a large population huddled in cities, and returns to the pastoral conditions for which most of us were designed.

Hutchison recognizes the same two groups, and believes that everyone has a tendency in one or the other direction. But he pictures the symptoms of these groups brought about by nervous instability acting in the hypersthenic group through the vagus, and in the asthenic group through the sympathetic. It may be serviceable to consider the two groups in detail.

The Hypersthenic Group

The hypersthenic individual is, commonly, a man whose long days in the city or in the practice of a competitive profession have left him, as the price of success, in a state of nervous irritability. The movements of his short stomach, which is tucked up under the ribs, are exaggerated, so that it is soon empty. The secretory output is likewise increased, but with less certain consequences. He complains of acidity when his stomach is full and of discomfort when it is empty.

The diet is planned to meet this secretory surfeit. But it is significant that, while some physicians give protein freely to fix the free acid and limit sugars and starches, others hold that protein foods only stimulate the gastric juice, and give carbohydrates with a full hand. Both are generous with fats and parsimonious of all substances that stimulate secretion—condiments, extractives in soups and gravies, alcohol, and the like.

The excessive secretion can be discouraged by moderation in smoking and by 5 minims of the tincture of belladonna given in an ounce of water before each meal, and be neutralized by an antacid powder, to which 6 grains of bromide are added, taken in water or milk one hour after food:

R	Sod. brom.	...	...	...	...	gr. vi
	Sod. bicarb.	}	...	...	...	āā gr. xv
	Mag. carb.					
	Bis. oxycarb.					

The Asthenic Group

In this group the patient is often a woman of poor physique, passing perhaps through the climacteric, who has never put up much of a fight against circumstance. She complains of sinking sensations when the stomach is empty, of flatulence and fullness. She may wear an abdominal belt, but, whatever is done for her, exhaustion and depression seem to be her lot. The movements of her long, low stomach are so impaired that it empties slowly. Digestion is slow and the secretions diminished.

The diet is drawn up to suit a stomach the movements of which are impaired. Certainly small, frequent, and dry meals (fluids being drunk between meals), without bulky foods such as cabbage, potatoes, and most vegetables, leave the patient more comfortable. All those foods which were banned in the first group because they

stimulated secretion are here rational. A prescription such as the following, taken before meals, stimulates gastric secretion, and is helpful.

R̄ Sod. bicarb. ... gr. x
Tinct. nuc. vom. ... m v
Inf. gent. co. ... 3 ii
Aq. chlorof. ... ad 3 i

It is a common practice to prescribe hydrochloric acid for members of this group.

R̄ Acid. hydrochlor. dil. ... 3 ss
Syrup. aurant. ... 3 ij
Aq. ... ad 3 ss

A tablespoonful in five ounces of water sipped at meal times.

Prescribed in this manner it makes a rather uninteresting beverage of the same complexion as weak lemonade. It is often given so that the unpleasant acid taste is not disguised, when it is likely to defeat the very end for which it was prescribed. If the bowels are obstinate, cascara, aloes, rhubarb, and senna are available. Rest in bed or a holiday, according to the severity of the case, is beneficial.

Re-education

It is all a little too tidy to be convincing. There is no proof that an increase in the acidity of the gastric juice can cause symptoms. Heartburn is not due to regurgitation of acid gastric juice into the oesophagus; it occurs with achlorhydria. Where there is impairment of function it would seem reasonable to counsel dietetic restrictions. If the stools are full of coarse undigested material which is found to be cellulose the symptoms may vanish when foods containing cellulose are left out of the diet, and the stools become normal. But with experience we look less and less to drugs and diets for salvation in this form of digestive disorder. They have been fully sampled before we are summoned. We have to wean these people from their preoccupation with the digestibility of various foods, not to encourage them in their researches. Their eyes follow what they eat in its passage through the alimentary tract; they must be turned to more useful occupations. We have to resist the importunate in their demand for medicine and a diet, to convince them that there is no easy path to health along these lines. It has been well said that the patient who is the subject of this article, like the diabetic, is curable only in so far as he can be educated and taught to care for himself.

In the hypersthenic group much may be done by a tenacious attempt to induce the man in question to conduct his life on more rational lines. Here is a man with a liking for ball games and some proficiency in them, living in Hampstead a few minutes from a golf links, who drags himself out of bed on a fine summer day at 8.45 a.m., when he might have risen with his servants two hours before and come to his breakfast with an appetite born of exercise before the good of the day has gone and he himself is used up. If he can be led to talk of his golf instead of his insides we shall not need our "do's" and "don't's" in dieting, our alkalis and belladonna, though we shall not necessarily earn the gratitude of his family by the change. We too readily accept the protestations of members of this group that they cannot possibly alter their lives. These men are often intelligent; they will listen to close reasoning—it is their own coinage—and they will respect the force of character which prevents their physician hurrying along the line of least resistance. It may be necessary to put it to them that they must choose between living with their symptoms and the reformation of their ways, remembering that few really efficient people are too busy to do what they wish to do.

Our task in dealing with the women in the asthenic group is of another and less hopeful kind. Here the affront to the laws of health is less patent, the inborn error in her architecture more obvious. This woman is

only busy with her discontents; if she had more to do we might hear less of them. At the best her recovery of health is going to be a long-drawn-out affair. But she, too, has a cause for her exhaustion; if it can be found and removed we shall be well rewarded.

I shall end as I began by affirming our independence of drugs and diets. In diagnosis we shall get what we want from the patient's discourse, and in treatment we depend on that sympathetic insight into another mind which is the beginning and the end of all achievement in this field.

A NEW NATIONAL LIFE TABLE

IMPROVED EXPECTATIONS OF LIFE

The Registrar-General's decennial supplement—England and Wales, 1931, Part I, Life Tables—has just been issued. It consists of a report by the Government Actuary, Sir Alfred W. Watson, on the mortality experience of England and Wales in the three years 1930, 1931, and 1932. The experience of males and females has been examined separately.

The principal table is a new national life table for England and Wales, which continues the series of life tables constructed after each successive census, and is henceforth to be known as English Life Table No. 10. An indication of the trend of mortality in recent years is afforded by the comparison of the expectations of life according to English Life Tables, No. 8, No. 9, and No. 10, which may be taken as representative of the conditions existing in 1911, 1921, and 1931 respectively.

Expectation of Life (Years) : English Life Tables

Age	Males			Females		
	No. 8 1910-12	No. 9 1920-2	No. 10 1930-2	No. 8 1910-12	No. 9 1920-2	No. 10 1930-2
0	51.50	55.62	58.74	55.35	59.58	62.88
10	53.08	54.64	55.79	55.91	57.53	58.87
20	44.21	45.78	46.81	47.10	48.73	49.88
30	35.81	37.40	38.21	38.54	40.26	41.22
40	27.74	29.19	29.62	30.30	31.83	32.55
50	20.29	21.36	21.60	22.51	23.69	24.14
60	13.78	14.36	14.43	15.48	16.22	16.50
70	8.53	8.75	8.62	9.58	9.95	10.02
80	4.90	4.93	4.74	5.49	5.56	5.46
90	2.87	2.82	2.63	3.16	3.13	2.98

Analysis of Mortality Rates

An examination of the rates of mortality at individual ages shows even more emphatically the great improvement in the vitality of the people which has taken place at all but the most advanced ages. The improvement is specially marked at the youngest ages. The probability that an infant will die within a year of birth has decreased between 1931 and 1911 by as much as 40 per cent. Out of every 1,000 boys born the number who died before attaining the age of 1 year was in 1911 about 120, in 1921 about 90, but in 1931 about 72 only. Out of 1,000 girls born the numbers of deaths in the first year of age were 98 in 1911, 69 in 1921, and 54 in 1931. During childhood and adolescence also the improvement in mortality experience though not so pronounced has still been substantial, while from ages 20 to 60 the 1931 mortality rates are much less than those for 1911, the difference being greatest round about age 40, where it is well over 30 per cent.

In contrast with this remarkable improvement in the mortality experience during the greater span of life it is found that at the advanced ages the 1931 rates of mortality are rather heavier than those for 1921 or 1911. The deterioration as compared with the 1921 rates first becomes apparent at age 69 in the case of males and at age 78 in the case of females. One cause of this phenomenon is probably the survival to old age in the present generation of many of the