THE PROBLEM OF SUICIDE*

BY

C. STANFORD READ, M.D.

LECTURER IN PSYCHOLOGICAL MEDICINE, BETHLEM ROYAL HOSPITAL ;
PSYCHIATRIST, WEST END HOSPITAL FOR NERVOUS DISEASES

There are but few human problems that do not, directly or indirectly, come within the purview of the general practitioner. The problem of self-destruction does not frequently present itself to him, but he never knows when he may have under his care some patient who may be potentially suicidal, whether that fact is recognized or not, or when he may be called to someone who has already attempted or completed the act. Popularly, suicide is usually regarded as rather a simple business. Somebody was in great trouble, severely depressed, or in acute pain. Life in such circumstances seemed unbearable and so the great refuge was sought.

A scientific study of the subject so often falls back upon barren statistical analyses, and medical literature ignores suicide as if it were scarcely entitled to recognition as a cause of death. One reads daily how this form of death is the logical consequence of circumstances, particularly ill-health, discouragement, financial loss, humiliation or shame, or unrequited love. It is assumed that the forces impelling suicide come wholly from without, whereas the paramount factors determining behaviour are the impulses from within. Few of us have been free from severe stress some time or other, but, though temporarily we may have seen life "as through a glass darkly," no impulse to end it took serious possession of us.

Yet there are many who voluntarily seek death on apparently slight provocation, even though they would seem to have much to live for. Evidently some strong inner compulsion overcomes the self-preservative instinct and leads them to this final goal. Psychologically, suicide is a complex act, but before its deeper mental factors are considered a survey of the many data concerning the subject will reveal much of interest.

Statistics of Suicide

It is generally assumed that the suicide rate is on the increase, and it has certainly shown an upward trend since 1920. Though during the great war period it was temporarily much diminished, there had been previous fluctuations which are by no means easy to explain. In England the rate is about 12.7 per 100,000 population.

It is interesting to note the great differences in the rate that are found in the various countries of the world. Austria, France, Switzerland, Czechoslovakia, and Japan have led the world in frequency of suicide. The high rate in Japan is partly due to the fact that the act is founded upon traditional custom and sanction. The United States, Sweden, Belgium, Denmark, and Australia hold a more middle position, while Norway, Holland, Spain, Portugal, Italy, England, and the Irish Free State are those with the lowest suicide mortality. In the Irish Free State it falls as low as 2.8 persons in every 100,000, while the figure for Spain is only 3.5. In Sweden, France, Denmark, and Spain suicide has decreased, while in all the other European countries the current tendency is decidedly upwards.

Contributory Causes

It is readily understandable that as age progresses and as we have less and less to look forward to, in this world at any rate, the tendency to self-destruction should become increasingly in evidence. More than half of all suicides occur among those of 45 years or over. Although in childhood suicide, as we might suppose, is not much of a

social problem, it is more common than one might expect, and in German countries there have been astounding outbreaks of it among school children that seemed almost epidemic. From puberty onwards the rate rises steadily.

On the whole, suicide may be looked upon as a masculine type of reaction. On the average about three and a half times as many men as women kill themselves, though the ratio differs a good deal at various ages. In men the rate rises steadily throughout life, the upward incline being marked at the advanced age period. In women the rate increases very slightly up to the age of 30, and then remains stationary afterwards.

In most European countries the rate is distinctly higher in the city population than in rural districts, although in Great Britain there is relatively little difference. Naturally one would be inclined to associate urban life with stress and factors tending to instability, and especially so when we bear in mind that insanity has a higher incidence in towns as compared with the country. However, the more we investigate all these data relating to suicide the more we shall find it difficult to correlate cause and effect amidst the multiplicity of possible factors.

Economic Status and Religion

Statistics tend to show that those of good social status and business positions end their own lives as frequently as poor people. In general, persons with good means are engaged in intellectual vocations, and those thus employed seem to be more liable to suicide than the manual labourer. Yet the study of occupational groups shows that this is by no means of universal application. In this country it is striking that among those with the highest rates are physicians, dentists, barristers, and solicitors, and other professional workers such as teachers have low rates, while clergymen of certain denominations stand near the bottom of the list. In the labouring classes a high rate is most frequent among the poverty-stricken, so that suicide appears to be most prevalent at both extremes of the social scale.

We shall not be surprised that religion has a potent influence, and statistics demonstrate that those countries where the population is predominantly Catholic have a much lower suicide mortality than where Protestantism largely reigns. The Irish Free State has only a rate of 2.8 per 100,000, Spain 3.5, and Italy 9.6; while the Protestant nations show very much higher figures—for example, Germany 27.8, Denmark 17.6, and Sweden 15. During the last century the Jewish people had a very low suicide rate, but it is now higher than the Catholics and Protestants. This is partly accounted for by the fact that the financial and industrial depression following the war affected the Jews very seriously, and that later there was a recrudescence of anti-Semitism.

Historical Factors

It is of interest to note the attitude of ancient forms of civilization to suicide. Most authorities seem to be agreed that it was not considered a sin in the Greek or Roman State; that it was not infrequently extolled in old legends and in Homer; that it was opposed by such early philosophers as Plato and Pythagoras, but that the later Stoics rather actively encouraged it. There were many factors in those days which raised suicide to an extraordinary prominence. Life generally was held cheaply.

On the advent of Christianity quite the reverse attitude began to be adopted gradually, and by the fifth century suicide was definitely condemned by ecclesiastical law. As life became regarded as a divine gift, he who took his own life sinned against God. Through the severe denunciation of the Church there arose those barbarous and revolting customs which were carried out upon the corpse of the suicide. At any rate, whether due to the

* Abridgement of a lecture delivered to the East Kent Division of the British Medical Association at Margate.

injunctions of the Church or the fear of these indignities, the fact remains that throughout the Middle Ages suicide was almost unknown.

By the seventeenth century, however, a new philosophy of life had begun to modify opinion on the subject, and the individualism of that period again placed the right to die in the hands of the person himself. Hume and Voltaire in their eighteenth century writings warmly defended suicide, and Schopenhauer pointed out that obviously no one had a greater right over anything in the world than over his own person and life. In the nineteenth century and subsequently the subject received an increasing recognition from the scientific aspect. The Orthodox Church still condemns the practice, and the law regards it as a felonious crime, while the coroner's jury commonly returns a verdict of "suicide while of unsound mind" to avoid the infliction of any penalty or stigma, notwithstanding the fact that usually there is little or no evidence pointing to such a conclusion.

In the Orient suicide has had an intimate relationship with traditional customs. Among the Eastern religions it would seem that both Brahminism and Buddhism have had encouraging effects, while Mohammedanism has severely condemned self-destruction. It is commonly supposed that the main sources of suicide lie somehow in the factors of civilization, and that it is unknown in primitive people. On investigation, however, we find that though in some regions it is non-existent, in others it occasionally occurs, and that there are still others in which it is quite prevalent.

Naturally enough, judging perhaps from our own feelings, we should expect that suicide would be most in evidence during the most gloomy months of the year. It is surprising, therefore, to find that in reality the death rate from this cause attains its highest limit in May and June. Through the following summer months the rate decreases until the lowest point is reached in November or December, and subsequently it gradually rises once more. Many conjectures have been made in an attempt to explain this seasonal variation, none of which, however, is satisfactory. It is plainly evident that suicide is not a reaction to the hardships associated with the winter months.

Suicidal Methods

Many points of interest arise when we review the various methods of committing suicide. As one might suppose from their instinctive make-up, women choose the less violent ways. The largest proportion of suicides among males is by hanging or strangulation, while among females drowning accounts for one-quarter of the total. Gas and other poisons come next with women, and in men cutting instruments and drowning. The use of firearms for this purpose here is comparatively infrequent compared with some countries, such as America, where among men it heads the list.

The enormous increase in the employment of coal-gas for this purpose has been very marked of late years. This can be well understood when it is realized how much more the gas stove has become a household requisite in modern times, and how the simplicity and handiness of such a method must appeal. Without doubt, from time to time, some special mode of self-destruction tends to become the vogue, and is imitated through its suggestive effect. In London the tube railway station tracks have had their allurements, but the digging of a deep pit beneath the rails seems to have stemmed the attempts in this particular way. The power of suggestion is often a big factor in determining not only the resolve to die but the method to be used, so not uncommonly we have almost an epidemic wave of suicide in which life is taken in a precisely similar manner.

The choice of how death shall come about depends often upon the most available and convenient means at hand, but the removal of such means from one who has made up his mind will only result in its being attempted in some other way. The determination of method, however, is frequently related to deep psychological factors, and the choice is thus perhaps unconsciously guided. One can the more easily conceive of there being at times some such hidden urge when we hear of some exceptional and dramatic methods being used, such as thrusting a red-hot poker down the throat, lying down before a steam-roller, leaping into a volcanic crater, plunging into molten glass, or self-crucifixion.

Psychological Undercurrents and Motivations

Through psychological study some light may be thrown upon the attendant mental undercurrents and the motivations that urge towards the act. Naturally enough, it will be regarded as a mode of refuge. There is much that can be said in support of such a point of view, for without doubt this is psychologically true. So frequently life is ended because of some difficult problem that seems impossible to solve. Poverty, ill-health, remorseful guilt, or loss of love are the circumstances that may finally constitute the crucial determination.

Nevertheless, we must look upon such causation as quite secondary, and seek the primary source in the hidden depths of the human personality. A little reflection tells us that there are many who are not thus defeated in the face of terrible ordeals and who continue to live on in circumstances which would seem to be intolerable. This at once shows us that the determining factor in suicide is really a subjective one. Though objective influences, of course, must be taken into account, they are not sufficient provocation alone to self-destruction; whether or no suicide results will depend mainly upon unconscious mental forces.

Those who are unacquainted with modern psychological knowledge will not adequately realize how much mental functioning takes place in an unconscious mind, and how much mental conflict can be operative without conscious awareness of the fact. In the course of development there are gradually built up in the depths of the mind buried tendencies of opposing forces related to the battles that have ensued in the past between instinctive desires and their fulfilment or frustration. All our minds are replete with contradictory emotions, which impel us in contrary directions. So, according to circumstances and environment, we may be torn between the urges related to love and hatred, pleasure and pain, activity and passivity, life and death.

These conflicting states are the residue or reactivation of the experiences of child life which centred around those who were near and dear. Through disappointment and the frustration of instinctive desires the child will hate and feel great hostility towards those who stand in the way of such gratification. Such hostility commonly involves a death wish against a near and dear one, but this becomes repressed into the unconscious mind because the contrary emotion of love that gains the upper hand renders hate and hostility repugnant to the personality. We must, however, understand that to the child's mind the desire for a person's death has not the significance it has to an adult. It simply involves the idea of removal from the environment. Those who constantly practise analytical psychology repeatedly meet with evidence of these death wishes that in early life had been repressed and in later years became the source of mental suffering and disorder.

When the young develop on normal lines these hostile mental components are lessened and subjugated, and much of their force is drafted off into useful and constructive

channels. In less favourable circumstances they are generated in the mind in an exaggerated degree, and lie repressed in its depths, where, indirectly, they are dynamic, and are apt to be productive of mental disharmony in some way or other. Because the idea of hatred in connexion with those whom we are supposed to love is repellent to us, it is but seldom that such a feeling, even if only temporarily felt, will be frankly acknowledged. It is only a question of the degree of hatred that may be felt and the degree of stability and control that is possessed which will determine whether aggressive action, or perhaps even killing, follows. Since we are all potential haters, we are all potential murderers. Psycho-analytic research has given us excellent evidence for being able to say that one reason for self-destruction is that the suicide desires the death of another person, and that in this way he acts out upon himself what he wished to inflict upon someone else.

Freud's "Death-instinct"

How is it, though, that such a powerful instinct as that of self-preservation is overcome? In order to have further insight into the problem we must consider several more psychological factors. Freud has put forward the interesting speculation that there is both a life-instinct and a death-instinct. He thinks that, besides that instinct which constrains every living thing to cling to life, there is also sometimes operative a desire for non-being and the achieving of a state of passivity free from all stress and conflict. Usually the life force is far stronger, because there is a self-love which is so powerful that at the menace of death the personality cannot connive at its own end.

It can be said that there may always be a death-instinct in man which in the healthy is kept in check by a life-instinct which is more or less synonymous with love. When in certain constitutional types of persons there arise seemingly insurmountable barriers in the path of Eros, the death-instinct is only too liable to assert itself. Love then turns to hate and aggression, and the death-instinct which had been directed towards the outer world is turned back against the individual himself.

We can well understand how aggression outwards and inwards may be intimately related. According to its direction homicide or suicide may be the outcome; not infrequently we hear of those who murder someone and end their own life directly afterwards. Recent criminal statistics showed that very nearly one murderer out of four committed suicide, and where the act was unsuccessful the figures were very similar.

The Guilt Factor

Besides these factors of unconsciously motivated hate, aggression, and death-instinct, a great sense of guilt is almost invariably a potent force. The old Judaic law of "an eye for an eye, a tooth for a tooth," etc., comes into play. The act of suicide then becomes one of retribution, a self-punishment, a necessary penance for the sinful hostile wishes entertained. It is the death penalty, sentenced upon the self and carried out by the self. The self-punishment may sometimes be related to the idea of revenge, and hence punishment to both subject and object. "See what you have driven me to; you will be sorry for the rest of your life," would express the feelings of many a suicide who regards himself as slighted in love, and in this childish way seeks to get even with the object of his passion. The staging of a drama, with himself as hero, seems to have a lure. Fears of various sorts which have a morbid origin and are commonly linked up with unconscious guilt are secondary motives to self-destruction.

More Obscure Factors

There are occasions when no outward provocation can be discovered. We may be sure, however, that there was present an inner mental problem which could not be faced, and that, however normal the individual may have seemed to others, he was a sufferer from some of those unconscious warring psychological forces mentioned previously. As an incentive there is not infrequently an idea that death will involve eternal union with a lost loved one, or some vaguer conception of a union with God, or becoming thereby in tune with the infinite.

We may speak with much truth of chronic suicide. There are many for whom, for some reason, life has little value, and who therefore pursue lines of conduct that so evidently lead to illness and death. It may be also that many who died from fatal accidents really had suicidal predispositions. Anyone thus mentally constituted becomes careless of his foothold when mountain climbing, disregards danger notices when bathing, and is generally indifferent to traffic warnings. Being neglectful of health in general may have the same unwitting purpose in view. The psycho-analyst looks upon suicide as an illustration of the inherent desire in the mind of man to evade the difficulties and problems of life, to return to the source of life, to return to the mother and again to be in that pre-natal state where all needs were supplied and no responsibility existed. In the face of disappointment and frustration of instinctive desires, the mind in those so predisposed may easily regress.

Suicide and Mental Disease

When we deal with the relationship between suicide and mental disease the first question that presents itself is whether those who attempt or commit the act are necessarily insane. On this point there are conflicting views, but it would seem that there are a fair number of so-called normal people who do end their own lives, though without doubt certain types of psychotics are much more liable to do so. It is useless to take the extreme standpoint that suicide itself is sufficient evidence of insanity. It is quite impossible to draw any hard-and-fast line between normality and mental disease. Similar mental mechanisms and processes are at work in both. We know very well that there is no such thing as "temporary insanity" according to the popular notion, and that the coroner's court only brings in such a verdict, with little, if any, supporting evidence, in order that the stigma and any legal penalties may be evaded.

It is quite clear that a good proportion of those who commit suicide are afflicted with mental disease of some sort. Dr. East, when he was at Brixton Prison, analysed 1,000 consecutive cases where the act had been attempted, and found that about one-fifth were certifiable as insane. We must in the so-called normal cases predicate a suicidal predisposition through the possession of those unconscious conflicts and tendencies which have been considered at some length. Neurotic people are the victims of mental conflicts, and tend to suffer from much unconscious morbid guilt, yet, considering how numerous they are, the proportion of those who actually commit suicide is surprisingly small. Some psychiatrists will go so far as to deny that any neurotic will take such an extreme measure, but my experience leads me to regard such a statement as inaccurate and over-sanguine. It is also a highly popular fallacy to suppose that anyone who talks about suicide will not carry it out. Experience can only enable one to differentiate the individual who really means what he says and another who in a childish way through such a threat is endeavouring to get sympathy and attention and dominate some situation.

I think we can broadly say that it would be well for the doctor to look upon any patient who has a marked depression as a potential suicide. Depression may be a

symptom of many abnormal mental states, and it requires much psychiatric experience to be able to distinguish them. It is in the mental disorder of melancholia, or what we should more strictly term the depressive phase of manic-depressive insanity, that morbid depression is most marked and that suicide is always not only a possible but even a likely happening. This is especially so in that form of the disease which may show itself at the involuntional period of life. Melancholia is certainly largely constitutionally determined, but apart from that it seems essentially a reaction to emotional stress, particularly as it occurs in family situations. There are good reasons for believing that the sufferer has identified himself with some loved object, and that the hate which has followed love through disappointment and frustration is thus directed on to the ego. The mental torments he inflicts upon himself gratify sadistic instincts, and the feelings of aggression which were at first aimed externally are turned inwards against the person himself. As Oscar Wilde wrote in his *Ballad of Reading Gaol*, "All men kill the thing they love." The melancholic does this to the object he once loved and then hated, which object became part of himself. There is not only the wish to kill but also the wish to be killed. There is not only the strong sadistic impulse but the contrary urge of submissive masochism. The great sense of guilt that is necessarily aroused by the murderous wishes demands a self-punishment and a punishment that will fit the crime—the death penalty self-inflicted.

Suicide in Other Forms of Psychosis

Suicide is not infrequent in that form of psychosis which is apt to show itself in adolescence—namely, dementia praecox. In some acute cases the attempts are not uncommonly very desperate ones, and the methods used may be particularly grotesque. When the attempt is unsuccessful, and the patient is questioned as to the reason for it, it may well be that he can give none, or the motive alleged be trifling and absurd. In this type of mental disorder the provocation is often impossible to predict. In other types any delusions will probably be in accord with any self-destructive impulse, and hallucinatory voices not infrequently order patients to kill themselves.

In more than one form of psychosis delusions of persecution may be in the foreground, and it can be well understood how such a victim to them may, in the course of time, become tired of the perpetual struggle against his supposed aggressors and seek the refuge of death. So sufferers from senile mental disorders, paranoia, or general paralysis of the insane may in such circumstances end their lives. From the psychological aspect we must regard the taking of alcohol in excess as a defence against some form of unconscious mental conflict, so that it is not surprising that alcoholism frequently involves a suicidal urge. In many cases the psychosis alone does not cause the suicide, but to it is added some concrete crisis which constitutes the immediate motive. The idea of thus acting as a means of escape is as evident in these individuals who so deal with an imagined trouble as in the case of more normal people who face a real one.

Suicide in General Practice

In general practice all these varied types may be met with. Where there is frank evidence of a severe mental disorder which patently needs supervision and control in a mental institution the needful steps for certification will be taken. For more than one reason, however, the medical man is often loath to take this procedure. The recognition of a suicidal possibility is not so easily grasped. Because of such difficulties I have long advocated the appointment of a medical officer of mental health in every area, who could be called upon in a consultative capacity and countersign any certificate.

Some of the milder cases who have some insight into their abnormal state may co-operate in their treatment and become voluntary boarders at a hospital. Where volition is not present it is possible to get cases admitted as "temporary patients" without certification where there is likelihood that recovery is probable. The Mental Treatment Act of 1930 made such a step legal, in order that by non-certification any stigma might be obviated and thus early treatment be encouraged. Only too often do we hear of suicides having taken place which might have been prevented if only such precautionary measures had been taken. All concerned are very wise after the event.

It is not possible to lay down specific rules by means of which an attempt at self-destruction may always be foretold. Where certain types of psychoses are present such a happening is always possible, but especially so in melancholia and any severe depressive state. When we are consulted by a neurotic who shows indications of depression and a sense of guilt our sympathy and endeavour to understand his difficulties and feelings will go far in banishing the idea of suicide from his mind. The attachment to the "all-knowing" physician will constitute a human link that will bind him in some degree to this world. When upon physical examination no organic signs are found to account for his hypochondriacal or other worrying symptoms do not say, as is so often done, "There is nothing the matter." The patient feels that if he is thus misunderstood, as of course he is, no other help is available, with the result that death may beckon. In cases where no psychosis is present it may be true to say that the death call will be robbed of its main potency when there remains some hope and an understanding friend exists. The feeling that somebody cares is a valuable prop to lean on.

Insomnia, of course, is a usual prominent symptom in patients whose mental conflicts are of a depressing nature. From the first this should be combated, and in this respect do not be afraid of the giving of suitable sedatives until the mind has gained a greater measure of peace. We hear much of the dangers of hypnotic drugs, but psychiatric experience recognizes no reason for such apprehension when rationally administered.

Psychotherapy can certainly do a great deal to help the individual to overcome the ravages of past unfortunate emotional experiences, and to adapt more rationally and more happily to the world in which he lives. Only in this way can there be an adequate possibility of lessening or removing the morbid foundations of guilt, hatred, sadism, and revenge, which loom so largely in the suicidal background. Analytic measures, however, should be supplemented by synthetic ones. The individual who has gained insight into those emotional factors that have played havoc with his mental well-being and obviated his being "master of his fate and captain of his soul" must be led to adopt some philosophy of life which can perpetually act as a guiding force and a support in times of strain and stress. To some spiritual help may be conducive to that end.

In the prevention of suicide our most rational prophylactic aim must be to try to lay down a healthy emotional foundation for the mental soil, so that in later years the mind shall be enabled to adapt adequately to the inevitable periods of painful reality that in some degree must come to all. Parents must be better educated in the manifold factors related to the character development of their children, and especially so in the direction of their love life. When disharmonies are manifested in early life child-guidance clinics can be, if necessary, consulted. Certain childish characteristics and symptoms may, to the expert eye, be indications of possible suicidal tendencies in later life.