BRITISH MEDICAL JOURNAL

LONDON: SATURDAY, APRIL 25th, 1936

THE PROBLEM OF TOXIC GOITRE*

BY

JOHN MORLEY, CH.M., F.R.C.S.

HON. ASSISTANT SURGEON, MANCHESTER ROYAL INFIRMARY ; PROFESSOR OF SURGERY (ELECT),
MANCHESTER UNIVERSITY

(With Special Plate)

I propose in this paper to make some remarks on the problem of toxic goitre, but in truth this fascinating and elusive disease fairly bristles with problems, and there are few conditions, if any, with which surgeons are commonly concerned, about which we have to admit so much uncertainty. The treatment of the disease is at present by common consent mainly a matter for the surgeon, and the practical results of operative measures have so far outstripped our pathological knowledge that our operations stand in some danger of the charge of empiricism.

My observations are based upon a personal series of 785 cases of toxic goitre that I have operated on, with twenty-nine deaths—a mortality rate of 3.7 per cent. All the goitres have been examined histologically.

Classification

With many diseases, as our knowledge grows, our classification, both clinical and pathological, tends to become ever more complex. With toxic goitre this is not the case; there has been a marked tendency during the past few years towards a simplification, amounting almost to unification. At first sight there appears to be a clear distinction between the cases of primary exophthalmic goitre occurring in young women about the period of adolescence and the so-called toxic adenomatous goitres. In the latter type a middle-aged woman who has had a lump in her thyroid for years finds after some emotional strain or some illness, such as an attack of influenza, that her heart is beating faster and she is losing weight and strength. She has less of the peculiar restless irritability than the case of primary Graves's disease, and little tendency to exophthalmos, but though the tachycardia may not be extreme the cardiac effects are graver, and she soon shows a marked degree of auricular fibrillation.

Are these clinical types, which we call primary Graves's disease in the one case and toxic adenomatous goitre or secondary toxic goitre in the other, to be regarded as different or as essentially similar diseases? Plummer of the Mayo Clinic has maintained that they are really different diseases. Not only has he stressed the points on which they differ clinically, but he has stated that in primary exophthalmic goitre the thyroxine molecule is deficient in iodine, whereas in toxic adenoma the thyroxine molecule is not deficient in iodine, but is secreted in excess; there is, in fact, a true hyperthyroidism. He has

sought to support his biochemical views by the results of iodine therapy, and claims that whereas iodine causes a remission of symptoms in primary Graves's disease it makes a patient with toxic adenoma worse.

Much as we owe to Plummer for the introduction of iodine medication, it must be admitted that his chemistry is unsound, and his statement as to the harmful effect of iodine in toxic adenoma is not in accord with clinical experience. Since Harington in 1928 not only established the exact chemical composition of the thyroxine molecule, but succeeded in synthesizing thyroxine, we must abandon Plummer's conception that there is something qualitatively abnormal in the thyroid hormone in primary Graves's disease. Clinically we know that we can as a rule obtain just as marked a remission in toxic adenomatous goitre by the administration of iodine as in primary exophthalmic goitre.

Difficulties in Classification

The more one sees of patients suffering from hyperthyroidism the more difficult one finds it to separate them into clearly marked groups, either on a clinical or on a pathological basis, with any real confidence. There are, it is true, certain obvious and outstanding differences, as I have already indicated, between the primary exophthalmic goitre at one end of the scale and a toxic adenomatous goitre at the other. But between them we see a large number of cases showing clinical and pathological features that lead us by degrees from one type to another.

It might be thought that the presence of one or more adenomata in a goitre marks it out as something quite distinct from the diffuse hyperplastic goitre of primary Graves's disease. But it is coming to be recognized generally that these so-called adenomata, in the majority of cases, are not adenomata at all in the sense of benign new growths. They are localized areas of hypoplasia with increased colloid storage. In the strict pathological sense the term "adenoma" should be limited to the type of encapsuled growth, consisting of interlacing columns of cells with little or no colloid, which is known as the foetal adenoma (Fig. 1 on Special Plate). There is no doubt that many of the cases which come before us as nodular goitres began years ago as diffuse hyperplastic goitres associated with all the symptoms of primary Graves's disease. It is the recurring cycles of hyperplasia and involution which have gradually transformed them into nodular goitres. Others, again, have started as the diffuse colloid goitres of adolescence without toxic symptoms, and

* A paper delivered before the Sheffield Medico-Chirurgical Society on February 13th, 1936.

gradually in the course of years have developed one or more so-called colloid adenomata before the onset of any toxic symptoms. The prolonged administration of iodine will gradually transform a diffuse hyperplastic goitre into a nodular goitre, though the nodules are usually small and numerous, and scattered more or less uniformly through the enlarged gland. Yet other difficulties of clinical classification that must be faced are the large proportion of cases that have genuine primary hyperthyroidism without exophthalmos, and a certain smaller number that show marked exophthalmos and constitutional symptoms without any obvious goitre.

An Illustrative Case

I cannot do better than illustrate the difficulty of classification by a recent case.

A married woman, aged 37, was referred to me with the following story. There was no family history of goitre, and she had had no evidence of thyroid trouble until nine years ago, when, after the emotional shock of having a still-born baby, she developed a goitre with exophthalmos and the other classical symptoms of primary Graves's disease. A slow and spontaneous improvement followed, but five years later she found her father in a fainting condition and thought he was dead. This second shock caused a severe relapse; but again she slowly recovered. Six months before she came under my care her two-year-old child was accidentally, by the blowing-to of a door, locked into a room where there was a pan of boiling water on the floor. In terror she broke a window to save the child from being scalded, and since then all her old symptoms had returned. But her goitre when I saw her was not the original diffuse form; she had a large single adenoma in the left lobe pushing the trachea well to the right, and the right lobe was not clinically abnormal.

The conclusion to which a study of this and other somewhat similar cases must lead us is that all the varying clinical types with which we meet are essentially one disease, which you may call toxic goitre or hyperthyroidism as you will. The clinical and pathological differences between them would appear to be conditioned to a large extent, but not entirely, by the age at which the toxic symptoms appear.

As we must have some classification let it be as simple as possible. Following Dunhill, I believe these cases are best classified as primary and secondary toxic goitres.

CLASSIFICATION OF TOXIC GOITRES

Primary.—(1) Early. (2) Late.

Secondary.—(1) Early. (2) Late.

By *primary* I mean that the onset of the toxic symptoms and the appearance of the goitre coincide more or less closely in time. By *secondary* I mean that the goitre precedes the toxic symptoms by a considerable time, and usually by several years. Each of these groups, the primary and the secondary, may conveniently be divided into two subdivisions, the early and the late. Early primary cases usually begin about adolescence. Late primary cases arise most commonly in women about the time of the menopause. Early secondary cases are those in which a simple diffuse colloid goitre of adolescence persists, and the toxic symptoms arise a few years later. Late secondary cases are those which are so commonly called toxic adenomatous goitre, where a simple nodular goitre with either one or many "adenomata" is present for many years before the onset of the toxic symptoms.

From a strictly pathological point of view any classification breaks down on the fact that microscopical sections will so often show areas of both hyperplasia and colloid adenomata in the same gland, and even inside the capsules of colloid adenomata we find areas of hyperplasia.

Histological Considerations

It is convenient at this stage to consider the various histological pictures and some of the problems that they suggest. To show the typical appearance of the primary toxic goitre it is necessary to go back to the material examined before the introduction of pre-operative iodine therapy (Fig. 2). We notice the almost complete absence of colloid storage, the resulting empty alveoli, and the columnar shape of the cells lining them.

The effect of the iodine is to increase the colloid storage (Fig. 3). Clinically it causes a pronounced remission of symptoms, shown by a drop in the pulse rate and the basal metabolic rate, with a rapid improvement in the nervous phenomena. This general improvement begins in two or three days, and reaches its maximum as a rule in from ten days to a fortnight. After that there is usually a definite though gradual tendency to relapse, whether the iodine is continued or stopped. If, as so often happens, iodine has been administered for months before, a remission cannot be obtained by its use unless it is discontinued for some weeks, and I believe that these later remissions are never so complete or so favourable as the first one, when the iodine is acting on virgin soil. These are the reasons why surgeons plead, too often as a voice crying in the wilderness, that iodine should be regarded, except in grave emergency such as maniacal attacks or severe vomiting, not as an agent for medical treatment, but merely as a temporary expedient to make operation safe. It is, of course, beyond question that the use of iodine has done far more than any other measure to bring down the operative mortality in toxic goitre.

I wish, in this connexion, to emphasize the inadvisability of the routine use of large doses of iodine as practised by some authorities. Adequate iodization of these patients can be assured by giving ten minims of Lugol's 5 per cent. iodine solution daily; the use of massive doses does involve danger of a severe exacerbation of the disease.

Significance of Colloid

The student is apt to think of the colloid secretion as the active principle of the thyroid gland. But when we see it as colloid the secretion is not in its active state but in a resting stage. It is stored up in the alveoli and inert, and has to undergo some change, the nature of which is unknown, before it can be reabsorbed by the alveolar cells and passed into the blood stream as active thyroxine. In the most active hyperplastic glands there is least colloid secretion visible. Such a gland contains less iodine than normal, while an excess of iodine is found in the blood and urine. The not inappropriate term thyroxine "diarrhoea" has been suggested for this condition.

A period of involution, during which the toxic symptoms become less evident, may coincide with an actual increase in the nodular swellings in the gland, and to this state we might give the term "constipation of thyroxine." It is to the active cellular hyperplasia that we must look for the main histological evidence of hyperthyroidism.

When we examine a series of sections taken from patients with toxic goitre we are at once perplexed by the variety of changes. In the same gland and often in a single section we find areas of hyperplasia, areas of involution and fibrosis, and so-called colloid adenomata. It is extremely difficult, if not impossible, particularly since the introduction of iodine therapy, to correlate the histological findings with the clinical symptoms.

Nature of So-called "Vacuoles"

I must draw attention to a very common appearance in the follicles of toxic goitre. We see in many sections a peculiar vacuolated appearance at the periphery of the alveoli between the colloid and the lining epithelium (Figs. 4 and 5). In contrast with the pink staining of the colloid these rounded areas are clear and colourless. Some authorities have regarded them as artefacts due to shrinkage of the colloid during fixation and staining, but Professor S. L. Baker, who has kindly prepared some special sections to investigate this point, assures me that they are certainly not artefacts. These "vacuoles," as he describes them, are considered by Boyd to be due to absorption of the colloid by the epithelial cells with which it is in contact. But if such absorption took place I think we should see merely a relative diminution of the colloid contents and a shrinking in size of the gland as a whole, and not a series of empty spaces.

It appears to me that they are not vacuoles at all, but droplets of recently poured out secretion which is in an active, non-staining state, and is no doubt capable of rapid absorption through the epithelial cells into the blood stream. I suggest that the emotional disturbance due to preparation for the operation may be in part the cause of this excessive secretion. It is not uncommon to observe a perceptible increase in the size and rotundity of the goitre on the morning of operation as compared with the preceding days. This pre-operative swelling of the gland was more noticeable before pre-operative iodine medication was introduced.

The interlobular fibrosis, which is a marked feature of the nodular goitres and of the old-standing cases of primary Graves's disease, is presumably developed in the various cycles of involution of the gland, and may be an attempt on the part of Nature to overcome the excessive vascularity that we see in the early primary toxic goitre. This fibrosis is responsible to a great extent for the peculiar hardness of an old-standing diffuse toxic goitre.

In some toxic goitres conglomerations of lymphocytes are a marked feature, and in many we see definite lymphoid follicles with typical germinal centres scattered throughout the gland (Fig. 6). Although a lymphadenoid goitre is often said to be associated with myxoedema all the specimens I have studied, and there are very many in my collection, have shown marked hyperthyroid symptoms. The function of this lymphoid tissue in toxic goitre is not in the least understood, and should be a fruitful subject for research.

Problem of Pathogenesis

Consider first this question. Are the various clinical phenomena of Graves's disease the result of an excessive output of a normal secretion—a true hyperthyroidism, or of the formation of an abnormal secretion—a dysthyroidism as it has been called? I am aware that there is a great weight of authority in favour of the theory of dysthyroidism. But there are certain considerations that seem to me to point strongly to the theory of a simple excess of the active principle. In the first place, as we have seen, the exact analysis of the thyroxine molecule by Harington, who has established its structural formula and has brought it to the proof of synthesis, makes it hard to conceive that the gland can have a secretion of variable composition.

We know, too, that the symptoms of Graves's disease can be induced by an excessive dosage of iodine or of thyroid extract—what is known on the Continent as "iodine Basedow's disease." I have seen several cases in which toxic symptoms were quite definitely due to the administration of thyroid extract in considerable doses

to patients with simple goitre, under the mistaken belief that the goitres were an expression (as some undoubtedly are) of iodine or thyroxine insufficiency.

Then again, as we note the wonderful effect on the disease of the surgical removal of a sufficiently large proportion of the gland and see how too drastic a removal will cause, though rarely, post-operative myxoedema, it is difficult to resist the conviction that the disease is associated with a simple excess of thyroxine. The analogy of the other glands, such as the pituitary, the parathyroids, the islands of Langerhans, and the adrenals, points to the same conclusion, for we think of all their disorders as due to simple excess or deficiency of their internal secretion.

Whatever may prove to be the ultimate truth of the matter I feel that in the present state of our knowledge we confuse our thought unduly by entertaining the theory of perverted secretion, and I believe that the theory of excessive secretion provides a safer and better working hypothesis to explain the phenomena of toxic goitre.

Predisposing Causes

Age, sex, and heredity all play their part as predisposing causes. As regards age, early primary toxic goitre is essentially a disease of adolescence. It is not uncommon, however, to meet with it in a severe form in quite young girls, and I have had occasion to operate on several cases under the age of puberty, the youngest being 9 years of age. There is a late primary form, occurring, as it seems to me, with increasing frequency, which shows itself about the period of the menopause.

Early secondary toxic goitre—a relatively uncommon form—occurs a few years after the onset of a simple colloid goitre of adolescence, usually between the ages of 25 and 35. The commoner late secondary type is met with mostly after the age of 30, and often in quite elderly patients. Primary toxic goitre in men tends to arise notably later in life than in women, a fact no doubt connected with the greater immunity to the disease shown by the male. As regards sex, it is generally agreed that about ten cases occur in females to every one in the male. My own series contains approximately eight females to one male.

It is difficult to assess the importance of heredity as a predisposing factor, but in a large number of cases there is a history of other goitres in the family, either toxic or non-toxic, and one cannot resist the conclusion that many of these patients have an inherited instability of their nervous mechanism, or thyroid apparatus, or both. A case illustrating this point and brought to my notice by Dr. A. Hillyard Holmes was reported by Dr. Dreschfeld¹ in 1896.

A female child, aged 2½ years, was startled by a large black dog. From that time she screamed and talked in her sleep and became very quiet and listless when awake. Soon after this her mother noticed emaciation and a heaving action of the heart, and she developed all the symptoms of primary exophthalmic goitre. She recovered spontaneously by the age of 7, grew up and married, and her daughter, at the age of 14, came under Dr. Holmes's care with exophthalmic goitre.

Exciting Causes

Given the hyperthyroid tendency there is still some exciting cause needed to produce the state of toxic goitre, and we find that this exciting cause very commonly takes one of two forms. It may be either emotional trauma or bacterial infection. Emotional trauma is undoubtedly the commonest exciting cause in primary toxic goitre. The large black dog of Dr. Dreschfeld's case is a typical instance. Let me mention some other examples. The sudden death of a near relative is a very common

precipitating event, as is the news of an accident to a relative. In one of my cases the doctor told a mother whose child was ill that he had no hope of the child's recovery. The child recovered, and the mother promptly developed Graves's disease.

Sexual offences or attempts at sexual offences against girls provide a not uncommon cause, particularly where the emotional disturbance of giving evidence in court is added. One patient said that her symptoms started with the news that her sister had had an illegitimate child. Sometimes the disease follows an operation, such as hysterectomy, appendicectomy, or complete removal of the teeth. Motor accidents provide a very frequent cause. One could elaborate the list almost indefinitely, but it is worth noting that in most of the cases I have mentioned the shock was a sudden one, and was not repeated.

Less commonly we find a history of the more prolonged emotional strain of a pregnancy, or of nursing a sick child or relative in a long illness, or in young girls the strain of working for examinations and failing in them. Examinations in music seem particularly prone to cause such troubles, but not examinations in medicine. An undue proportion of cases are found in school mistresses, theirs being an occupation that involves much nervous strain. In men it is the prolonged strain of business worries or anxiety about employment that causes trouble rather than any sudden shock, and it will be recalled that we did not see many cases among the shell-shocked during the war, though there were undoubtedly a few.

Among bacterial infections influenza comes easily first, and the epidemic of 1933 has been providing surgeons with thyroidectomies ever since; less common illnesses, but still not infrequent, are recurrent acute tonsillitis and acute rheumatism. All these infective conditions seem specially prone to produce toxic symptoms in patients with old-standing simple nodular goitres, though we occasionally see them producing primary Graves's disease.

Though diet has a proved and important place in the aetiology of simple goitre, there is little evidence that it counts greatly in toxic goitre. Dr. Kletz and others have stressed the importance of a diet rich in carbohydrates in pre-operative treatment, in order to restore the depleted reserves of glycogen in the liver.

Pituitary Relationships

To enumerate the exciting causes of the disease plainly brings us no intellectual satisfaction. We must inquire how they act. I fear that in the present state of our knowledge we can only inquire and wait patiently for the final answer. What is the part played by the central nervous system, by the sympathetic nervous system, or by the pituitary gland in the causation of the disease? It is plain that the experimental physiologists have the task of answering these more fundamental questions, and it is to the results of their labours that we must turn.

Let us first consider the evidence of a close relationship between the pituitary and the thyroid gland. It has been established experimentally in a number of animals that extirpation of the pituitary leads to some degree of atrophy of the thyroid, and, conversely, that continued injection of anterior lobe extracts of the pituitary causes hyperplasia of the thyroid. It is also known that thyroidectomy in animals causes hypertrophy of the anterior lobe of the pituitary.

Some interesting experiments have been recorded by Schockaert² on the duck. He found that daily injection of bovine anterior pituitary extract into young male ducks caused hypertrophy of the testes and increase in the size of the thyroid. After three weeks the thyroid reached thirty times the size of normal control glands; and histologically the gland developed features closely resembling primary toxic goitre. When the injections had

continued for more than three weeks the birds showed exophthalmos, loss of down, and increase in the weight of the heart. These effects are attributable to the thyrotropic hormone which has been isolated from the anterior pituitary.

Thyrotropic Pituitary Hormone

Collip and Anderson,³ working with this thyrotropic hormone, have shown recently that daily injections into rats cause hyperplasia of the thyroid with an increased basal metabolic rate. This effect did not persist, however, and after the thirty-seventh day the basal metabolic rate was greatly below the normal, and lower even than that of hypophysectomized rats. When the serum of these rats, which had been thus rendered immune or resistant to the hormone, was injected into hypophysectomized rats, these latter animals were found to show none of the hyperplastic thyroid reaction or increased basal metabolic rate which the thyrotropic hormone normally induces in them. Similar results were obtained in the guinea-pig. Collip suggests that this opens up the possibility of a serum treatment of hyperthyroidism.

Marine⁴ has shown that the thyrotropic hormone is independent of nerve impulses, since it stimulates thyroid transplants in any part of the body to approximately the same degree that the non-transplanted thyroid is stimulated. Recently Eitel, Krebs, and Loeser⁵ have shown that the thyrotropic hormone stimulates thyroid cells *in vitro*.

While the great importance of the thyrotropic hormone of the anterior lobe of the pituitary as a cause of thyroid hyperplasia has received increasing attention of late, there is some evidence that the thyroid may be stimulated to over-activity by a purely nervous mechanism, and this work has hardly received the attention it deserves. In 1915 Cannon, Binger, and Fitz⁶ showed that in cats suture of the proximal end of the divided phrenic nerve to the distal end of the cervical sympathetic produced exophthalmos, increased basal metabolic rate, diarrhoea, and an irregular pulse—all the essential phenomena, in fact, of hyperthyroidism. This ingenious experiment, though proving beyond doubt that the thyroid secretion is to some extent under the control of the sympathetic nervous system, cannot be said to solve the riddle of toxic goitre, for there is no reason to believe that a continuous rhythmical discharge of stimuli, such as this phrenic anastomosis provides, ever passes down the sympathetic nerves to the thyroid in cases of toxic goitre in man.

Hypothalamic Relationships

The fundamental problem of toxic goitre seems to me to be this: How are we to explain the mechanism by which a transient emotional shock, or a brief bacterial invasion, sets up a state of hyperactivity in the thyroid which often lasts for years? Some kind of vicious circle must be postulated in the mechanism that brings this about—but what? It is here that experimental work on the hypothalamus brings us an uncertain glimmer of light.

The condition of "sham rage" manifested by the decorticated dog in Goltz's classical experiment has been investigated by Bard,⁷ and the reactions constituting this condition have been proved to depend on a centre in the hypothalamus. This state of "sham rage" is associated with raised blood pressure, increase in blood sugar, sweating, and outpouring of adrenaline. This, however, is only a part of the evidence which tells us that there is in this region of the brain a higher centre for the sympathetic system. Recent research has shown that this small area of brain under the third ventricle controls the primary emotions, or perhaps we should say the expression of those emotions, the metabolic processes of the body, and the

sympathetic nervous system. It is also, as Cushing has shown, in very close relationship with the pituitary body. The hypothalamus, according to Cushing, by its close neural connexion with the pituitary, is capable of stimulating that organ to activity through the agency of nerve fibres running down the pituitary stalk. We must not yet regard this hypothalamus-pituitary relationship as certainly proved, but it is a theory which grows in probability with increasing knowledge, and Langdon-Brown⁸ has recently summarized the growing evidence in its favour.

A Theory of Pathogenesis

Let me be bold enough to attempt a theory of the pathogenesis of toxic goitre. I would suggest that an

A diagrammatic sketch of the "vicious circle" theory of toxic goitre. A = Hypothalamus. B = Pituitary. C = Sympathetic system. D = Thyroid. E = Adrenals.

emotional trauma, or bacterial infection acting on a hypothalamus that is rendered peculiarly susceptible by hereditary and possibly by dietetic influences, stimulates that hypothalamus to over-activity. The hypothalamus in its turn stimulates the pituitary by nervous pathways. From the anterior lobe of the pituitary the thyrotropic hormone is poured out in excess, and doubtless the adrenotropic hormone also, and, passing through the circulation, initiates a process of active hyperplasia in the thyroid. Thyroxine normal in quality and excessive in quantity is

poured out into the blood, and produces its symptoms by some action on the sympathetic system.

Whether the thyroxine acts centrally on the sympathetic centre in the hypothalamus, or peripherally on the sympathetic nerve terminals, or both, seems quite uncertain, though the occasional occurrence of acute mania in toxic goitre and the constant emotional instability of these patients provide suggestive evidence of an action on the centres in the hypothalamus. But it would appear quite plain that all the cardinal symptoms of the disease—exophthalmos, tachycardia, sweating, tremors, and the increased basal metabolism—are due to the over-stimulation by thyroxine of the sympathetic system. In this over-stimulation adrenaline would seem to play an essential part, and it may be that there is a closer connexion than has hitherto been believed between the thyrotropic and the adrenotropic hormone of the pituitary. This theory has the merit of suggesting how an initial emotional shock or bacterial infection may set up a vicious circle of nervous and endocrine disturbances which keeps the disease in activity, and which it is the object of treatment to break.

Exophthalmos

Some recent experimental work suggests that while thyroxine alone is unable to produce exophthalmos in animals it can do so when given in combination with drugs, such as adrenaline or ephedrine, which are themselves capable of stimulating the sympathetic system.⁹ Further it has been shown that this experimental exophthalmos is produced with greater ease after thyroidectomy—a fact which suggests that the thyrotropic and/or the adrenotropic hormone of the pituitary may be of greater importance in its production than thyroxine itself.

I must here elaborate my reasons for including exophthalmos among the results of sympathetic stimulation. The exophthalmos has been attributed by some, including no less an authority than Dr. G. R. Murray, to increased intra-orbital fat, but the post-mortem evidence for this theory has never been convincing, and it seems improbable that a wasting disease could cause such a local excess of fat.

There are, however, certain arguments that point directly to a muscular origin of the exophthalmos. It will sometimes, though rarely, develop in a few hours after a sudden shock. This could only happen by a neuro-muscular mechanism. Further, the opposite condition, enophthalmos, is caused, as we all know, by division of the cervical sympathetic—a lesion that we often saw during the war, and now produce deliberately when we do a cervical sympathectomy. This paralytic enophthalmos is convincing evidence that the position of the eyeball in the bony orbit is dependent on muscular tone.

Jaboulay's operation of division of the cervical sympathetic in the neck has long become obsolete as a means of treating Graves's disease because it does nothing to reduce the basal metabolic rate and other constitutional symptoms, but it still has some value in the rare cases with exophthalmos so extreme as to threaten ulceration of the cornea.

If we conclude, as I think we must, that the exophthalmos is due to a muscular mechanism innervated by the cervical sympathetic, we must next attempt to indicate the precise muscles involved. Without stopping to consider the various muscles which have been held responsible, I must briefly state my own view. The check ligaments are arranged in continuity with the capsule of Tenon. The check ligaments, as Sappey showed long ago, are chiefly composed of smooth muscle fibres, and the capsule of Tenon, though largely fibrous, can be shown by differential staining to contain a considerable proportion of smooth muscle. These structures, the check ligaments and the capsule of Tenon, combine to form a sort of catapult, which when it contracts projects the eyeball forward. The retraction of the upper lid, which is a separate phenomenon from the exophthalmos, can be explained by stimulation of the smooth muscle element in the levator palpebrae superioris.

If I am asked why it is that exophthalmos, after a successful operation for Graves's disease, so often persists when all other symptoms have vanished, I should attribute it to the organic shortening which this muscular sling, like any other muscle in prolonged spasm, has undergone.

Clinical Problems

It would be wrong to leave this imperfect attempt to elucidate the mechanism of exophthalmos without admitting that the view I have attempted to set forth is still largely hypothetical, and further investigations may well lead to some modification of it. No theory of exophthalmos can be regarded as entirely satisfactory when it fails to account for the occasional occurrence of a strictly unilateral exophthalmos in toxic goitre. The explanation of this phenomenon remains a complete mystery.

There are two clinical problems connected with toxic goitre to which I can refer only briefly. A considerable proportion of these patients, and particularly the older ones, develop some degree of chronic arthritis which they describe as "rheumatism." The arthritis usually begins before operation, and sometimes continues unchecked or even becomes worse after operation. The second problem is that many of them suffer from a type of abdominal pain that sometimes simulates the pain of peptic ulcer. I believe that both these conditions offer a promising

field for systematic clinical research, but it is unlikely that we shall arrive at a satisfactory understanding of them until more is known concerning the influence of the sympathetic system upon the joints and upon the mechanism of visceral pain.

Conclusion

I am conscious that these remarks have raised various problems and have done little to solve them, but it is perhaps well that we should realize what gaps still remain in our knowledge of this most fascinating disease. On the question of operative treatment, with which I have not attempted to deal, it would be unwise to assume that we have reached finality. Although the surgical treatment of toxic goitre can at present claim such results that it is regarded as the method of choice for all save the mildest cases, I never do a subtotal thyroidectomy without feel-

ing what a crude and imperfect method it is. Who can doubt that before many years have passed the biochemist will place some weapon in the physician's hand that will enable him to break the vicious circle of toxic goitre as certainly as the surgeon can, and without any of the risk which we cannot, try as we may, altogether eliminate from these anxious and delicate operations.

REFERENCES

- ¹ Dreschfeld, J.: *Practitioner*, 1896, iv, N.S., 135.
- ² Schockaert: *Amer. Journ. Anat.*, 1932, xlix, 379.
- ³ Collip, J. B., and Anderson, E. M.: *Lancet*, 1934, i, 76.
- ⁴ Marine, D.: *Journ. Amer. Med. Assoc.*, 1935, civ, 2253.
- ⁵ Eitel, H., Krebs, H. A., and Loeser, A.: *Klin. Woch.*, 1933, xii, 615.
- ⁶ Cannon, Binger, and Fitz: *Amer. Journ. Physiol.*, 1915, xxxvi, 363.
- ⁷ Bard: *Ibid.*, 1928, lxxxiv, 490.
- ⁸ Langdon-Brown, W.: *Lancet*, 1935, ii, 1155.
- ⁹ Brain, W. Russell: *Ibid.*, 1936, i, 182.

DIAGNOSIS OF PHYSICAL DISORDERS IN THE INSANE

BY

RUBY O. STERN, M.D.

PATHOLOGIST TO ST. ANDREW'S HOSPITAL, NORTHAMPTON

J. McLEMAN AND B. F. M. BOND

M.B., Ch.B.

M.B., B.S.

ASSISTANT MEDICAL OFFICERS TO THE HOSPITAL

(With Special Plate)

The well-recognized fact that psychotic patients may suffer from serious organic disease without uttering a word of complaint by no means constitutes the chief difficulty in the diagnosis of physical disease in the insane. As Hall¹ has pointed out, physical signs may be few, or even absent, in grave medical or surgical conditions which in the mentally normal give rise to unmistakable clinical pictures. If this be true in such common disorders as appendicitis and gastric ulcer, to take only two examples, it is unlikely that rare diseases will be diagnosed in the insane during life, especially as the use of accessory methods of investigation may not be possible owing to lack of co-operation or active resistance of the patient, and even a complete physical examination may have to be abandoned.

During the past year four patients came under observation at this hospital in whom the physical diagnosis was doubtful. As the conditions revealed at necropsy were somewhat unusual these cases were considered of sufficient interest to record.

Case I

A man aged 82 had been insane since 1888. He was transferred to this hospital in 1929 in a state of secondary dementia. He then had arterial and myocardial degeneration, but there were no signs of active heart disease. For the last two years of his life he had been bed-ridden and so demented that he was unable to notice any change in his physical well-being. The terminal illness began on July 15th, 1934, with a rigor, which was followed by high fever. The heart sounds became almost inaudible, and crepitations appeared at the bases of the lungs. He gradually became unconscious, and died on July 20th without developing any further signs. There was, however, some diminution in the quantity of urine passed during the last few days of life. Examination of the urine the day before death showed a large amount of albumin, many granular casts, a few pus cells, and many coliform bacilli.

A necropsy was performed on the day of death, when it was found that the heart was enlarged, weighing 14 oz. The muscle was red and firm, and the left ventricle was hyper-

trophied. On the left cusp of the aortic valve there was a mass of vegetations about the size of a sixpence. This mass was reddish in colour, and, though not easily friable, neither was it hard and fibrotic. It ulcerated the margin of the valve. No other vegetations were present. There was only slight atheroma of the cusps of the aortic and mitral valves and of the proximal part of the thoracic aorta. The coronary arteries were patent throughout.

Both kidneys presented a "flea-bitten" appearance, due to multiple petechial haemorrhages scattered over their surfaces. On section many more tiny haemorrhages were seen in the cortex. In the centre of some of these were tiny white spots which exuded pus on pressure. The cortex of both kidneys was greatly diminished, and the cortical markings were indistinct. In the right lung many alveoli were dilated, giving a bullous appearance. On section a little clear, frothy fluid was expressed. The left lung was well aerated throughout. The brain and spinal cord were normal to naked-eye examination, and nothing abnormal was noted in the other organs, except in the spleen, which was unduly soft and friable. No infective focus other than the vegetation on the aortic valve could be traced.

Microscopical examination revealed an area of acute inflammation in this vegetation, in which polymorphonuclear leucocytes and diplococci could be seen. Miliary abscesses, also of recent origin, were present in the cortex of both kidneys. During routine examination of sections of the brain many miliary abscesses were found around the vessels in the cerebral cortex, both in the grey and in the white matter (Fig. 1 on Special Plate).

The unusual features of this case were the age of the patient, the absence of signs of active pre-existing heart disease, and the short duration of the terminal illness. It is probable that the endocarditis was due to a pneumococcal infection, since this form of bacterial endocarditis is more common in later adult life, is very rapidly fatal, and, according to Thayer,² in only 37.5 per cent. of cases is there a history of pre-existing heart disease. Focal areas of suppuration, such as we found, are common in pneumococcal endocarditis—Thayer found them in 72 per cent. of his cases—while they are rare in any other form.

Case II

A man aged 74 was admitted on December 5th, 1934, in a condition of arteriopathic dementia. He had advanced generalized arteriosclerosis. His blood pressure was 180/122. The urine contained a trace of albumin but no casts, while the blood urea was 85 mg. per cent. He had incontinence of urine, not suggestive of retention overflow, but attributable to his mental state. There was a constant purulent nasal discharge, which on examination was found to be due to large nasal polypi. An operation for their removal was considered necessary in spite of his poor physical state, and this was performed on December 20th under sodium evipan anaes-

JOHN MORLEY: THE PROBLEM OF TOXIC GOITRE

FIG. 1.—Foetal adenoma of thyroid (high power). Interlacing columns of cells with relatively little colloid.

FIG. 2.—Early primary toxic goitre untreated by iodine. Almost complete absence of colloid storage with empty alveoli lined by columnar cells.

FIG. 3.—Early primary toxic goitre modified by iodine preparation. Note the increased colloid storage.

FIG. 4.—Primary toxic goitre, showing so-called "vacuoles" at the periphery of the alveoli between the colloid and the lining epithelium.

FIG. 5.—Secondary toxic goitre, showing so-called "vacuoles" lying again between the colloid and lining epithelium of the alveoli.

FIG. 6.—Primary toxic goitre with lymph follicles, showing germinal centres.