A NEW ARSENICAL FOR THE TREATMENT
OF SYPHILIS AND TRYPARSOMIASIS

BY

WARRINGTON YORKE, M.D., F.R.C.P., F.R.S.

AND

F. MURGATROYD, M.D., M.R.C.P., D.T.M.

In Collaboration with

F. GLYN-HUGHES, M.B., CHB.,

H. M. O. LESTER, M.R.C.S., L.R.C.P., D.T.M. & H.

AND

A. O. F. ROSS, M.D., D.P.H.

During the last seven or eight years we have examined, for the Chemotherapy Committee of the Medical Research Council, the trypanocidal activity of numerous new compounds, including a long series of aromatic arsenicals of the general formula—

prepared in Professor Morgan's laboratory of the Department of Scientific and Industrial Research at Teddington. The initial steps in this investigation consisted in examining the trypanocidal activity of the compounds in mice infected with a standard trypanosomal infection, the response of which to old and well-tried drugs, such as atoxyl and tryparsamide, had been thoroughly investigated. For practical purposes we selected tryparsamide with the object of comparison because it is probably the most widely used of the aromatic arsenicals in the treatment of trypanosomiasis, and is also employed on a considerable scale in the treatment of neurosyphilis.

As a result of prolonged investigation we discovered that about half a dozen of these new compounds had in trypanosome-infected mice a therapeutic activity at least equal to that of tryparsamide. After conferring with Professor Morgan regarding the probable ease and relative cost of manufacturing these compounds on a commercial scale, we selected one of the most active for more extensive trial on larger animals, and eventually on man. The compound selected was sodium succinamido-methylamide-*p*-arsonate—

it is referred to as *Neocryl** in the present paper. It is a white crystalline substance readily soluble in water, and its mode of preparation has already been described by Morgan and Walton (1931).

Experiments on Mice

The drug was administered to mice intraperitoneally, each dose being dissolved in 0.5 c.cm. of sterile distilled water.

Toxicity.—Neocryl is, like tryparsamide, tolerated by mice in relatively large doses. The maximum well-tolerated dose of tryparsamide for mice is about 40 to 50 mg. per 20-gram mouse, and the new compound is well borne in such doses. In order to obtain some idea of the relative toxicities of neocryl and tryparsamide thirty mice were each given 100 mg. per 20-gram mouse of tryparsamide, and thirty mice the same dose of neocryl. Of the thirty which were given tryparsamide seventeen (57 per cent.) died in seven days, and of those given neocryl only five (17 per cent.) died in a similar period. It appears, therefore, that neocryl is definitely less toxic for mice than is tryparsamide.

* This product will be put on the market as "Crylarsan Brand of Neocryl."

Therapeutic Action.—The minimum effective dose (M.E.D.) of neocryl for mice infected with our laboratory strain of *T. rhodesiense* is about 6 mg. per 20-gram mouse, and the minimum curative dose (M.C.D.) about 12.5 mg. per 20-gram mouse. The M.E.D. of tryparsamide for mice infected with this strain was found by us (Yorke, Murgatroyd, and Hawking, 1931) to be 6 mg. per 20-gram mouse, and the M.C.D. about 20 mg. per 20-gram mouse. These results for tryparsamide agree closely with those recently published by Strangeways (1935), who, working with mice infected with our strain of *T. rhodesiense*, also found the M.C.D. of tryparsamide to be 20 mg.

Experiments on Rabbits

The drug was administered to rabbits intravenously in a 20 per cent. solution in sterile distilled water.

Toxicity.—Nine rabbits were given a single dose of neocryl of 1.5 grams per kilo of body weight. Four of the animals tolerated the dose well, but the others showed toxic signs and died within ten days. Eleven animals were then given 1 gram per kilo of body weight; of these seven remained perfectly well, but the remaining four lost weight and died within three weeks; smaller doses were well borne by all rabbits. These figures are comparable with those obtained with tryparsamide.

Therapeutic Activity.—A series of rabbits infected with our own laboratory strain of *T. rhodesiense* was treated by various courses of neocryl. Treatment was commenced about the end of the third week of the infection, when the animals showed well-marked signs of disease, such as subcutaneous oedemas, patches of indurated and ulcerated skin, ulceration of the nares, and genital lesions; trypanosomes were present in the blood, usually in small numbers, and in considerable numbers in the various lesions. The rabbits were kept under observation for at least three months after cessation of treatment.

TABLE I.—Showing the Results of Treating Rabbits Infected with *T. rhodesiense* with Single and Repeated Doses of Neocryl and Tryparsamide Respectively

Method of Treatment and Dose in Gram per Kilogram of Body Weight	Neocryl		Tryparsamide	
	Number Treated	Number Cured	Number Treated	Number Cured
0.25 gram, single dose	4	0	3	0
0.5 gram, " " " " " " " "	5	4	5	3
0.125 gram, 4 doses at three-day intervals	5	1	5	1
0.25 gram, " " " " " " " "	9	9	12	9
0.4 gram, " " " " " " " "	5	5	6	6

From this table it can be seen that with neocryl a single dose of 0.5 gram per kilo body weight sufficed to cure four out of five infected rabbits, while all animals receiving four doses of 0.25 gram or more per kilo body weight at three-day intervals were cured. Comparison of these results with those obtained with tryparsamide and the observations on the relative toxicities of the two drugs suggest that neocryl is somewhat more effective than is tryparsamide in the treatment of rabbits suffering from trypanosomiasis.

In vitro Observations

In a previous paper (Yorke and Murgatroyd, 1930) it was shown that our normal strain of *T. rhodesiense* maintained in nutrient medium (equal parts of de-activated rabbit serum and Ringer-glucose solution) at 37° C. was completely destroyed within twenty-four hours by a concentration of 1 in 1,600 tryparsamide, and by a concentration of only 1 in 100 millions of reduced tryparsamide thioglycollate. The corresponding values of

neocryl and its reduced homologue are 1 in 6,400 and 1 in 100 millions respectively. As was to be expected from our earlier observations on drug resistance (Yorke, Murgatroyd, and Hawking, 1931) neocryl was found to exert as little action on arsenic-fast strains of trypanosomes as did tryparsamide or the other aromatic compounds of arsenic.

Conclusions from Experimental Work

All this preliminary experimental work suggests that there is a close resemblance between the therapeutic action of neocryl and tryparsamide. Both drugs are of relatively low toxicity, and are tolerated in large doses by mice and rabbits. Judged from experiments on mice, neocryl appears to be even less toxic than tryparsamide, since 83 per cent. of mice tolerated a dose of 100 mg. of the former, whilst only 43 per cent. of mice survived a similar dose of the latter.

In both infected mice and infected rabbits neocryl proved to have a somewhat greater trypanocidal activity than tryparsamide. The minimum curative dose of neocryl for infected mice was 12.5 mg. per 20-gram mouse, as compared with 20 mg. in the case of tryparsamide; a short course of neocryl, consisting of four doses of 0.25 gram per kilo body weight, given at three-day intervals to infected rabbits, sufficed to cure all of nine animals, whereas a similar course of tryparsamide cured nine of twelve rabbits. Having reached the conclusion from this experimental work that neocryl is rather less toxic and rather more active than tryparsamide, we felt that we had established a *prima facie* case for testing the drug in man.

Trial on Man Infected with Syphilis or Trypanosomiasis

Experiments on animals having shown that neocryl is if anything less toxic than is tryparsamide, it was decided in the initial trials on man to give the drug in doses similar to those employed for tryparsamide—namely, in amounts of 2 to 4 grams per week. Experience soon showed that the drug was easily tolerated in these weekly doses, and that they could be continued for prolonged periods without producing toxic signs. Even advanced

cases of neurosyphilis appeared to tolerate the drug well, and none developed signs of visual disturbance, although among similar cases treated with tryparsamide ocular troubles were fairly frequent. Apart from very occasional nausea and vomiting immediately following an injection, the only toxic signs observed in our series of cases were mild arsenical dermatitis in two patients, and jaundice, which subsequently disappeared, in two or three very advanced cases of neurosyphilis after prolonged courses of the drug.

Neocryl was administered to man by the intravenous injection of a 15 to 20 per cent. solution in sterile distilled water. The drug has been tested on patients suffering from primary and secondary syphilis, tertiary syphilis, and from various forms of neurosyphilis; it has also been tried on a number of cases of human trypanosomiasis in Nigeria.

Primary and Secondary Syphilis

It seems to be commonly agreed that tryparsamide is of little use in the treatment of any stage of syphilis except neurosyphilis. Brown and Pearce (1924) write: "A few patients with active primary or secondary lesions were also treated, but in these the lesions were either unaffected or their activity was increased to such an extent that, for the time being, the use of the drug was discontinued in patients showing such lesions." Reese (1933) states that "Tryparsamide has a very feeble spirochaetidal activity. It is inert in the first and second stages of syphilis, in syphilitic aortitis, in tabetic arthropathies, and in tertiary manifestations in general, which precludes its therapeutic use in these types." Moore (1933) in his recent work on the modern treatment of syphilis records the opinion that tryparsamide "is of no value in the treatment of early syphilis," and that it "has practically no effect in the healing of lesions in early syphilis in double the average therapeutic dose."

In Table II we have summarized the results obtained by us in the treatment of early syphilis with neocryl, either alone or in association with bismuth. For the sake of comparison we also record the results of treatment of a few similar cases with tryparsamide.

TABLE II.—Comparing the Action of Tryparsamide and Neocryl Respectively on Cases of Primary Syphilis

Name	Lesion	Spirochaeta pallida	Blood W. R.	Date of Treatment and Amount of Drug	Immediate Result	Remarks and Subsequent History
P.378	Chancre	—	+	<i>Tryparsamide</i> Feb. 18–March 28, 1935—36 grams	Slow healing (possibly healing commenced before treatment started)	May 27, muco-cutaneous relapse. W.R. +
274	Chancre	+	+	Feb. 18–March 5, 1935—15 grams	Primary not healed. Rash developed	Spirochaetes persisted throughout. Healed later by N.A.B. and bismuth
279	Chancre	+	—	March 1–14, 1935—10 grams	Lesion extended	Later healed by N.A.B. and bismuth
E.222	Chancre	+	—	<i>Tryparsamide and Bismuth (Bisozyl)</i> Aug. 12–Sept. 30, 1935—24 grams tryparsamide, 1 gram bismuth	No improvement	Oct. 2–23, 9 grams neocryl, 0.3 gram bismuth (bisozyl); ulcer healing. Defaulted
E.269	Chancre	+	—	Aug. 19–Sept. 6, 1935—14 grams tryparsamide, 0.7 gram bismuth	Not healed	Defaulted
W.	Chancre and rash	+	+	<i>Neocryl</i> Feb. 16–June 6, 1935—60 grams	Chancre healed, rash faded	Relapse in June, which responded to N.A.B. and bismuth
R.	Chancre and rash	+	+	Feb. 15–March 21, 1935—33 grams	" " "	Relapsed with secondary signs
A.	Chancre and rash	—	+	Feb. 28–May 3, 1935—26 grams	Healed	Went to India
Al.	Chancre	+	+	March 7–May 9, 1935—32 grams	"	Went to sea
E.309	Chancre	+	—	<i>Neocryl and Bismuth (Bisozyl)</i> Aug. 30–Oct. 14, 1935—36 grams neocryl, 1.2 grams bismuth	Healed on Sept. 16, 1935	Feb., 1936. Still healed, but has had further routine treatment with bismuth. W.R. negative
D.	Chancre	+	+	Sept. 13–Oct. 22, 1935—24 grams neocryl, 2.4 grams bismuth	Healed	Feb. 24, 1936. Still healed. W.R. +
E.357	Chancre	—	+	Sept. 23–Nov. 18, 1935—24 grams neocryl, 0.8 gram bismuth	Healed on Nov. 4, 1935	Jan. 2, 1936. Still healed. Not reported since
E.396	Chancre	+	—	Oct. 1–Nov. 22, 1935—36 grams neocryl, 1.2 grams bismuth	Healed on Oct. 15, 1935	Mar. 20, 1936. Still healed. W.R. negative
E.437	Chancre	+	—	Oct. 22–Dec. 31, 1935—36 grams neocryl, 1.2 grams bismuth	Healed on Nov. 5, 1935	Mar., 1936. Still healed. W.R. negative
E.340	Chancre	+	+	Sept. 25, 1935–Jan. 6, 1936—45 grams neocryl, 1.5 grams bismuth	Healed on Sept. 30, 1935	Went to London Jan. 14, 1936
E.510	Chancre and rash	+	+	Nov. 22, 1935–Jan. 27, 1936—36 grams neocryl, 1.2 grams bismuth	Healed on Dec. 9, 1935	Mar. 16, 1936. No signs of syphilis. W.R. negative
P.	Chancre and rash	+	+	Dec. 17, 1935–Jan. 24, 1936—24 grams neocryl, 1.6 grams bismuth	Healed and rash faded by Jan. 3, 1936	Feb. 5, 1936. Still healed. W.R. +. Went to sea

TABLE III.—Showing the Action of Neocryl on Cases of Tertiary Syphilis

Name	Lesion	Blood W. R.	Date of Treatment and Amount of Neocryl	Immediate Result	Remarks and Subsequent History
K.	Ulcer, leg	+	Nov. 29, 1934-Jan. 17, 1935—22.5 grams	Healed	
E. S.	Ulcer, foot	+	Feb. 14-Mar. 21, 1935—20.5 grams	Healed	Untraceable
F.	Ulcer, leg and foot	+	June 13-July 25, 1935—19.5 grams	Healed	Went to Southampton to go away to sea
H.	Tertiary rash	+	July 22-Aug. 30, 1935—25 grams	Healed	Jan. 5, 1936: Still completely healed. Wife reports at work regularly since Aug., 1935
W.	Generalized tertiary rash	+	July 22-Aug. 30, 1935—23 grams	Healed	Jan. 31, 1936: Perfectly well; has been working since August
Z.961	Ulcer, tonsillar fossa	+	July 22-Oct. 15, 1935—36 grams	Completely healed October 15	Had previously resisted bismuth and kharsulphan. Nov. 12, 1935-Feb. 6, 1936—2.4 grams bismuth (oil sol.). April 16: Still healed
E.241	Ulcer, tongue	+	Aug. 8-Sept. 9, 1935—18 grams	Healed on Sept. 9	Feb., 1936: Still healed, but has had further treatment with N.A.B. and bismuth
McQ.	Rupial rash	+	Sept. 6-Oct. 22, 1935—24 grams	Healed	Untraceable
E.282	Ulcer, right arm	+	Oct. 29-Dec. 3, 1935—15 grams irregularly	Slow improvement	Aug. 22-Oct. 22, 1935: Course of 27 grams tryparsamide, during which ulceration was spreading. Jan. 15, 1936: Broken down again, but said to be healing with bismuth 0.8 gram (bisoxyl). Healed by Feb. 19, 1936
E.375	Ulcer pre-arm, sinus at acromio-clavicular joint, generalized adenitis	+	Sept. 25-Dec. 9, 1935—36 grams	Ulcer healed Oct. 14	Dec. 9: Ulcer broke down again, then put on neokharsivan and bismuth and improved
McG.	Ulcer, scalp	+	Oct.-Dec., 1935—22.5 grams	Healed	Mild dermatitis; rapidly subsided. Feb. 5, 1936: Now quite well
E.	Ulcer, leg	+	Oct.-Nov., 1935—16.5 grams	Healed	Died apoplexy
Y.189	Subcutaneous gummata, thigh	+	Oct. 8-Dec. 17, 1935—30 grams	Gummata disappeared by Oct. 13	Seen on Jan. 7, 1936, with an arsenical dermatitis. Feb. 6: Dermatitis now subsided
E.431	Large ulcer, leg	+	Oct. 14, 1935-Jan. 2, 1936—45 grams	Healed Dec. 12	Feb. 10: Still healed
Z.869	Ulcer, nose	+	Oct. 1-Dec. 10, 1935—30 grams	Very slow improvement	Oct. 16, 1934-Jan. 15, 1935: Kharsulphan. Feb.-Mar., 1935: Pot. iod. Mar.-Sept., 1935: Neokharsivan and bisoxyl. These courses produced no improvement in lesion. Healed with novostab, after course of neocryl
Y.184	Ulcer, soft palate and lip	+	Oct. 1-Dec. 10, 1935—33 grams	Not healed	Healed after 0.75 gram novostab
M.485	Ulcer, soft palate	+	Oct. 3-Dec. 18, 1935—20 grams	Healed	Dec. 27, 1935: Broke down and put on sod. iod. and bismuth
E. A.	Tertiary rash	+	Nov. 1-Dec. 13, 1935—21 grams	Healed	Jan. 31, 1936: Working perfectly well
M.	Ulcer, arm	+	Nov. 25 Dec. 2, 1935—4.5 grams	Healed after second injection	Defaulted
N.	Ulcer, leg	+	Nov., 1935-Feb. 26, 1936—28.5 grams irregularly	Healed after five injections	General health much improved
Ca.	Ulcer, leg	+	Nov. 5, 1935-Jan. 8, 1936	Greatly improved, but did not heal completely	Healed with novostab
G.	Ulcer, foot	+	Dec. 27, 1935-Jan. 17, 1936—10.5 grams	Healed Jan. 17, 1936	Defaulted. Mar. 11 seen again. Healed and well
E.520	Ulcer, right arm	+	Nov. 27, 1935-Jan. 6, 1936—24 grams	Healed by Dec. 30	Mar. 18: Still healed
E.470	Annular syphilide	+	Nov. 14-Dec. 27, 1935—27 grams	Little, if any, improvement	Healed by 1.65 grams novostab
M.509	Tertiary syphilide, papulo-squamous	+	Nov. 15, 1935-Jan. 25, 1936—25 grams	Infiltration disappeared	Jan. 25: W. R. +

It is seen that, in conformity with the generally accepted views, tryparsamide had no action in four of the five cases of primary syphilis; even in the fifth its action was doubtful, since healing was very slow and had possibly commenced before treatment was instituted; in all the twelve patients treated with neocryl, however, the primary and secondary lesions disappeared. We do not, of course, wish to suggest that neocryl alone is an adequate treatment for this stage of syphilis; the fact that of the four cases so treated two at least quickly relapsed is strong evidence that by itself it is not adequate. Whether in combination with bismuth it is capable of producing permanent benefit at this stage of the disease is a matter which only time and further observations can reveal. These observations, however, suffice to indicate that neocryl, in contradistinction to tryparsamide, has a very definite action upon early syphilitic lesions.

Tertiary Syphilis

The results of treating twenty-five patients, suffering from various tertiary manifestations, with the new drug are summarized in Table III. In twenty cases the lesions were completely healed before the cessation of the course of treatment. In two of these (E.375, M.485) the lesions broke down again after the cessation of treatment, and were subsequently treated with N.A.B. or sodium iodide, and bismuth. It is interesting to note that one patient (Z.961), who was cured with neocryl, and another (Z.869),

who showed slight improvement under the drug, had previously resisted prolonged courses of sulpharsphenamine (kharsulphan), neoarsphenamine (neokharsivan), and bismuth. Of the five cases which showed little or no improvement as a result of treatment with neocryl one (E.282) improved under treatment with bismuth, and four (Z.869, Y.184, C., and E.470) rapidly responded to neoarsphenamine (novostab).

The general conclusion from this work appears to be that neocryl is definitely active in tertiary manifestations of syphilis. It should be noted that these patients were not prepared by any preliminary treatment with iodides, and that two, who had resisted previous treatment with kharsulphan and bismuth, responded to the new drug. The only toxic effects noticed in this long series of cases was mild arsenical dermatitis in two patients; this trouble subsided in both cases. There would appear to be a *prima facie* case for believing, therefore, that neocryl has a definite place in the therapy of tertiary syphilis.

Neurosyphilis

The drug was tried on a number of cases of neurosyphilis of different types. Two very early cases of neurosyphilis, without symptoms of nervous involvement, were treated with neocryl.

Wil.—This patient was admitted on January 30th, 1935. His general condition was poor, and he had been losing weight; he did not feel well. The reflexes were normal.

The Wassermann reaction of the blood was ++ and that of the cerebro-spinal fluid +++-; in the latter case the protein content was normal, the cell count 1 per c.mm., and the Lange reaction 0012211000. From March 7th to July 23rd he was given weekly doses of 3 grams of neocryl, to a total of 36 grams. On April 10th he was feeling very well; he had gained 27 pounds in weight, was back at work, and had no visual disturbance. On July 3rd he was very well and working; his sight was normal. The blood Wassermann reaction was ++; that of the cerebro-spinal fluid was ++++, and the protein content and cell count were normal, while the Lange reaction was 0000000000. On January 24th, 1936, he was very well, and had been working continuously. The blood Wassermann reaction was ++, but the cerebro-spinal fluid was normal in all respects.

Da.—This patient was admitted on July 8th, 1935, with a climatic bubo, profound anaemia, and cachexia, but no clinical signs of syphilis. The blood Wassermann reaction was +, that of the cerebro-spinal fluid was -, the protein content 0.015, the cell count 16 per c.mm., and the Lange reaction 4322100000. From August 23rd to October 25th he was given neocryl weekly in 3-gram doses, the total administered being 30 grams. His physical condition had then improved remarkably. He refused further treatment, and insisted on returning to India.

The following three cases of tabes dorsalis were treated with neocryl alone.

Hor.—This patient was admitted on August 9th, 1934, with advanced tabes. He had been pensioned owing to inability to work because of ataxia and lightning pains, but was mentally normal. Romberg's sign was present, as were the pupillary signs; the knee-jerks were absent, but the vision and fundi were normal. The Wassermann reaction was negative for the blood, but in the cerebro-spinal fluid it was +++-; the protein content was 0.02, the cell count 2, and the Lange reaction was 0001210000. From August 16th to November 11th he received 36 grams of neocryl in weekly doses of 3 grams. On January 14th, 1935, some slight clinical improvement was noted, and the sight was normal. The blood Wassermann reaction was negative; that of the cerebro-spinal fluid was +++-, the protein content was 0.015, the cell count 2, and the Lange reaction 0001000000. Between January 17th and April 4th he received 36 grams of neocryl in weekly doses of 3 grams. On April 12th the Wassermann reaction of the blood was negative. Between May 30th and August 9th he received a third series of neocryl in the same doses, and on August 28th some clinical improvement was noticed; the sight was normal. The Wassermann reaction of the cerebro-spinal fluid was +++-, the protein content 0.02, the cell count 2, and the Lange reaction negative. The patient was discharged on September 23rd feeling very well. On January 23rd, 1936, he reported himself as fit and working as a hospital orderly.

Mas.—This patient was admitted on November 12th, 1934, with advanced tabes and complete ataxia. He was unable to stand, but was mentally normal. He had Argyll Robertson pupils, the fundi were normal, and the knee-jerks absent. The blood Wassermann reaction was ++; that of the cerebro-spinal fluid was ++++, the protein content was 0.015, the cell count 8, and the Lange reaction 544443210. Between November 29th, 1934, and February 7th, 1935, he received 36 grams of neocryl in weekly doses of 3 grams. On February 8th the fundi were normal. The Wassermann reaction of the blood was ++, and of the cerebro-spinal fluid ++++; the protein content was 0.02, the cell count 2, and the Lange reaction 3211110000. Between March 28th and May 30th he was given 30 grams of neocryl in weekly doses of 3 grams. On June 4th the fundi were normal, the blood Wassermann reaction was ++, and that of the cerebro-spinal fluid +++-; the protein content was 0.03, the cell count 4, and the Lange test was completely negative. Between June 22nd and August 2nd he received 18 grams of neocryl in weekly doses of 3 grams. He was discharged as able to work on October 14th, and on January, 1936, he was at work as a ship's engineer.

Gre.—This patient was admitted on December 8th, 1934, with early tabes and gastric crises; he had had a laparotomy. The fundi were normal. The blood Wassermann reaction ++, and that of the cerebro-spinal fluid ++++; the protein

content was 0.025, the cell count 5, and the Lange reaction 1123221000. From January 17th to March 22nd, 1935, he was given a total of 30 grams of neocryl in weekly doses of 3 grams. On March 22nd he was clinically much improved. The fundi were normal. The blood Wassermann reaction was ++, and that of the cerebro-spinal fluid was ++++; the protein content was 0.03, the cell count 2, and the Lange reaction 1123210000. Between April 11th and May 6th he was given 18 grams of neocryl in weekly doses of 3 grams, and between August 2nd and September 6th a total of 15 grams in the same doses. On August 6th the blood Wassermann reaction was +. Between September 21st and November 22nd he received a total of 30 grams of neocryl in 3-gram weekly doses. On January 23rd, 1936, he was back at work as a tram conductor, much improved clinically.

The following patient was treated first by malaria therapy and subsequently with neocryl.

Pat.—This patient was admitted on July 11th, 1935, with advanced tabes, lightning pains, and ataxia; he was bed-ridden and unable to walk. There was a deep ulcer over the left ankle; Argyll Robertson pupils were present, and the knee-jerks were absent. The blood Wassermann reaction was ++, and that of the cerebro-spinal fluid ++++, the protein content was 0.04, the cell count 4, and the Lange reaction 1111100000. He received malaria therapy from July 25th to August 9th. The ulcer was unchanged on September 26th, when the Wassermann reaction of the cerebro-spinal fluid was ++++, the protein content 0.05, the cell count 6, and the Lange test 4432210000. Between September 27th and November 29th he received a total amount of 30 grams of neocryl in weekly doses of 3 grams. On December 2nd he was clinically improved, and the ulcer had almost healed. The sight was normal. The blood Wassermann reaction was ++, and that of the cerebro-spinal fluid ++++; the protein content was 0.03, the cell count 2, and the Lange reaction was completely negative. On December 5th he was discharged, the ulcer having entirely healed. On January 27th, 1936, he was re-examined. He was found to be clinically improved and able to get about; the ulcer remained healed. The blood Wassermann reaction was positive, and that of the cerebro-spinal fluid ++++; the protein content was 0.05, the cell count 12, and the Lange reaction 4221110000.

The following four cases of taboparesis and general paresis were treated with neocryl alone.

Wes.—This patient was admitted on October 26th, 1934, with taboparesis, aortitis, precordial pain, and dyspnoea; an x-ray examination showed aortic dilatation. There were sensory disturbances in the legs with slight ataxia, but the vision and fundi were normal. The knee-jerks were absent. The Wassermann reaction of the blood was negative, but in the cerebro-spinal fluid it was +++-; the protein content was 0.035, the cell count 50, and the Lange reaction 4555332100. From November 1st, 1934, to January 17th, 1935, he received a total amount of 36 grams of neocryl in weekly doses of 3 grams. On January 25th he was clinically greatly improved, and the fundi were normal. The Wassermann reaction of the blood was negative, and that of the cerebro-spinal fluid was +++-; the protein content was 0.025, the cell count 12, and the Lange reaction 5555433100. He refused further treatment because he felt so well, and returned to Africa.

Ro.—This advanced taboparetic patient was admitted on December 3rd, 1934, with pronounced mental symptoms and catalepsy. He had Argyll Robertson pupils, and the knee-jerks were absent. The Wassermann reaction of the blood was +, and of the cerebro-spinal fluid ++++; the protein content was 0.06, the cell count 12, and the Lange reaction 553333211. He was given weekly doses of 3 grams of neocryl to a total of 69 grams from January 8th to June 20th but did not improve, and was sent to a mental hospital.

McN.—This patient was admitted on January 4th, 1935, with advanced general paralysis and insanity, and Argyll Robertson pupils. The Wassermann reaction was +, and that of the cerebro-spinal fluid ++++; the protein content was 0.2, the cell count 144, and the Lange reaction 5555443210. From January 17th to May 2nd he received weekly doses of 3 grams of neocryl, to a total amount of 48 grams. On

April 8th the blood Wassermann reaction was +, and that of the cerebro-spinal fluid +++++; the protein content was 0.09, the cell count 14, and the Lange reaction 5555543210. In consequence of progressive mental deterioration he was sent to a mental hospital.

Mor.—This advanced taboretic was admitted on June 5th, 1934, in a very depressed mental condition, with hypotonia and Argyll Robertson pupils. The knee-jerks were absent, and there was primary optic atrophy. The blood Wassermann reaction was negative, and that of the cerebro-spinal fluid was +++++; the protein content was 0.035, and the cell count 9. From July 12th to September 14th he received a total amount of 30 grams of neocryl in the usual weekly doses. On September 14th the blood Wassermann reaction was negative, and the optic atrophy had not been adversely affected by the neocryl treatment. On January 28th, 1936, he was reported to be bed-ridden and dirty in habits, mentally impaired, and refusing further treatment.

The following two cases were treated by courses of neocryl followed by malaria therapy.

Hol.—This patient with taboparesis was mentally abnormal when admitted on August 7th, 1934. He had lightning pains, the knee-jerks were absent, and there was slight Rombergism. The blood Wassermann reaction was +, and that of the cerebro-spinal fluid was +++++; the protein content was 0.08, the cell count 12, and the Lange reaction 5554433321. From August 30th to November 15th he received 36 grams of neocryl in the usual weekly doses. On November 16th the blood Wassermann reaction was negative, and that of the cerebro-spinal fluid was +++++; the protein content was 0.05, the cell count 12, and the Lange reaction 5544333300. From January 3rd to March 21st, 1935, he received another series of weekly doses of neocryl amounting to 36 grams. On March 26th the blood Wassermann reaction was ++, and that of the cerebro-spinal fluid +++++; the protein content was 0.07, the cell count 8, and the Lange reaction 5555543100. From April 18th to May 3rd he received 9 grams of neocryl in the usual weekly doses, and from May 10th to the end of June malaria therapy. On June 26th there was mental and general improvement. The blood Wassermann reaction was ++, and that of the cerebro-spinal fluid +++++; the protein content was 0.04, the cell count 4, and the Lange reaction was completely negative. A further total amount of 30 grams of neocryl was given in the usual doses from July 26th to September 6th, and from February to May, 1936, another course of 30 grams in the usual weekly doses. He is being discharged to his work.

Dev.—This patient was admitted on October 7th, 1934, with general paralysis, attacks of giddiness, Argyll Robertson pupils, slight Rombergism, and bilateral interstitial keratitis. The blood Wassermann reaction was ++, and that of the cerebro-spinal fluid +++++; the protein content was 0.05, the cell count 30, and the Lange reaction 4422110000. He was given 36 grams of neocryl in the usual weekly doses from October 18th, 1934, to January 3rd, 1935. On January 7th the Wassermann reaction of the cerebro-spinal fluid was +++++; the protein content was 0.03, the cell count 4, and the Lange reaction 4431000000, and he was discharged. He was readmitted in September with condition unchanged. Malaria therapy was employed from October 23rd to November 16th. On December 13th the Wassermann reaction of the cerebro-spinal fluid was +---; the protein content was 0.025, and the Lange reaction was completely negative. From December 13th, 1935, to February 14th, 1936, he was given 30 grams of neocryl in the usual weekly doses. On March 4th the keratitis had improved, and the patient was very well mentally and physically.

The following patients were treated with malaria therapy, followed by neocryl.

Be.—This patient, admitted on August 23rd, 1935, had advanced general paralysis; he was fatuous, had pronounced tremors, and the reflexes were exaggerated. The blood Wassermann reaction was ++, and that of the cerebro-spinal fluid +++++; the protein content was 0.05, the cell count 12, and the Lange reaction 5432110000. He had been treated previously from January to September, 1933, with 72 grams of trypanamide. In September, 1935, he received malaria therapy. On October 9th the Wassermann reaction of the

cerebro-spinal fluid was +++-; the protein content was 0.03, the cell count 4, and the Lange reaction 5443210000. On October 18th he was discharged, no longer fatuous; the tremors had disappeared, and there was considerable general improvement. From that date until December 20th he received a total amount of 30 grams of neocryl in the usual weekly doses, and on February 26th this treatment was continued. The Wassermann reaction of the blood was +, and his wife reported that he was greatly improved, though not quite normal.

Wat.—This patient, admitted on July 24th, 1935, was a taboparetic with loss of memory; he had had epileptiform seizures, and now had a loss of vibration sense in his legs, Argyll Robertson pupils, and no knee-jerks. The Wassermann reaction of the blood was ++, and of the cerebro-spinal fluid +++++; the protein content was 0.03, the cell count 2, and Lange's reaction was completely negative. From August 9th to September 2nd he had malaria therapy. On September 26th the Wassermann reaction of the cerebro-spinal fluid was +++++, and the protein content, cell count, and Lange reaction were normal. From September 26th to February 26th he had a total amount of 51 grams of neocryl in irregular weekly doses. He improved, and returned to work as a taxi-driver. On March 22nd he returned with jaundice, which subsequently disappeared.

Abstract of Results

The results of treating cases of early neurosyphilis and tabes are distinctly encouraging. Two early cases of neurosyphilis without definite nervous symptoms improved greatly in general condition, and in one the cerebro-spinal fluid (Wassermann and Lange tests) became normal. Three well-marked cases of tabes treated with neocryl alone improved very greatly in physical condition, and the patients are now all back at work. In another case of tabes a course of neocryl was administered after malaria treatment; this was followed by marked improvement. This patient at the commencement of treatment was a very advanced bed-ridden case with a deep, penetrating ulcer at the ankle, but since treatment has so far recovered as to be able to get about the ward, and the penetrating ulcer has now healed. In this case it is, of course, impossible to assess the relative parts played by the malaria and the drug. The malaria therapy had certainly a good deal to do with the improvement, but the subsequent course of neocryl appeared to have a further beneficial effect, and it is noteworthy that the penetrating ulcer, which was apparently unaffected by malaria therapy, cleared up under treatment with neocryl.

A comparatively early case of taboparesis (*Wes.*) was treated with a single course of neocryl, and was so much improved that he insisted upon returning home and was lost sight of. Two more advanced cases (*Hol.* and *Dev.*) received but slight benefit from the first courses of neocryl and were subsequently treated with malaria, which was followed up by further treatment with the drug. These patients derived definite benefit from this therapy. Three other patients (*Ro.*, *McN.*, and *Mor.*) did not appear to receive the slightest benefit from the drug, and went steadily downhill. Two more patients (*Be.* and *Wat.*) were treated by malaria therapy followed by neocryl. One of these patients, who was completely fatuous on admission, derived very great benefit from the treatment, and the other recovered sufficiently to return to his work as a taxi-driver. The notes of the cases given above show, however, that no small part of the benefit resulted from the malaria therapy.

Nigerian Trypanosomiasis

A quantity of the new drug was sent to Dr. H. M. O. Lester, deputy director, Tsetse Investigation, Nigeria, who had kindly consented to try it on cases of Nigerian sleeping sickness. The preliminary results of this trial are summarized in Table IV.

TABLE IV.—Summary of Results Obtained in the Treatment of Cases of Nigerian Sleeping Sickness with a Single Course of Neocryl

Case No.	Duration of Symptoms	Condition at Commencement of Treatment				Treatment		Condition Immediately after Treatment			Remarks	
		Clinical	Blood Tryps.	Glands Tryps.	Cerebro-spinal Fluid		Dates	Total Amount of Neocryl Given	Clinical	Cerebro-spinal Fluid		
					Globulin	Cells				Globulin		Cells
4503	3 mnths	Weakness, severe headache, sleeplessness at night, sleeping during day, adenitis	+	+	+	28	28/6/34 to 18/9/34	28 grams (2 gram doses)	Quite strong, no headache, sleeps well at night, not in daytime; adenitis gone, blood negative	+	219	
4521	2 "	Weakness, headache, loss of weight, somnolence, adenitis	+	—	++	238	7/7/34 to 24/9/34	27 grams (2 gram doses)	Quite strong, no headache, sleeps normally, put on weight; no adenitis, blood negative	Slight haze	39	
4510	1 year	Weakness, very severe headache, somnolence, dullness, slurring speech, emaciated	+	—	+++	332	5/7/34 to 24/9/34	25 grams (2 gram doses)	Quite strong, no headache, sleeps normally, appears and looks perfectly normal; no adenitis, blood negative	—	1	
B.K.	mnths	Weakness, severe headache, somnolence, dullness, some adenitis, well nourished	+	—	Normal	Normal	26/6/34 to 9/8/34	15 grams (2 gram doses)	Quite strong, no headache, sleeps normally; no adenitis, blood negative			Absconded 12/8/34
K.S.	1 month	Weakness, somnolence, slight adenitis	+	—	"	"	26/6/34 to 15/9/34	29 grams (2 gram doses)	Quite strong, sleeps normally; no adenitis, blood negative			Absconded 15/9/34
A.G.	6 mnths	Weakness, headache, somnolence, dullness, loss of weight, some adenitis	+	+	+	42	30/6/34 to 23/8/34	19 grams (2 gram doses)	Quite strong, no headache, sleeps normally, blood and gland juice negative			Absconded 24/8/34
M.D.	6 "	Headache, well nourished, adenitis	—	+	Normal	Normal	10/7/34 to 9/8/34	11 grams (2 gram doses)	No headache, feels quite well, blood and gland juice negative			Absconded 12/8/34
K.G.	4 "	Weakness, severe headache, somnolence, some adenitis	—	+	+	128	10/7/34 to 26/9/34	29 grams (2 gram doses)	Stronger, no headache, no somnolence, able to do light work, blood and gland juice negative			Absconded
S.J.	6 "	Weakness, headache, somnolence, dullness, adenitis; history of syphilis six years previously and of "eye trouble" two months previously	+	+	+	70	9/3/35 to 30/4/35	31.5 grams (3 gram doses)	Quite strong, no somnolence, appears and feels perfectly normal; blood and gland juice negative	—	3	
1	3 "	Weakness, headache, fever, somnolence, dull, emaciated	—	+	++	296	?	28.5 grams (3 gram doses)	Improved, but still somnolent	+	150	
2	4 "	Weakness, headache, fever, emaciated	+	+	+	21	?	28.5 grams (3 gram doses)	Normal	—	0	

Dr. Lester, writing about the earlier cases treated with neocryl, stated that the stimulant action of a long course of treatment with the drug was very pronounced, and led to the same difficulty commonly met with in experimental work with tryparsamide. Patients felt so much better that they became tired of the prolonged treatment, and absconded before they had completed their course of injections. Of the first eight cases treated with neocryl five disappeared before the final examination of the cerebro-spinal fluid could be made. In every case patients showed very rapid clinical improvement, and thought themselves completely cured after they had had five or six injections.

Taken as a whole, the results were distinctly promising. No late cases were included in the series, but it is noteworthy that one very advanced case that had failed to react to full courses of tryparsamide, of Bayer 205, and of antrypol, for some reason or another improved markedly on being given a course of this new drug.

Summary

1. As a result of examining the trypanocidal activity of a large number of arsenicals of the general formula $\text{NaO.HO.O.As} \begin{array}{c} \diagup \diagdown \\ \text{NHCO} \end{array} (\text{CH}_2)_n \text{CO.R}$, one, sodium succinylomethylamide-*p*-arsonate (neocryl), was selected for further more detailed therapeutic trial.

2. In experiments on laboratory animals this compound compared favourably with tryparsamide in that it was found to be rather less toxic and of somewhat greater trypanocidal activity than the latter drug.

3. In man neocryl was well tolerated in doses which it is customary to employ for tryparsamide—namely, in weekly amounts of from 2 to 4 grams. The usual course consisted of the administration of this amount weekly until a total quantity of from 30 to 36 grams had been given. A number of patients had several such courses, the total quantity ranging from 66 to 141 grams, while one patient had an uninterrupted course of 69 grams, without showing any toxic signs. Apart from very occasional nausea and vomiting, the only toxic signs observed

were mild arsenical dermatitis in two patients and temporary jaundice in two or three very advanced cases of neurosyphilis after prolonged courses of the drug. Visual disturbances have not been encountered, but it would be unwise to assume that there is no risk of their production with neocryl, in view of the close chemical relationship of this drug to other pentavalent arsenicals known to produce such disturbances. In further trials, therefore, the same unrelaxed attention will have to be given to the possibility of this danger.

4. Neocryl exhibited in a pronounced degree the stimulating action which one is accustomed to associate with tryparsamide and the other pentavalent aromatic arsenicals.

5. In contrast with tryparsamide the new drug exerts a definite action in primary, secondary, and tertiary syphilis. In primary syphilis neocryl by itself is inadequate since, although the lesion cleared up rapidly, secondary manifestations subsequently developed. When neocryl was combined with bismuth the effects on primary syphilis seemed to be more permanent, and so far none of our patients treated in this way has relapsed.

6. Neocryl has a very definite action on tertiary manifestations of syphilis, the lesions disappearing completely in twenty out of twenty-five cases.

7. In early neurosyphilis and in tabes neocryl gave very satisfactory results.

8. Of eleven cases of Nigerian sleeping sickness treated by a single course of neocryl ten became clinically normal and the other was improved. Eight of these cases had at the time of treatment pathological spinal fluids; in three of them the fluid had become normal after the single course of treatment; in two it had become considerably improved; and in one pathological changes had apparently increased, although clinically the patient had become quite normal; the other two cases absconded owing to their clinical improvement before the completion of the treatment. In addition, a twelfth very advanced case, which had failed to react to very full courses of tryparsamide, of Bayer 205, and of antrypol, for some reason or other improved remarkably after a course of neocryl.

9. Whilst, of course, it is too early to make any definite claims for neocryl in the therapy of syphilis and trypanosomiasis, we consider that we have produced sufficient evidence to show that the drug is of definite value in the therapy of these diseases, and that a more extensive trial is warranted.

BIBLIOGRAPHY

- Brown, W. H., and Pearce, L.: *Journ. Amer. Med. Assoc.*, 1924, lxxii, 5.
 Moore, J. E.: *The Modern Treatment of Syphilis*. Baillière, Tindall and Cox, London, 1933, p. 63.
 Morgan, G. T., and Walton, E.: *Journ. Chem. Soc.*, 1931, lxxxiii, 615.
 Reese, H. H.: *Journ. Nerv. and Ment. Dis.*, 1933, lxxviii, 354.
 Strangeways, W.: *Ann. Trop. Med. and Parasitol.*, 1935, xxix, 231.
 Yorke, W., and Murgatroyd, F.: *Ibid.*, 1930, xxiv, 449.
 Yorke, W., Murgatroyd, F., and Hawking, F.: *Ibid.*, 1931, xxv, 313.

ESTIMATION OF PERCENTAGE OF ETHYL ALCOHOL IN THE BLOOD

BY

H. KENNETH V. SOLTAU, M.D.

The difficulty of deciding whether or not a person is under the influence of alcohol has led to the description of numerous tests for drunkenness.¹ That these are not infallible, however, is shown by the number of occasions on which even medical opinion differs when persons are accused by the police of being "under the influence of drink." The position in this country is thus by no means satisfactory, and, owing to the alleged association of alcohol with road accidents, the matter has lately assumed greater prominence.

Nature of the Test

From time to time² attention has been drawn to a method which has been in vogue in Sweden for some years, by which the percentage of alcohol circulating in the blood at the time of examination can be determined, and by which, it is claimed, a clinical diagnosis of "under the influence of drink" can be more readily sustained or disproved. This method was introduced by Professor Widmark of the University of Lund some years ago, and was published in detail in 1932.³ I have recently had the opportunity of studying this method and of applying it to patients who were under my care as medical superintendent of a voluntary home for alcoholics.

The method is based on the oxidation of the ethyl alcohol contained in a known weight of blood by a mixture of potassium bichromate and sulphuric acid. The distillation takes place under specified conditions, and the resulting distillate is then titrated.

Description of the Method

This may be described briefly in four parts.

Collection of Blood.—Special U-shaped capillary pipettes are used, each weighing about 300 mg. when empty, and holding 100 to 150 c.mm. of blood. They may be obtained already oxalated and fitted with rubber caps or with sealed glass ends. Blood is obtained from a prick of the finger or lobe of the ear, previously cleansed by a 1 in 1,000 solution of perchloride of mercury, not by ether or alcohol. The pipette is filled by capillary attraction, aided if need be by suction, and if the specimen has to await examination the ends are sealed.

Weighing of the Blood.—An extremely accurate balance must be used, such as Bunge's aperiodical analytical balance with damping device, which weighs rapidly to 0.0001 gram. The sealed ends are removed, and the pipette, full of blood, is hung on the hook of the balance and weighed. (Later the empty pipette is weighed, and the weight of the blood to be investigated is thus determined.) The blood is then expelled by blowing gently into a small glass cup fitted beneath the ground glass stopper of the triangular distillation flask, and situated from 1/2 to 1 cm. above the bottom of the flask.

The Distillation.—Into each flask is put 1 c.cm. exactly of a solution of known strength of potassium bichromate and sulphuric acid. The stoppered flask is made further air- and water-tight by means of a rubber cap. Three or four flasks, containing the bichromate acid mixture only, provide controls for each batch. The flasks are placed up to the neck in a water-bath, and kept at a temperature of 50° to 60° C. for two hours. They are then taken out and the stoppers cautiously removed. No fragments of dried blood must be allowed to fall into the acid mixture, as this would invalidate the test.

The Titration.—The contents of each flask are then diluted with 25 c.cm. distilled water. The mixture is titrated by N/100 sodium thiosulphate solution run in from a 10 c.cm. burette divided into fiftieths of a cubic centimetre. Half to one minute before titration a known amount of a 5 per cent. solution of potassium iodide is added. Starch is added later as an indicator. The end-point of the reaction is announced by the sudden disappearance of an intense blue colour, and is very distinctive. The difference between the number of cubic centimetres of thiosulphate solution used by the specimen containing blood and by the average control is noted, and by a simple calculation the percentage of alcohol in the blood can be ascertained.

The Test Employed

The relation between the percentage of alcohol in the blood and the clinical condition corresponded with that already noted by other observers.⁴ Thus a blood-alcohol percentage of 0.25 denotes drunkenness, whilst 0.15 points to moderate intoxication, but the percentages round about 0.1 provide the cases in which it is difficult to tell whether a man is "under the influence" or not. The minimum percentage which gave rise to this condition was noted as being 0.088.

Of the 100 estimations in the series:

- 63 were below 0.1 per cent.
- 18 were between 0.1 per cent. and 0.2 per cent.
- 13 were between 0.2 per cent. and 0.3 per cent.
- 5 were between 0.3 per cent. and 0.4 per cent.
- 1 was over 0.4 per cent.

The lowest value obtained was 0.00 per cent., and the highest 0.43 per cent.

Estimation of Tolerance

Gettler⁵ defines alcohol tolerance as oxidizing power, and remarks on the power of the habitué to oxidize alcohol very quickly. Below are twenty-two cases divided into two groups. Group A consists of habitual drinkers; Group B of bout drinkers.

A similar amount of diluted alcohol was given to each, and a specimen of the blood was taken at the time stated. Those patients in Group A, who had acquired a considerable tolerance for alcohol, tend to show a lower percentage of blood alcohol than those in Group B.

		<i>Group A</i>		
		Interval between time 2 oz. whisky given and sample of blood taken		Blood alcohol per cent.
Sex	Age			
M.	49	2.25	hours	0.122
M.	47	2	"	0.038
M.	46	2	"	0.088
M.	43	2.5	"	0.002
M.	48	1.5	"	0.016
M.	54	2	"	0.025
M.	56	1.75	"	0.045
M.	35	2	"	0.000
M.	44	2.5	"	0.110
M.	42	3	"	0.005
M.	52	2.5	"	0.006
		<i>Group B</i>		
M.	39	4	hours	0.113
M.	35	1	"	0.235
M.	50	3	"	0.270
M.	65	2.5	"	0.140
M.	40	1	"	0.187
M.	34	1.75	"	0.175
M.	29	2	"	0.144
M.	37	2.5	"	0.154
M.	52	2	"	0.160
M.	51	2.5	"	0.092
M.	55	4	"	0.120