TREATMENT OF FUSO-SPIRRILLARY INFECTION OF THE GUMS

BY

C. BOWDLER HENRY, L.R.C.P., M.R.C.S., L.D.S.

DENTAL SURGEON AND LATE LECTURER IN ORAL SURGERY, ROYAL DENTAL HOSPITAL, LONDON; LATE ARRIS AND GALE LECTURER AND HUNTERIAN PROFESSOR, ROYAL COLLEGE OF SURGEONS OF ENGLAND

Now that agranulocytosis is exciting interest all over the world it may be opportune to draw attention to an aetiological factor which is commonly present in the anginal form—namely, the fuso-spirillary symbiosis, or Vincent's infection. The tenacity of this infection, once it has gained a place in the mouth, is remarkable, and leads to a chronic state of so persistent and uniform a degree that the pathogenic potentiality of the disease, unless it flares up, is forgotten. The microbes are commonly regarded as normal saprophytes of the mouth and necessary concomitants of the common type of paradental degeneration which the average adult of both sexes sooner or later exhibits. That they are not so, however, is shown by any sequence of bacteriological examinations of the gums of different individuals. Indeed, many patients have considerable pyorrhoeic destruction and "pocketing" with a flora confined to leptothrix and cocci. On the other hand, it must be admitted that the fuso-spirillary pair are often found in even trivial gingivitis, but this does not absolve them from pathogenicity. As an examination of the literature shows, many a patient has paid for harbouring them, sometimes with his life. Hannson,¹ Kritchevsky and Seguin,³ Gornouec,⁴ Thompson,⁵ Palmer and Carr,⁶ to quote only a few instances, report cases where the removal of a tooth, especially the lower wisdom tooth, has been followed by a severe fulminating Vincent's stomatitis. Five of the cases cited ended in death, and they show several points in common with some of the cases of "agranulocytosis" excited by dental extraction,^{7, 8} or by the slighter trauma of "scaling."

Three features of the disease are worthy of emphasis. One is the way it persists in a subacute or chronic form after an acute attack, and leads at times to remote secondary lesions. Goadby,⁹ for example, relates a fatal case of sudden fuso-spirillary empyema in an apparently robust man eighteen months after an acute attack of Vincent's stomatitis. The second feature is its ability to gain an insidious entrance to the gingival tissues, so that a widespread and painless infection involving all the teeth is well established before any symptoms appear. The third feature is the predisposition—like that of recurrent agranulocytosis—to intermittent fulmination in the mouth and fauces, following periods of quiescence which sometimes last many months.

Epidemiology

The epidemiological characteristics of the disease were particularly apparent during the great war, when "trench mouth" was responsible for a large amount of invalidism in the troops.^{10 11 12 13} In 1917 the epidemic at Adrianople caused a number of deaths.¹⁴ While its communicability has always been recognized, the disease has not been made notifiable in this country, but in the U.S.A. Vincent's infection has been reported monthly by at least seventeen States since 1926.¹⁵ It is alleged to have been spread there by home-coming troops from Europe, and some alarm is felt at its continued increase.

Here in England, in hospital practice since the war, there seems to have been a steady dribble of cases, accompanied, as shown by Stallybrass,¹⁶ with seasonal rises; but an increased incidence has certainly become apparent in private practice, especially in the fashionable and cultured. This spread to a class hitherto immune

coincides with the appearance of agranulocytosis in this country—that is, from 1929 onwards. Adams, quoting from Krache and Parker,¹⁷ notes that in the U.S.A. agranulocytosis "affects mostly the better-class section of the community. Kitchin,¹⁸ discussing periodic or recurrent agranulocytosis, describes a tendency to chronicity associated with the common presence of Vincent's organisms, and Hartfall¹⁹ also speculates upon their aetiological significance. The close affinity of Vincent's infection to noma should also be borne in mind.

Importance of Prophylaxis

If these diseases are connected early recognition and treatment of fuso-spirillary infection of the mouth—especially in the subacute and latent cases—should not only lower the mortality from Vincent's disease, but should also reduce the gravity of agranulocytosis. Moreover, an attack of Vincent's does not confer immunity, and patients become "carriers," so it is important to treat this infection of the gums with greater thoroughness than is usual—which, indeed, its potentiality for harm demands. Smith, in the introduction to his monograph²⁰ on the various manifestations of fuso-spirochaetal disease about the body, emphasizes its preventability. He says:

"Finally, this is a preventable disease. The source of the infection is in diseased tonsils and diseased gums, and if these foci are eliminated then all the local and general infections with these anaerobic organisms might be prevented."

Nevertheless, from a dental point of view, treatment is difficult. Most topical and intravenous treatment, as subsequent bacteriological smears show, is only partially curative, and patients are reluctant, even after a severe attack, to consent to a prophylactic clearance of the teeth, though they might readily part with their tonsils. It seems worth while, therefore, to describe the only method which in my hands has yielded total and permanent elimination of the infection. It is sometimes laborious, but it is successful, as bacteriological rechecking proves, and it is based upon twenty years' experience. Smith postulates:

"Since the essential organisms in the disease are anaerobic, open drainage and free access of air are indicated."

This is the foundation of my technique.

Treatment of Acute Stage

When the case is first seen the diagnosis is confirmed by smears. A differential blood count should be made to detect any concurrent leukaemia or agranulocytosis, which should receive appropriate treatment at the same time. The patient should be isolated in a nursing home, where saliva-bearing articles can be effectively controlled and uninterrupted treatment maintained. The ulcers are very lightly scraped and washed clear of the pseudo-membrane by a thorough syringing with hydrogen peroxide, or, better, by a de Vilbiss dental spray containing normal saline and connected through a reducing valve to a cylinder of oxygen. It is important that the operator and attendant nurse should be protected by gloves, gown, and mask. If this cleansing is done delicately but copiously the necrotic debris is entirely removed, and a bleeding surface at the base of the ulcer is revealed. This raw area is painted with an escharotic, preferably pure carbolic acid, and the patient washes the mouth every half-hour during the daytime with a warm sodium perborate solution (one drachm to a tumbler of water). This treatment will rapidly reduce the acute symptoms; in twenty-four hours the ulcers will be clean and fibrin-covered, and in a few days epithelialized.

After meals the mouth, and especially the interdental spaces, should be sprayed to remove food debris. Bowman's mixture²² of vin. ipecac. 3 iv, glycerin 3 j,

liq. arsen. (Fowler) ad 3 j, or some other arsenical preparation; applied locally, may accelerate resolution by attacking the treponemata. Intravenous kharsivan and similar products are probably of great use in helping to arrest widespread ulceration of the cheek and fauces, but are particularly disappointing in the treatment of the gingival form, mainly because the organisms lodge in mechanical crevices and in the necrotic tissue of the gingival troughs, out of reach of the circulation. Indeed, the essence of treatment is to remove by physical means

Reproduction of Vincent's original illustration of a smear.²¹

the felt-like protection of slough and apply the medicaments to the freshly invaded surface, which is then kept clean and aerated by the oxygen-liberating mouth-wash. It is of practical importance, therefore, that the patient be encouraged to maintain for the first two days a continuous atmosphere of oxygen in the mouth.

After the first twenty-four hours the gums are no longer tender, and the gross tartar, which is a hindrance to easy flushing of the interdental spaces and the gingival troughs, should be removed. Scaling should not be attempted on the first day, but every succeeding day should see a progressive clearance of tartar. As the healed surface hardens the patient carries out an appropriate toilet with interdental tape and a soft toothbrush sterilized in spirit.

Bacteriological Examination

The acute ulceration disappears rapidly and easily, but it is a fallacy to suppose that the fuso-spirillary infection has thus been banished from the average mouth. At this stage it is wise to check the bacteriology by smears taken from representative sections of the denture. A negative result may be expected in the freely cleansable places, and a positive one in such uncleanable sites as pre-existing pyorrhoea pockets, under the margins of leaking or ill-fitting crowns and fillings, around dead roots, and in the deep circumvallate gingival trough of partially erupted and misplaced wisdom teeth. The crypts of the tonsils should also be examined. These places are the sites of persistent infection in the subacute and chronic cases (common in private practice), where the fuso-spirillary organisms have gained a habitat insidiously without provoking acute ulceration, and it should be recognized that the average acute case treated with mouth-washes, no matter how thoroughly, does no more than resolve itself into one of this kind. I have never yet seen the organisms totally eliminated from all parts of the mouth so long as these catch-pits remain to become foci from which sporadic fulminations occur. Therefore, unless smears from many sites taken at this stage are negative, treatment must be continued as next described.

Sub-acute Stages

These are the stages most popularly confused with parodontosis or pyorrhoea, with the result that the patient is often dismissed with a diagnosis of "gingivitis" and told to use an antiseptic mouth-wash. A bacteriological examination is necessary to confirm the diagnosis, but certain characteristic clinical features are usually present. In the subacute stage the gums are swollen and purplish-red at the gingival margins, but not necessarily round the whole denture. Only certain teeth, where stagnation is more prevalent, may be affected. There is a strong tendency to bleed upon light digital pressure or spontaneously. Although in the later stages the loosened teeth may become tender, there is no pain in the earlier stages nor undue tenderness in the gums. Very often the patient's first symptom is to find one morning a blood-stained patch of saliva on the pillow, or to notice during conversation a sudden and irritating seepage of blood into the mouth. The characteristic fetor may be present in the breath, although it might be unperceived in more fastidious patients.

Chronic Stage

In the chronic stage the gums are shrunken rather than swollen, and are normal in colour, but around the more grossly affected teeth there are extensive pockets, which bleed easily on the introduction of a fine blunt seeker. Patients of this subacute and chronic group are "carriers." They are a menace not only to other people, but to themselves as well, for they do not acquire immunity. At any time an acute and fatal exacerbation in the mouth or throat may supervene, or remote disease may be started.

"Situating in the tonsils and about the teeth they [Vincent's organisms] are in a strategic position to initiate disease locally in the tonsils, gums, cheeks, tongue, pharynx, and larynx; or to invade the eyes, ears, or sinuses. They may be aspirated and infect the trachea, bronchi, or lungs; or may be swallowed and cause disease in the oesophagus, appendix, or colon; by means of fingers contaminated with saliva these anaerobic organisms may be carried to the skin or to the external genitalia. Through emboli they may reach the brain, lungs; or other internal organs" (Smith).

Furthermore, the lesions around the teeth produced by fuso-spirillary destruction are favourable sites for intercurrent invasion by the strepto-staphylococcal group, and therefore superadded to the menace of fuso-spirochaetal disease are those of acute local pyogenic infection, such as osteomyelitis of the jaw and spreading cellulitis of the neck, and of chronic toxæmias and bacteraemias. At best the persistence of Vincent's organisms in the mouth leads to gross destruction of the alveolus of the dental sockets, and admits at the same time the organisms producing the common chronic toxæmias. At worst it leads to unexpected death either of the patient or of a contact.

Diagnosis

Chronic Vincent's infection (in which is included the subacute stage) is a disease of filth and stagnation. It is never seen in the edentulous mouth. It is not seen in all parts of the averagely clean mouth, but only in those places where effectual toilet is lacking, and where, *per contra*, miniature cesspits exist—that is, where the patient is lazy or where uncleanable "pockets" prevent his attention being effective, however hard he tries. Likewise, it is found in the depths of the tonsillar crypts. The incidence and persistence in unhygienic areas of the gums is stressed by many writers: D. T. Smith,²³ Colyer and Sprawson,²⁴ Mead,²⁵ Gill,²⁶ Dalitsch,²⁷ Puterbaugh,²⁸ Goadby, Bunting,²⁹ Davenport,³⁰ and others. Consideration of the morbid anatomy and the predisposing local factors, therefore, indicates the line of effective treatment.

Diagnosis is confirmed by smears, and is easy and positive. The only difference between the stages to be recognized in the smears is the greater predominance of the fuso-spirillary organisms in the acute phase. There appears to be no significant change in shape, size, or staining properties to indicate whether the specimen is from a fulminating mortal case or from a "carrier." In the more chronic stages the proportion of the two organisms to each other, which is roughly equal in the acute conditions, tends towards the reduction of the fusiforms, and at the same time other organisms common to septic "pockets" creep in. A full set of dental radiographs is required to show the degree of destruction of the alveolus, the presence of "pockets," dead teeth, obscure carious cavities, and ill-fitting crowns, fillings, and bridges, and the state of partially erupted teeth. A minute clinical dental examination, along with skiagrams for comparison, enables a scheme of treatment to be drawn up.

Scheme of Treatment

In the first place, on the principle that ultimate success depends upon an extreme degree of oral cleanliness, the patient must be instructed in the *raison d'être* of the therapeutic measures, and his complete co-operation in prophylaxis must be enlisted. He should be shown a method of oral hygiene which has as its object the meticulous cleansing of the gingival sulcus around each and every tooth, by means of toothbrush, dental tape, and spray.

All old amalgam fillings are removed and carious holes are cleaned, the cavities thus left being filled with fresh cement until satisfactory permanent fillings can be inserted. Fixed bridges, harbouring putrefying food, and ill-fitting shell crowns are also taken off, leaving the spaces cleansable. "Pockets" are noted for excision and dead teeth for removal. Measures should also be taken to counteract persistent mouth-breathing. Finally a critical scaling of all subgingival tartar is performed so that the necks of the teeth are left perfectly clean.

Dental Technique

Under intratracheal anaesthesia an attempt is made to deal with the whole mouth at one session. The dead teeth having been removed and the peridental sockets having been excised to their full depth and curetted, an electrocautery point is passed into the gingival sulcus around every tooth. The resulting coagulum is scraped and irrigated away. By this means, and by this means only, is a clean granulating surface left. Mouth-washes of hot sodium perborate solution and frequent irrigations with hydrogen peroxide are instituted during healing, which takes place rapidly and without discomfort. After a few days a searching bacteriological examination is again performed. The result should show only a few cocci and pus cells, significant of a healthy reaction. Should any spot disclose spirilla or fusiforms it must be treated again under an injection of novocain. The object of the treatment is, in the first place, to open up and clean out all stagnant areas, and, in the second place, to deal an effective blow at the Vincent's organisms by cauterizing the whole of their habitat at one operation. It is commonly thought possible to save the patient the expense and trouble of a nursing home by dealing with the mouth in stages in the dental chair under block novocain. The obvious disadvantage, which is proved by experience, is that the newly cauterized area becomes reinfected from the untreated places, and eventually it becomes obvious that no progress has been made.

A common focus of recurrence is found in the third mandibular molar. Even when fully erupted this tooth

rarely emerges so clear of the gum that it does not have a deep uncleanable sulcus behind it. When it is semi-erupted or impacted the pocket left under the hood of mucous membrane forms a nidus which, in several recorded cases, has not only led to repeated recurrence of the acute infection, but has been responsible for the death of the patient. These teeth should be extracted whilst the condition is under control.

Finally, when the mouth has healed and is bacteriologically clean, the tonsils, if they still harbour the organisms, should be removed before the gums become reinfected from them.

Conclusion

Vincent's infection of the gums is a preventable and a treatable disease, but it is essential that every detail of the denture be freely cleansable by the patient. The difficulty is that where many teeth are filled and fixed appliances are worn few patients can afford the large amount of therapeutic and dental restoration work which is entailed, and half-measures are useless.

Case Reports

CASE I.—ACUTE CASE UNCOMPLICATED BY FILLINGS, ETC.

October 8th, 1931.—A male aged 22 had gums previously very healthy and teeth more than normally clean; he had completed a dental overhaul two months previously. He contracted acute ulcerative gingivitis in Paris; the gums were painful and were bleeding easily. The teeth were festooned all around the denture by marginal ulceration; typical appearance and fetor. Bacteriological report on smears from several places (Dr. I. H. Maclean): "Films show a large percentage of red cells. Main organisms present: Vincent's spirilla 90 per cent., *B. fusiformis* 9 per cent., cocci 1 per cent. Pus cells rare. This film is the picture of an acute phase of ulcerative stomatitis when the spirilla are in the ascendancy, and there is very little reaction by way of pus cells by the patient."

October 8th, 1931.—Treated as outlined above, but with immediate cauterizing of the complete series of gingival troughs, after preliminary cleansing of the ulcerated area.

October 13th, 1931.—Smears made from sixteen representative areas around the teeth. A few pus cells were seen and cocci in all films, except that of one upper molar space. Comment: "The films show a great improvement on those of October 8th, and the only part where the infection can now be recognized is in the molar region 76/'' (I. H. Maclean). This isolated region was cauterized again and rapid healing occurred, leaving the gums perfectly healthy in colour, texture, and alignment.

December 20th, 1935.—Patient reported by letter from abroad that he continues to employ the method of dental toilet prescribed, and has had no return of the gingival symptoms.

CASE II.—CHRONIC VINCENT'S GINGIVITIS. REINFECTION FROM TONSILS

A female aged 35 complained of bleeding from the gums. She had been treated for "pyorrhoea," but without avail. A diagnosis of chronic Vincent's infection was made. Six sites were sampled for bacteriological examination (March 26th, 1935), and the same six sites, after cauterizing in a nursing home (April 5th, 1935), were later resampled (April 10th, 1935). These were examined by Dr. J. D. Benjafeld, who reported as follows:

Site	March 27th, 1935. Before treatment			April 10th, 1935. After treatment		
	Pus	B. fus.	Spir.	Pus	B. fus.	Spir.
1	+	+ (few)	+	nil	nil	nil
2	++	++	++	+	..	..
7	++	++	++	+	..	..
1	++	++	++	+	..	..
1	+	+	+	nil	..	..
7	+	+	+	+	..	..

Comment on report (April 10th): "The amount of pus present was very slight."

Healing occurred uneventfully and the patient remained well for two months, when she suffered from laryngitis with aphonia. Re-examination of the gums showed slight sponginess in the lower incisor teeth. Six suggestive sites from the gums and both tonsils were examined. Dr. Benjafield reported (June 28th, 1935): "All smears negative for Vincent except 32/ and 23/." It had not been possible to obtain satisfactory smears from the tonsillar crypts because of gagging, and the tonsillar films showed only "some pus cells and a mixed flora of organisms." Dr. Benjafield commented: "I am of the opinion that the tonsils constitute a definite septic focus from which the gums are being, and are likely to be, reinfected. The fact that Vincent's organisms were not found in the films from the tonsillar exudate does not necessarily exclude the possibility of their being present in the deeper part of the tonsils. In any case the rather poisonous flora grown in cultures justifies seriously considering the advisability of tonsillectomy."

CASE III.—INEFFECTIVENESS OF MOUTH-WASHES, SPRAYS, ETC.

A female aged 34, in bed making a difficult recovery from concussion after a motor accident, complained of acute soreness of the gums with typical fetor (July 31st, 1935). Smears from six sites were all heavily positive and typical of acute Vincent's infection. Dr. I. H. Maclean's comment is: "From the large numbers of spirilla and *B. fusiformis* this case suggests that there is some acute spirillary infection." Her dental surgeon reported that the patient had been a most fastidious woman, who regularly attended every two months for prophylactic polishing of the teeth, and that her gums had always been healthy. Presumably inability to attend to oral hygiene during the period of disablement (four and a half weeks) had allowed a state of stagnation of food which permitted the Vincent's organisms (source untraced) to gain a habitat. The patient was treated with intensive irrigation, spraying, and mouth-washes of hydrogen peroxide and sodium perborate solution, etc., as detailed above under the treatment of acute cases. The acute symptoms subsided and she was not cauterized. She returned on October 1st, 1935, for further examination after a sea trip for convalescence. In spite of assiduous home treatment, when re-examined the gums presented evidence of chronic fusio-spirillary infection, and ten sites selected for bacteriological examination (Dr. Lionel Whitby) were all positive for Vincent's infection. Dr. Whitby's comment reads: "The smear results show a heavy infection with Vincent's organisms around the 7/ and 8/, and a moderate infection round the 6/ and 7/. All other specimens showed small numbers of the same organisms, from which it may be deduced that the condition has at one time affected the whole mouth and is as yet not completely eliminated."

CASE IV.—INEFFECTIVENESS OF MOUTH-WASHES, SPRAYS, ETC.

A female aged 35 was admitted to a nursing home with acute Vincent's infection around all the gums. The infection was ultimately traced to a house-party where the host and hostess proved to be the "carriers," and of the guests three others besides the patient are said to have caught the contagion. This patient first became aware of having an oral lesion by repeated spontaneous haemorrhage from the gums. Smears were taken six weeks later during an acute attack. Dr. John Matthews reported (July 22nd, 1935) that all seven sites examined were positive. The patient was treated with scaling, spray, mouth-washes, etc. The acute symptoms rapidly subsided; the gums healed and appeared normal. In view of the encouraging improvement she decided not to be cauterized, but to see what she could achieve by her own efforts. She carried out her home routine enthusiastically at first, but relaxed later. When re-examined (January 23rd, 1936) the mouth and teeth were generally clean, but close inspection indicated signs of chronic infection. Smears taken from eighteen sites showed pus (sometimes very profuse), with leptothrix and streptococci in all the sites, and Vincent's organisms (sometimes in prodigious numbers) in eight of them.

CASES V AND VI.—NON-CAUTERIZING VERSUS CAUTERIZING

A female aged 38, and her husband aged 47, sought treatment because of bleeding and puffiness of the wife's gums. Clinically both patients presented symptoms of chronic Vincent's infection. Smears of wife (April 19th, 1933) and of husband (April 24th, 1933) were both heavily positive from several sites. Dr. I. H. Maclean's blood counts and commentaries were:

Wife's blood count (April 26th, 1933): red cells, 3,800,000 per c.mm.; white cells, 10,000 per c.mm.; haemoglobin, 76 per cent.; differential count—polymorphs, 56 per cent.; lymphocytes, 40 per cent.; other cells, 4 per cent. "The persistence of the *B. fusiformis* tends to show that the condition has reached a chronic stage, and there is a larger percentage of pus cells than usual in most ulcerative stomatitis. There is some anaemia, which is associated with an increased number of lymphocytes. These also show the chronicity of the disease, and that in the past the patient has not made a very good response to her infection."

Husband's blood count (April 26th, 1933): red cells, 4,920,000 per c.mm.; white cells, 7,000 per c.mm.; haemoglobin, 96 per cent.; differential count—polymorphs, 62 per cent.; lymphocytes, 36 per cent.; other cells, 2 per cent. "There is definite evidence of ulcerative stomatitis present in this patient. In the molar region it has reached the subacute stage with no evidence of healing. In the incisor region there is only a mild infection. The disease has so far had only local effects, and there is no evidence by the blood count that the patient's general health has suffered."

The wife, an American married one year to an English husband, gave a history of having suffered from an acute attack of Vincent's angina several years previously. This had been treated upon orthodox lines and was presumed to have been cured, but evidently persisted in her gums and she became a "carrier." The inference on the clinical, the bacteriological, and the haematological evidence is that she infected her husband. It was obviously necessary to treat both patients. Both were treated alike according to the method for subacute and chronic cases—that is, meticulous scaling, the removal of all crowns and leaking fillings, topical applications, and an intensive home routine. The husband consented to be cauterized (May 4th, 1933), but the wife, promising to redouble her efforts, refused. Forty-six days later smears were made from both patients from seven sites in their gums and from the crypts of both tonsils. The husband was negative, except for a dead tooth and a partially erupted wisdom tooth, which were immediately extracted. The wife was still positive both in the mouth and in the right tonsil.

REFERENCES

- Hannson, A.: *Centralbl. f. Chir.*, 1892, xix, 845.
- Marshall, John S.: *Journ. Amer. Med. Assoc.*, 1899, xxxiii, 1593.
- Kritchewsky, B., and Seguin, P.: *Dental Cosmos*, 1918, ix, 781.
- Gornouec: *Rev. de Stom.*, 1929, xxxi, 18.
- Thompson, L. E.: *Journ. Amer. Med. Assoc.*, 1929, xciii, 1063.
- Palmer, B. B., and Carr, M. W.: *Journ. of Dent. Research*, 1935, xv, 93.
- Hill, Harold P.: *Calif. and Western Med.*, 1926, xxv, 609.
- Adams, E. W.: Ministry of Health, Britain, Reports on Public Health, No. 76, 1935, p. 6.
- Goadby, K. W.: *Diseases of the Gums*, London; and *The Oral Mucous Membrane*, 1931, p. 51.
- Bouty, R. T. C.: *British Medical Journal*, 1917, ii, 685.
- Campbell and Dyas: *Journ. Amer. Med. Assoc.*, 1917, lxviii, 1596.
- Kritchewsky, B., and Seguin, P.: *Rev. de Stom.*, 1920, xxii, 613.
- Lange: *Deut. zahnartzl. Woch.*, 1915, 15.
- Sauerwald: *Berl. klin. Woch.*, 1917, liv, 111.
- Public Health Reports, Treasury Department, U.S.A., 1926 et seq.
- Stallybrass, C. O.: *Proc. Roy. Soc. Med.*, 1928, xxi, 1193.
- Krache, R. R., and Parker, F. P.: *Journ. Lab. and Clin. Med.*, 1933, xix, 799.
- Kitchin, P. C.: *Journ. Dent. Research*, 1934, xiv, 315.
- Hartfall, S. J.: *Lancet*, 1934, i, 62.
- Smith, D. T.: *Fusio-spirochaetal Disease*, London, 1932.
- Vincent, H.: *Ann. de l'Inst. Pasteur*, 1896, x, 510.
- Bowman, F. B.: *Lancet*, 1917, ii, 536.
- Smith, D. T.: *Op. cit.*
- Colyer, J. F., and Sprawson: *Dental Surgery and Pathology*, London, 1931, p. 468.
- Mead, S. V.: *Diseases of the Mouth*, fourth edition, London, 1932, p. 517.
- Gill, W. D.: *Texas State Journ. of Med.*, 1933-4, xxix, 11.
- Dalitsch, W. W.: *Journ. Amer. Dent. Assoc.*, 1934, xxi, 1257.
- Puterbaugh, P. G.: *Ibid.*, 1934, xxi, 1924.
- Bunting, W.: *Oral Pathology*, 1929, p. 322.
- Davenport, S. E., jun.: *Journ. of Dent. Research*, 1933, xiii, 65. (Discussion on A. Merritt's paper.)