TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some of the major medical disorders met with in general practice

THE TREATMENT OF ASTHMA

BY

GEORGE W. BRAY, M.B., M.R.C.P.

In using the term "asthma" I refer to that type of dyspnoea brought about by a spasmodic constriction of the bronchial muscle; at first paroxysmal in nature and principally expiratory in character; frequently occurring in members of families where other allergic diseases occur; and often accompanied in the past or present by other allergic conditions in the individual sufferer.

The condition is characterized clinically by wheeziness followed by intervals of apparent health, generally excited by an abnormal hypersensitivity to some foreign substance which is absorbed by inhalation (animal emanations, dusts, pollens, moulds, and fungi); ingestion (foods and drugs); injection (serums, drugs, bites, and stings); infection (bacteria and worms); or by contact (fabrics, chemical or physical agents); and frequently predisposed to by some psychic, nasal, toxic, endocrine, or dietary factor. Such a definition excludes the inspiratory difficulty associated with cardiac or renal disease, and the respiratory embarrassment associated with bronchitis, thymic enlargement, congenital laryngeal stridor, laryngismus stridulus, retropharyngeal abscess, and enlarged lymph glands in children. In adults it excludes the tachypnoea of hysterical subjects; the difficult breathing associated with chronic pulmonary tuberculosis, pulmonary new growths, and pathological states of the pleura; aneurysm; foreign body in the lung; and the severe type of bronchitis or low grade of pneumonia in advanced age.

The treatment of asthma may be discussed under two headings: first, the immediate treatment of the acute attack; and, secondly, the intensive treatment of the asthmatic sufferer to prevent further attacks.

Immediate Treatment of the Acute Attack

Inject subcutaneously in a child 2 minims, or in an adult 5 minims, of a 1 in 1,000 solution of adrenaline chloride. No additional benefit is to be derived from the more expensive proprietary preparations which contain adrenaline mixed with other constituents; frequently they cause intense palpitation, which is more discomforting than the asthma, as generally they are injected in doses of 1 cubic centimetre, and after they have been used for any length of time the asthma appears to be unrelieved by the usual adrenaline solution unless given in very large doses.

Adrenaline is rapid in its relief of bronchospasm, but unfortunately its effect is of correspondingly short duration. Hence, in severe attacks it is always the best procedure to leave the needle in position after giving the first dose, because smaller additional amounts are sure to be required to prevent the reappearance of the spasm. A 1 c.cm. Record syringe can be left filled with the adrenaline chloride solution, and a relative be instructed to attach the syringe to the needle and give two minims of the solution at the end of one hour, two hours, four hours, and eight hours, when as a rule the attack will be at an end and the patient have remained free from any recurrence in the meanwhile. In "status

asthmaticus" one or more minims of the adrenaline solution will need to be repeated more frequently—even as often as each minute until relief of the spasm is obtained, and then each quarter of an hour for the first hour, each half-hour for two hours, each hour for three hours, and each two to four hours until bedtime. In severer cases large amounts (but small individual doses) of adrenaline are necessary, and relief from the drug appears slower and less adequate. In these cases an additional solution is necessary, made up as follows:

Morphine sulphate	...	...	...	...	1/4 grain
Atropine sulphate	...	...	...	...	1/100 grain
Adrenaline chloride solution (1 in 1,000) to 10 minims.					

One minim of this special solution is injected every fifth minute instead of adrenaline alone, and the process is continued until the spasm is relieved or until the 10 minims of the solution have been injected. It is unwise to give more than 1/4 grain of morphine, even by this slow method; for, should the cough reflex be abolished and the spasm subside, large amounts of mucus will be poured out from the obstructed glands, and unless this is expectorated the patient will be drowned in his own sputum. More patients die from injudicious doses of morphine than from the severity of the asthmatic spasm. Once the spasm has been relieved by these means the plain adrenaline solution is continued as above until the patient has remained free for twenty-four hours. The maximum benefit from adrenaline is obtained when it is given at the earliest possible moment, in small doses, and repeated at gradually increasing intervals until the patient has been free for twenty-four hours.

Treatment After Cessation of Spasm

As soon as the patient begins to experience any relief the body temperature should be brought back to normal by additional blankets and hot-water bottles, and a tablet of ephedrine hydrochloride given. For an adult 1/2 to 3/4 grain may be ordered, and repeated each six hours if necessary; while for a child the dose is 1/8 grain under 1 year, 1/4 grain over 1 year, and 1/2 grain over 7 years. Unfortunately ephedrine causes unpleasant symptoms in a large proportion of cases, and many pharmaceutical efforts have been made to overcome these side effects. The synthetic ephedrine, such as pseudo-ephedrine manufactured by Burroughs Wellcome and Co., or ephedronin made by Merck, overcome these undesirable reactions in some patients, while others combine with the ephedrine such drugs as amytal (dose 3/4 to 1½ grains), or phenazone, as in the ephazone tablets.

If the patient is exhausted stimulants are indicated, and strong coffee in the morning, or towards evening strong hot alcohol, is advisable. Fluids are essential, and if the patient cannot take much fluid or glucose water by mouth, a glucose saline enema (3 ounces of glucose in 1 pint of normal saline) will serve a triple purpose; for not only does it suffice temporarily for food and fluid, but in the presence of flatulent distension of the abdomen or constipation relief may be afforded in addition.

The use of adrenaline, cocaine, or aromatic oils in sprays, or stramonium in the form of inhalations, powders, or cigarettes, is to be deprecated, as the relief afforded

is only temporary, and they have an extremely irritating effect on the bronchial mucous membrane; this, in turn, increases the liability to further attacks. They tend to change an asthmatic with acute attacks and perfectly free intervals into a chronic, constantly wheezing individual, though admittedly the spasms are less severe.

Intensive Treatment of the Asthmatic Sufferer

In order to prevent the recurrence of the asthmatic spasms two groups of factors need detailed consideration: first, the specific sensitizing substances peculiar to the individual case; and, secondly, a group of non-specific "catalysing" factors that tend to fire off any case.

Specific Sensitivity

In dealing with the specific sensitizing substance steps may be taken to ensure its removal or avoidance, or to reduce the patient's susceptibility to it either by specific or by non-specific means. In determining the specific cause a careful history, skin reactions, or elimination diets play a most important part. Here a few generalizations may be of assistance. Seldom does a food play an important part as a cause of asthma unless the spasms had their onset during the first decade, or are accompanied by skin lesions. In other words, a pure uncomplicated case of asthma commencing after childhood is most likely due to sensitization to an inhalant substance. In a non-seasonal case the four commonest causes are feathers, dust, animal hair, and orris root; while in seasonal cases pollens may also be considered. Bacterial allergy plays very little part in the majority of patients; even in cases always commencing with a cold the bacteria merely lower the resistance of the mucous membranes of the respiratory tract and so make the absorption of the common inhalant sensitizing substances more readily brought about, especially if the patient is confined to the house or put to bed.

Most asthmatics, when properly tested with potent solutions by the intradermal method, will react to one or more of the feathers, dust, animal hair, and orris root tests; so as a general rule in any asthmatic it is wise to remove these as far as possible. Feather or down pillows, eiderdowns or bolsters, feather, horsehair, or rabbit hair mattresses, should be replaced by a rubber mattress manufactured by the Dunlop Rubber Company, and by the best Java kapok pillows, bolster, and eiderdown. Both may be obtained from S. Brown and Sons, 10, Houndsditch, E.C.3, to fit any sized bed. Orris-root-free cosmetics may be obtained from Boutalls, Ltd., 150, Southampton Row, W.C.1. In avoiding dust a minimum of ledges and furniture in the bedroom is essential, and electric vacuum cleaning is advisable. Occupational hypersensitiveness can be avoided only by a complete change of occupation.

Desensitization

If all these factors cannot be removed or avoided, desensitization is necessary. In such a case it is better to employ a solution of these four common irritants, as one or more of them will be specific for the case, while the others will have a non-specific effect. By using concentrated solutions of the four inhalant substances, and preventing any general reaction by the simultaneous injection of 3 to 4 minims of adrenaline chloride (1 in 1,000 solution) in the earlier doses, it is possible to desensitize the patient very successfully and rapidly. This solution, together with directions as to dosage and technique, may be obtained from C. L. Bencard (1934) Ltd., 152, Great Portland Street, W.1. This specific solution is far more efficacious in the majority of cases of asthma than any of the previous non-specific methods such as peptone, tuberculin, milk, or vaccines.

Diet in Asthma

Food may affect an asthmatic in one of four ways. First, it may contain an article to which the patient is specifically sensitive at all times. Secondly, the patient may be specifically sensitive to the food only when his power of digestion is impaired: quite a large number of asthmatics, and especially children, who are more commonly food-sensitive than adults, show a diminished secretion of acid gastric juice or evidences of mucous gastritis, and so improperly digested foods gain easier access to the blood stream and hence produce reactions. Thirdly, a gross attack of indigestion may make the patient less resistant to some specific substance in the diet. Fourthly, the stomach and upper intestinal tract are supplied by the vagus nerve, so gastric distension may irritate the vagus and so set up a reflex bronchial spasm.

A basic diet avoiding all the commoner specific causes of allergic reactions has been published previously (*British Medical Journal*, 1933, ii, 43). As the commonest cause of indigestion in an asthmatic is a deficient secretion of hydrochloric acid, this should be rectified. One aklorep tablet (Roberts and Co., 76, New Bond Street), containing the equivalent of 20 minims of dilute hydrochloric acid, may be given immediately before each meal, swallowed with a wineglassful of water or orange juice. To prevent gastric distension three small, easily digested, simply compounded meals a day are best. A good early breakfast, a substantial hot midday meat meal, and a light early supper several hours before retiring should be the rule. No food or drink should be taken after supper, and especially not just before or after getting into bed.

Medicines

As regards medicaments, the chronic asthmatic needs a strong antispasmodic and sedative linctus at bedtime, such as tincture of stramonium 15 minims, syrup of Virginian prune 1 drachm, syrup of codeine phosphate 1 drachm. One dose should be taken every night without fail for several months, until the patient has been free from asthma for three months at least. The majority of asthmatics do not require any medicines during the day, and while glucose helps from an energizing point of view calcium in any form is of no proven value.

Non-specific Factors

The non-specific "catalysers" that help to fire off the asthmatic spasm form a very motley group. Whilst the general tendency is for the body to overcome its inclination to react with the specific sensitizing substance, its resistance may be broken down at any time by a group of non-allergic factors of a psychic, endocrine, nasal, toxic, dietary, or environmental nature. These factors explain why an asthmatic attack comes on with a cold, during states of emotion or excitement, following dietary indiscretions, or in certain atmospheres. Diet and bacterial infections have been touched upon already. Regarding nasal abnormalities, most recent observations prove that they are only coincidental or directly secondary to nasal allergic manifestations; so treatment should be conservative and non-surgical along allergic lines until as good results as possible are obtained. Psychic stimuli can only cause asthmatic attacks in a patient who is already asthmatic: in other words, as a result of his allergy the onset of the spasm may become a conditioned reflex, which, after repeated exposures, is capable of being fired off solely through nervous channels without the intervention of the specific sensitizing substance. When, as the result of desensitization, the patient is shown that he can experience exposure to the original cause without reaction all the psychic elements disappear. Endocrine dysfunction is rare in definite cases of allergy, yet

menstruation, marriage, and the menopause play an important part in women.

Week-end asthma, especially in school children, is due largely to the hot bath and syrup of figs on a Friday night, after which the child comes into immediate contact with his specific sensitizing substance—probably a feather or down pillow. Hence it is wise to insist on a morning bath and a morning saline. Early morning breathing exercises, especially of an expiratory nature (counting, humming, hissing, or blowing), are very beneficial if not continued up to the point of fatigue. As regards climate, the sufferer should be shielded from extremes of temperature, draughts, cold wind, and fogs.

Nova et Vetera

RECOVERY AFTER JUDICIAL HANGING

There must have been good reasons for the introduction of the "long drop" instead of the ancient method of hanging by the neck "until he be dead." It was only in 1874 that this new mode of execution was introduced by Marwood, the hangman. His predecessor, Calcraft, was said to have been very rough in his method of hanging, and doubtless many accidents had occurred during his tenure of office. One reason for this change over must have been the numerous cases of recovery after the offender had been "turned off the ladder." This form of capital punishment was really death by asphyxiation, and it is well known that many people have not only escaped from such a death, but certain individuals have made it a means of livelihood by giving exhibitions of being "strung up."

So often was death slow in coming to relieve the unfortunate one of life, that the executioner was in the habit of adding his weight to that of the victim by hanging on to his legs or by lifting the body up and then jerking it down. In these efforts the hangman was often assisted by the friends of the prisoner or by the interested spectators. The hangman, in spite of the wide experience which the law afforded him, was often ignorant and uncouth and, as well might be expected, under the influence of drink, with the result that the rope not infrequently slipped off or broke, and the unfortunate had to suffer all the horrors of a fresh suspension. How could such accidents fail to happen when processions to Tyburn took place almost every day, or when batches of ten or twelve prisoners were executed together at the Old Bailey? The notorious pirate Captain Kidd had to be hanged twice before he died, and the same fate befell another pirate, John Gow.

That there were many cases of actual recovery after the legal process of hanging had been carried out is indubitable. There is the case of the hanged woman who brought fame to the surgeon and diplomat, Sir William Petty, in the reign of Queen Anne. The body was lying in the anatomy theatre in Oxford awaiting dissection, when Sir William, noticing signs of life, bled her, and put her to bed along with another woman; later, spirits and cordials were administered, with the result that the woman revived and continued to live for some fifteen years afterwards. In 1658 a female servant was hanged in the same town, and, to make sure that she was dead, the body was allowed to hang an unusually long time. Yet she revived after being taken down, but only to be hanged by the neck until she was dead the following day.

A man bearing the very ordinary name of Smith was condemned to death in 1705. A reprieve arrived unfortunately just after he had been hanged. He was taken down, bled, and revived. He described his feelings at the moment of death as if his "spirits" forced their way up to his head and seemed to go out at his eyes with a great blaze of light, and then all pain left him. On his resuscitation the blood "and spirits forcing themselves into their proper channels gave him such intolerable suffering that he could have wished those hanged who

cut him down." William Duell was hanged in 1740, but when the body was laid out for dissection a man who was washing it noted signs of life. Duell was bled, and in two hours he was able to sit up in a chair. He was taken back to Newgate, and as an act of clemency was transported instead of being hanged again. Cases of recovery after hanging became so well known that it was a common practice for the friends of the hanged one to carry the body at once to a surgeon, who by bleeding and by other means made an endeavour to bring back life. This was done to the body of Dr. Dodd, who was hanged for forgery in 1777, but the effort was vain.

In some cases recovery occurred during the conveyance of the body in a cart. The roads were very rough, and there being no springs to the wagon, a sort of artificial respiration took place, with the result that in some instances the hanged person revived. The Edinburgh case of "half-hangit Meg" is well known locally. She was a married woman who, convicted of the crime of concealment of pregnancy, was hanged in 1724. A broadside of the time gives in grim detail an account of the hanging: how the hangman carried out his "usual office" of dragging down her legs, how the body after hanging the allotted time was put in a coffin, and how, after a scuffle with some apprentice surgeons who wished to secure the body for the anatomists, the friends of Margaret Dickson succeeded in placing the coffin in a cart. The sad journey to Musselburgh was interrupted at Peffermill by Meg beginning to show some signs of returning life. By the time that Musselburgh was reached she was almost well. Crowds of folk flocked to see such a wonder. She lived to become the mother of several children, kept an ale-house, and sold salt in the streets of the capital for many years after her suspension in the Grassmarket. I find it stated that jolting in a cart became so well known as a means of resuscitation that it was carried out in Ireland in almost every case subsequent to hanging. It seems to have been the custom there for the friends of the prisoner to hold him up by the waistband so as to prevent the weight of the body drawing the rope tight, the sheriff at the same time finding something to distract his attention.

At the beginning of last century a soldier was executed in Jersey, but unsuccessfully from a legal point of view. As death did not occur sufficiently quickly to satisfy the hangman, he hung on to the soldier's legs. This, however, did not seem to hasten matters, so he got upon the shoulders of the man, with the result that the rope broke. "To the great surprise of all who witnessed the dreadful scene, the poor criminal rose straight upon his feet, with the hangman on his shoulders, and immediately loosened the rope with his fingers." The sheriffs sent for another rope, but the spectators interfered, and the man was sent back to prison and, later, received pardon. Not long after this occurrence, the ropes broke in the case of two offenders who were being executed at Chester. In the afternoon, however, new ropes having been procured, they were successfully "turned off." William Snow, while being executed at Exeter, slipped from the gallows owing to the carelessness of the hangman and fell to the ground. He rose to his feet and, addressing the crowd, said, "Good people, do not be hurried; I am not, I can wait."

Recently I came across the following interesting account of the case of recovery at Oxford mentioned above:

News From The Dead: In the Revival of Anne Greene, Hanged at Oxford in 1650 with near 40 Ingenious Poems on the subject in Latin, English, etc. by the Prime Wits of that University. Phoenix Britannicus. Vol. I. 1732.

That none may be deceived and that so signal an Act of God's Mercy and Providence may never be forgotten, I have here faithfully recorded it.

In the House of Sir Thomas Read at Dunstew in Oxfordshire, there lived a Maid named Anne Greene; being about twenty-two years of Age, of a middle stature, strong, fleshy, and of an indifferent good Feature. She was seduced by Mr. Jeffery Read, Grandchild to the said Sir Thomas, and was delivered of a Man Child which she was alleged to have murdered. In an extreme cold and rainy Day, she was sent to Oxford Jail; where having passed three Weeks in continual