Conclusion

The results of thoracic surgery are most encouraging and the work must be supported, but there is no need to be carried away in our enthusiasm to provide surplus possibilities for the treatment of one case while there are not adequate facilities for treatment of the ninety-nine others.

In conclusion, I wish to state that the above views are my own personal opinions, and have no bearing on, or connexion with, the scheme or work of any public authority, and that I have brought them forward merely with a view to promote discussion on what I consider an interesting and important problem.

REFERENCES

¹ *Brompton Hospital Reports*, 1933, ii.

² *Public Health*, No. 1, xlviii, 37.

³ Ellman, P., and Tecton: *Architectural Journal*, January 18th, 1933.

A REVIEW OF ONE HUNDRED CONSECUTIVE GALL-BLADDER OPERATIONS

BY

J. MURRAY BLACK, M.B., F.R.C.S.ED.

HONORARY SURGEON, DUNFERMLINE AND WEST FIFE HOSPITAL

The cases under review have all been operated on by the writer, and sufficient time has elapsed since operation to enable one to judge from a clinical standpoint the success or otherwise of the operative procedure.

In Table I the cases have been divided into four groups. Included in the first group are cases of empyema of the gall-bladder and tension gangrene of the gall-bladder, so that the term "acute" denotes the urgency of the condition rather than its duration, since many of the acute cases were associated with, or superimposed on, a chronic condition. The other three groups are self-explanatory.

TABLE I

Type	No. of Cases	Sex		No. of Deaths
		Male	Female	
1. Acute	14	5	9	3
2. Chronic cholecystitis with stones	50	4	46	6
3. Chronic cholecystitis without stones	34	6	28	0
4. Carcinoma of gall-bladder ...	2	1	1	1
Totals	100	16	84	10

Sex Incidence, Age, and Habitus of Patient

In this series 84 per cent. of the patients were females, and of these fifty-three had borne children. This corresponds with what is generally known. The average age of the eighty-four female patients was 49 years, while that of the sixteen males was 52 years. Of the whole series forty-six of the patients were of the asthenic type of bodily habitus, and fifty-four of the sthenic type.

Symptoms and Associated Lesions

In eighty-five of the cases there was a history of flatulent indigestion; thirty-three cases had suffered from attacks of biliary colic; and in thirty-one of the cases jaundice had been present either at the time of treatment or on some previous occasion. Fifteen of the patients had definite symptoms or signs of "cholecystitic heart"—namely, dyspnoea on exertion, cyanosis, oedema of the ankles, or bronchitis. Four cases suffered from attacks

of bronchial asthma. Pyelitis was an accompaniment in sixteen of the cases, and was due to *B. coli* in every instance. Duodenal ulcer was present in addition to the gall-bladder lesion in four cases. Chronic appendicitis was present in ten instances, and in one case hour-glass stomach accompanied the gall-bladder infection.

Type of Operation and Percentage Mortality

Cholecystostomy was performed in the fourteen acute cases; in the two cases of carcinoma of the gall-bladder cholecystectomy was the operation carried out; and in the eighty-four chronic cases the choice of operation performed is shown in Table II.

TABLE II

Type of Operation	No. of Cases	No. of Deaths
1. Cholecystectomy	56	4
2. Cholecystectomy with drainage of the common bile duct	3	1
3. Cholecystostomy	25	1
Totals	84	6

In the acute cases there were three deaths out of fourteen, a mortality of 21 per cent. The cause of death in two cases was peritonitis, and in the third myocarditis. All three patients were over 68 years old. In the eighty-four chronic cases the mortality rate was 7 per cent. In two cases the cause of death was a technical error—namely, ligation of a crushed cystic duct with necrosis at the site of the ligature and extravasation of bile into the peritoneal cavity on the eighth and tenth days after operation respectively. Pulmonary embolus accounted for another fatality, and subphrenic abscess for a fourth. The remaining two deaths occurred in patients over 70 years of age, and were due to myocarditis and bronchitis respectively.

Pathology and Late Results in the Chronic Cases

In fifty of the eighty-four chronic cases stones were present, and in these the gall-bladder, if removed, was not examined microscopically, stone formation being sufficient evidence of a diseased organ. In the remaining thirty-four cases in which the gall-bladder was removed histological examination was carried out, and the results are tabulated below.

TABLE III

Gall-stones {	single	...	...	...	13 Cases
	multiple	...	...	...	37 "
					50
Chronic cholecystitis					22 Cases
Cholesterosis and fibrosis					12 "

Associated liver changes were noted at the time of operation in forty-six cases. The liver showed "whitening" in the region of the gall-bladder, and this discoloration was of value in determining cholecystectomy when the gall-bladder did not appear to be grossly diseased to naked-eye inspection. (E. S. Judd, writing in the *Mayo Clinic Collection* of 1930, states: "Hepatitis may occur as a primary condition, and the symptoms which result from it are similar to those of cholecystitis. Furthermore, removal of the gall-bladder in these cases of primary hepatitis will relieve the symptoms.")

There is still a difference of opinion regarding the effects on the health of the patient of removal of the gall-bladder. This review was undertaken with the aim of finding out the late clinical after-results of cholecystectomy as compared with those after cholecystostomy. The recurrence of previous symptoms, the onset of new

symptoms following operation, and the general health of the subject at a period at least three years after the operation were investigated, and the results are shown in Table IV.

TABLE IV

No. of Cases	Type of Operation	Health Satisfactory	Required Further Operation	Still Have Symptoms	Not Accounted For
59	Cholecystectomy	51 (86.4 %)	1 (1.8 %)	2 (3.3 %)	5 (8.4 %)
25	Cholecystostomy	10 (40 %)	6 (24 %)	7 (28 %)	2 (8 %)

The case requiring further operation after cholecystectomy is of interest, and a similar condition following cholecystostomy has been observed and reported on by Judd and White in the Mayo Clinic publication of 1928.

In my case, at the first operation, a thick-walled gall-bladder, containing stones, was removed; no stones were palpated in the common bile duct, which was not opened. Four months later the patient, a stout female of 54 years, had recurrence of severe attacks of pain with jaundice. It was surmised that she must have a stone in the common duct. At the second operation the common duct was opened and no stone was present, but the pancreas was hard, nodular, and enlarged. Drainage of the common duct was carried out by means of a Deaver's T-tube for about six weeks, her jaundice and pain disappeared, and she has now been free of symptoms for two years.

Of the 24 per cent. of cases requiring further operation after cholecystostomy five had stones in the gall-bladder at the second operation, and one developed acute pancreatitis, requiring redrainage of the gall-bladder.

Summary

Up till the time of Wilkie's experimental work there was room for performing cholecystostomy in the belief that drainage might cure the condition of the gall-bladder. With the realization that the gall-bladder infection is intramural this operation is now reserved for acute cases. C. H. Mayo states: "It may be wise to preserve and drain the gall-bladder in cases of acute illness." A further indication for cholecystostomy may be the presence of thickening and enlargement of the pancreas where prolonged drainage of the gall-bladder is necessary.

Judging from the clinical results in this small series one has no hesitation in concluding that cholecystectomy is the operation of choice.

The report for 1933 of the Malaria Advisory Board, published as a supplement to the *Federated Malay States Government Gazette*, announces the issue of an up-to-date guide to the identification of anopheline adults and larvae. It is stated also that bamboo subsoil drainage had been tried for breeding-places in seepages from a banked-up pool. The bamboos were laid in a semi-green condition, and it was expected that they would last from two to three years, but even if they had had to be replaced yearly the cost of this would have been less than that of oiling. It is suggested that this method should therefore be particularly useful in the more remote places, where the skilled supervision required for oiling is not available. In view of the present financial difficulties more economical means of oiling had been carefully discussed, but where only a minimum of oil is used grave danger is to be anticipated unless constant, skilled, and intimate supervision is supplied. A modification of Dr. W. T. Quaife's brush method is being used throughout the Kuala Lumpur oiling area. This method is said to have the following advantages over ten-day oiling and older methods: equal or increased safety; much lower cost; thorough mixing of oil, water, and larvae; drains automatically swept clear of obstructions; and minimum damage done to the sides. An investigation showed that antimalarial oiling presented no particular danger to poultry.

PREVENTIVE IMMUNIZATION IN AN INSTITUTIONAL EPIDEMIC OF SCARLET FEVER

BY

F. L. SCOTT, M.R.C.S., L.R.C.P., D.P.M.

SENIOR MEDICAL OFFICER

AND

W. MORTON, M.B., CH.B.

ASSISTANT MEDICAL OFFICER

BRACEBRIDGE MENTAL HOSPITAL, NEAR LINCOLN

During the early part of 1934 an outbreak of scarlet fever occurred at the Bracebridge Mental Hospital affecting the female patients and nursing staff. The fever was of an unusually mild type, but none the less serious from the point of view of complications and infectivity. The first case, that of a nurse, occurred on January 27th, and during the ensuing six weeks sixteen other nurses and twenty-six patients contracted the disease. During the same period there were also twelve cases of septic tonsillitis and numerous others of a follicular type.

Type of Case and Complications

The majority of cases were generally atypical in character, the temperature seldom rising above 100° F., though the pulse rate was comparatively high; circumoral pallor was rarely seen, the tongue was only lightly coated, and palatine spots were absent. The rash was typical but usually transient, and was followed by normal desquamation. There was little constitutional disturbance, and of the thirty-one cases treated here only four developed any albuminuria (eleven nurses were treated by the public health authorities). So mild was the usual case that we should often have hesitated to diagnose scarlet fever had we not known that such a type was prevalent in the neighbourhood.

Otitis media occurred in two cases, multiple arthritis in seven, and bronchopneumonia in one case, which was fatal in a patient aged 71. Nephritis was not a complication in any case.

TABLE I.—Showing Age Incidence

	Up to 20	21 to 25	25 to 30	31 to 35	36 to 40	41 to 45	46 to 50	50 and Over	Total
Female patients	2	3	1	2	4	5	4	5	26
Female nurses	6	9	2	0	0	0	0	0	17

It will be observed that fourteen of the twenty-six patients affected were over the age of 40, two being 71 and 76 respectively, but it must be remembered that the average age of the patient population is comparatively high.

Preventive Measures

Cases were isolated as soon as diagnosed, but in an institution with 200 staff and 1,350 patients, many of whom are employed in different parts of the building, isolation of all contacts was found impracticable. It was decided, therefore, to immunize all susceptibles as demonstrated by the Dick test, the test being applied to the whole female staff and to all direct patient contacts. In order to save valuable time, however, ninety-two patients working in the kitchen and laundry were immunized without preliminary testing.

Immunization was effected by giving a course of four weekly intramuscular injections of scarlet fever prophylactic toxin (Burroughs Wellcome and Co.). This method was used in preference to the injection of one massive