There are cases both of the inflammatory and of the degenerative types in which a spasmodic element may be so definite as to lead to difficulty in an accurate diagnosis, and in eventualities of this kind the tests which I have mentioned are of such value that I think it is obligatory to employ one or other of them before assuming that we can afford these cases some measure of relief by procedures designed to secure vaso-dilatation.

Treatment, apart from amputation, is frankly discouraging. Vaso-dilators of a chemical nature, such as muscle extract, peptone, and acetylcholine, have but a temporary value, and that only when the spasmodic element exists. The alternate suction and pressure treatment of Landis and Gibbon has not been employed in this country to any extent so far as I am aware, but promising results have been claimed by these authors in cases of thrombo-angiitis. Sympathectomy, which at one time appeared to offer a solution, has failed to yield the results which were hoped for. I do not say that it is valueless—by no means; it increases the degree of comfort, it lessens the pain, and it probably delays the onset of skin ulceration and gangrene—but it cannot be said to have any permanent effect upon the progress of the disease. And when we consider the pathology which underlies these lesions it could not very well be otherwise, for the vasospasm which sympathectomy relieves is but an incidental and often a minor part of the pathological picture.

RELATIONSHIP OF JOCHMANN'S AND OTHER ENZYMES TO THE ENCEPHALITOGENIC AGENT IN LYMPHADENOMATOUS LYMPHATIC GLANDS

BY

IAN MACKENZIE, M.B., CH.B.ED.

DAVIDSON RESEARCH FELLOW IN APPLIED BACTERIOLOGY, EDINBURGH UNIVERSITY; ASSISTANT PATHOLOGIST, ROYAL INFIRMARY, EDINBURGH

AND

C. E. VAN ROOYEN, M.D.ED.*

HALLEY STEWART RESEARCH FELLOW, EDINBURGH UNIVERSITY DEPARTMENT OF BACTERIOLOGY

(From the Department of Bacteriology, University of Edinburgh)

It has been shown by Gordon (1932-3) and van Rooyen (1933-4) that the intracerebral inoculation of rabbits with emulsions of lymphadenomatous tissue is followed by highly characteristic nervous lesions in the animal. Thus, after a definite incubation period lasting from three to four days, the animal develops signs of ataxia, incoordination, muscular spasms, head retraction, and generalized encephalomyelitis with usually a fatal result. A number of animals, however, that are affected to a lesser degree pass into a chronic state of disease, in which progressive atrophy of muscles and general emaciation are the principal features.

Such effects as these cannot be produced by the intracerebral inoculation of lymph glands affected with lymphosarcoma, leukaemia, carcinoma, tuberculosis, or syphilis, and thus the phenomenon has served as a biological test for the diagnosis of Hodgkin's disease. The value of the test has been adequately demonstrated by several workers (see van Rooyen, 1933-4; Ogilvie and van Rooyen, 1933-4; Gow, 1934; and van der Hoeden and Hulst, 1934), and consequently no further reference need be made to this aspect of the subject in the present paper.

Although the encephalitogenic effect in the rabbit of lymphadenomatous tissue appears to be a well-established fact, considerable dubiety exists with regard to the nature

of the pathogenic agent present in this tissue. Gordon (1933) has suggested that the behaviour of this agent is analogous to that of certain viruses in man and animals. Van Rooyen (1934) has concluded that the effects produced in the rabbit resemble more those of toxic action than of a virus infection, and has also pointed out that inability to transmit the condition from rabbit to rabbit, and the absence of intracellular inclusion bodies and of any immunity reaction, renders it difficult to assign this encephalitogenic agent to the infective viruses.

Further light has been thrown on the problem by the work of Friedemann (1934), who showed that the effects in the rabbit of intracerebral inoculation of lymphadenomatous tissue may also be produced by the inoculation of emulsions of normal splenic tissue and bone marrow, and of normal leucocytes. He showed that similar materials derived from cases of acute leukaemia and pernicious anaemia may likewise yield the characteristic effects. The encephalitic reaction has been found to be particularly marked with a specimen of bone marrow derived from a case of acute myelogenous leukaemia in which there were also chloromatous changes (van Rooyen, 1934).

The question has therefore arisen whether the agent present in lymphadenomatous tissue is identical with that present in these tissues, and whether, as Friedemann points out, both agents can be identified with the proteolytic enzyme extracted by Jochmann and Lockemann (1908) from leucocytes, splenic tissue, and bone marrow.

Thus it has been clearly shown that the characteristic encephalitic syndrome can be produced by the intracerebral inoculation of a product derived from normal tissues and cells; but, on the other hand, we have obtained evidence to show that the encephalitogenic agent in lymphadenomatous glands is not identical with Jochmann's enzyme, although both of them exhibit the same degree of resistance towards certain physical and chemical agents. We have also ascertained whether certain bacteria and vaccinia virus can withstand these agents.

The extraction and properties of the enzyme have been fully described by Jochmann and Lockemann (1908), and Jochmann (1913), and further reference has been made to its presence in tissues by Friedemann (1934). We have carried out a number of experiments with tissues extracted according to the method adopted by these workers for separating the ferment, and the extracts have been carefully examined for proteolytic properties and their ability to produce pathogenic effects when inoculated into the brain of the rabbit.

Technique for Extraction of Jochmann's Enzyme

We have employed the same method as was adopted by Friedemann (1934) for this purpose. It is as follows.

Tissue is weighed, placed in an oven at 55° C. for forty-eight hours, and 5 volumes of a mixture consisting of 2 volumes of absolute alcohol to 1 volume of ether added. After twenty-four hours at room temperature the supernatant fluid is discarded and the residue desiccated to dryness *in vacuo*. The desiccate is thereafter triturated in a 50 per cent. aqueous solution of glycerol, allowed to stand for twenty-four hours at room temperature, centrifuged at 2,500 revolutions per minute for twelve minutes, and the supernatant fluid withdrawn with a sterile pipette and added to an alcohol-ether mixture of the previous proportions. After twenty-four hours at room temperature the precipitate is removed by centrifugalization, resuspended in 2 to 3 c.cm. of 0.86 per cent. saline, and used for injection.

The alternative method (advocated by Friedemann, 1934) for the extraction of the enzyme by the use of acetone and filter paper was also tried, and found to give similar results. Unextracted tissue emulsions were also used for inoculation experiments; these constituted 1 in 10 suspensions obtained by

* By aid of grants received from the Science Committee of the British Medical Association and the Medical Research Council.

finely mincing and grinding material in broth of pH 7.1 and thereafter allowed to stand for seven days in the ice-chest prior to injection. As a precautionary measure against bacterial contamination those tissues which had been obtained at necropsy were initially subjected to the superficial application of heat and alcohol and then triturated in 0.5 per cent. phenol broth according to the method described by van Rooyen (1933). In some cases the broth emulsions were also heated at 60° C. for sixty minutes to exclude all possibility of bacterial contamination.

The technique used for intracerebral inoculation of rabbits was similar to that described by van Rooyen (1933), and, stated briefly, consisted of injecting 0.4 c.cm. of tissue emulsion into the occipital lobe of the animal to a depth of 3 mm. from the skin surface. The site chosen for perforating the skull was approximately 2 mm. equidistant from the right (or left) lambdoidal suture postero-laterally and the sagittal suture medially.

Interpretation of Results

Only those animals which exhibited the characteristic syndrome described by Gordon (1932) have been regarded as yielding positive results. The features of this condition are well known and require no further description, but we would emphasize that the incubation period of three to four days prior to the development of the syndrome and the absence of any cultivable bacterium in the brain of such an affected animal are perhaps the most striking features of the encephalitic condition. We would also like to place on record that "spasmodic contractions of the muscles of the neck, distorted position of the head, frequent rotations around the body axis" (Friedemann and Elkeles, 1933) are not a characteristic feature of Gordon's syndrome. In our experience, gleaned from the results of intracerebral inoculation in over 1,000 rabbits, such signs are the outcome of brain injury.

Effect of the Jochmann-Lockemann Method Applied to Histologically Typical Glands

Twelve enlarged lymphatic glands removed from cases of clinically and histologically characteristic Hodgkin's disease, which had given a positive reaction on intracerebral inoculation, were treated by the Jochmann-Lockemann method and then tested in the same way. On every occasion the product obtained after extraction produced the same effect in the rabbit as was noted with freshly emulsified suspensions of similar tissue. In the case of two freshly removed glands which yielded a weakly positive effect when emulsified in broth, an appreciably greater reaction in the animal was elicited after extraction. But the feebly enhanced effects of concentration by this method were not as great as that obtained with broth emulsions prepared from similar glands which had been first subjected to desiccation. The experiments have, however, demonstrated that the encephalitogenic agent is capable of withstanding consecutively 55° C. for forty-eight hours, alcohol (2 volumes) and ether (1 volume) for forty-eight hours, and 50 per cent. glycerol for twenty-four hours' duration.

EXPERIMENTS ON LYMPH GLANDS FROM TYPICAL CASES YIELDING A NEGATIVE BIOLOGICAL TEST

Elsewhere van Rooyen (1933) has pointed out that this test was negative in 25 per cent. of cases of clinically and histologically characteristic Hodgkin's disease. It was, moreover, suggested that the absence of the reaction in such cases may possibly have been due to the effects of repeated irradiation or excessive fibrosis of the glands examined. Further work has been done on this aspect of the subject, and an attempt made to ascertain whether the extracts obtained from negatively reacting glands by the Jochmann-Lockemann procedure may yield positive reactions.

Six enlarged Hodgkin lymph nodes of over 2 mg. in weight, which had previously given a negative reaction,

were extracted by the Jochmann-Lockemann technique, and the extract was injected intracerebrally into rabbits. Fifteen rabbits were injected with such emulsions, and the dosage varied from 0.5 to 1 c.cm. in size, but, in spite of the employment of such comparatively large amounts, no effects were produced. These results have indicated that the preliminary application of the Jochmann-Lockemann technique to lymphatic glands giving a negative biological test fails to extract an encephalitogenic product from them.

Examination of Lymphatic Glands affected with Lymphadenoma and other Pathological Conditions

In order to ascertain whether Gordon's syndrome is due to Jochmann's proteolytic enzyme, we have investigated a number of lymphatic glands affected with a variety of different diseases, including Hodgkin's disease, for the presence of this enzyme. Thus twenty-four enlarged lymphatic glands have been examined for proteolytic action on coagulated serum (see Müller and Jochmann, 1906) in gelatin and litmus milk, and the results tabulated as follows.

TECHNIQUE

This test consisted in placing a large loopful of the tissue extract (vide supra) on the surface of a Petri dish containing Loeffler's coagulated serum medium which was incubated for forty-eight hours at 55° C., and then examined for liquefaction. Positive results were signified by the formation of a crateriform area of depression with liquefaction at the site of addition to the medium (see Müller and Jochmann, 1906).

To 5 c.cm. of litmus milk medium 1 to 2 c.cm. of the tissue extract was added, which was then incubated at 55° C. for three days. In the case of post-mortem material

TABLE I.—*Examination of Lymphatic Glands for Jochmann's Enzyme correlated with Pathogenic Effect on Intracerebral Inoculation of Tissue Emulsions and Tissue Extracts*

Case	Histology	Result of Intracerebral Inoculation of		Proteolytic Action of Extracts on	
		Broth Emulsion (Gordon's syndrome)	Extract prepared by Jochmann-Lockemann Method	Loeffler's Serum	Litmus Milk
H 12	Hodgkin's disease	Positive	Positive	No liquefaction	No change
H 18	" "	"	"	"	"
G	" "	"	"	"	"
H 40	" "	Strongly positive	Negative	"	"
E	" "	Negative	"	"	"
J	" "	Positive	Positive	"	Slight acidity
NH 2	" "	Negative	Negative	"	No change
NH 3	" "	"	(?) Positive	"	"
F	Case clinically resembling lymphadenoma but unconfirmed on histological examination, and no diagnosis given	"	Negative	"	"
C		"	"	"	"
D		"	"	"	Acid with digestion of milk
G 2		"	"	"	No change
NH 7	" "	"	"	"	"
LL 2	Lymphatic leukaemia	"	"	"	"
LL 3	" "	"	"	"	"
LL 4	" "	"	"	"	"
NH 8	Secondary carcinoma	"	"	"	"
I	" "	"	"	Large area of liquefaction	Acid with digestion of milk
C	Undifferentiated carcinoma	"	"	"	Slight acidity
H	Casating tuberculosis	"	"	"	No change
F	" "	"	"	"	"
B	Chronic tuberculosis	"	"	"	"

TABLE II.—Results obtained with Spleen and Bone Marrow

Case	Pathological Condition	Tissue	Intracerebral Inoculation of Broth Extract	Intracerebral Inoculation of Jochmann Extract	Proteolytic Action of Jochmann Extract on		
					Loeffler's Serum	Litmus Milk	Gelatin
B (W.G.H.)	Myelogenous leukaemia	Bone marrow	Positive	Negative	Liquefaction	No action	—
B (W.G.H.)	" "	Spleen	"	Positive	No liquefaction	"	—
LL 1	Lymphatic leukaemia	Bone marrow	"	Negative	"	"	—
LL 2	" "	"	Negative	"	Liquefaction	"	—
LL 2	" "	Spleen	Positive	"	Slight liquefaction	Marked activity	—
LL 3	" "	Bone marrow	Negative	"	No liquefaction	No action	—
LL 3	" "	Spleen	Positive	Positive	Active liquefaction	Marked activity	—
LL 4	" "	Bone marrow	Negative	Negative	No liquefaction	No action	—
LL 4	" "	Spleen	"	"	"	"	—
A	Removed at necropsy from patients dying of various conditions, one of which were blood diseases	"	Positive	Positive	Liquefaction	"	Negative in acid, neutral, and alkaline media
B		"	"	Negative	No liquefaction	"	
C		"	"	Positive	"	"	
D		"	"	"	"	"	

the presence of bacterial contamination was excluded by appropriate methods of examination as a precautionary measure. The slightest trace of redness in the indicator as compared with the uninoculated control was regarded as a positive result, and the appearance of marked acidity with digestion of the milk was interpreted as a strongly positive reaction. Then 0.5 c.cm. of tissue extract was added to a tube containing 2.5 c.cm. of gelatin, incubated at 55° C. for forty-eight hours, and allowed to stand for one day at room temperature (15° to 20° C.). The amount of liquefaction produced was measured by means of a calibrated pipette, and any bacterial growth was excluded by cultural examination. The results obtained were found to be identical with those noted in the case of coagulated serum, and consequently they have not been incorporated in the tables.

The findings in Table I indicate that Jochmann's enzyme is absent in lymphadenoma and the other conditions studied. They also show that the encephalitogenic action of Hodgkin lymph nodes cannot be identified with Jochmann's enzyme; thus several glands were found to be markedly encephalitogenic, but possessed no apparent proteolytic activity. From the results obtained with splenic and bone marrow tissue (Table II) it will be observed that though the broth extracts generally reproduced the encephalitic syndrome, the corresponding extracts obtained by the Jochmann-Lockemann method were by no means always active. Further analysis of the results reveals that there was no relation between proteolytic activity on the one hand and encephalitogenic effect on the other. From these experiments we have concluded that the encephalitogenic agent which occurs in lymphadenomatous lymphatic glands also occurs in the spleen and bone marrow.

TABLE III.—Pus Cells* and Normal Leucocytes

Case	Clinical Condition	Intracerebral Inoculation of Broth Extract	Intracerebral Inoculation of Jochmann's Extract	Proteolytic Action of Jochmann's Extract on	
				Loeffler's Serum	Litmus Milk
1	Pyogenic skin abscess	Positive	Negative	Active liquefaction	Very slight action
2	" "	"	"	Liquefaction	No action
3	" "	"	Positive	Active liquefaction	Slight action
4	" "	"	"	"	Marked action
5	" "	"	Negative	Liquefaction	"
6	" "	"	"	"	Very marked action
7	Normal leucocytes†	"	Positive	"	Marked action

* Sterilized by heating at 60° C. for sixty minutes.

† Obtained by differential centrifugalization of normal human blood which had been citrated.

Our experiments with human pus and leucocytes have confirmed the early work of Müller and Jochmann (1906), who drew attention to the marked proteolytic activity of polymorphonuclear leucocytes derived from normal blood and inflammatory exudates. It will again be observed, however, that there was no correlation between proteolytic activity and encephalitogenic effect; thus in four cases the Jochmann-Lockemann extracts were found to be non-pathogenic to the rabbit, but possessed definite proteolytic powers. Further, this method of extraction proved detrimental to the pathogenic agent, since only three out of seven cases yielded a positive result with Jochmann-Lockemann extracts, whereas positive results were obtained in all cases by the use of broth extracts. Similar results have been noted by Gordon (1934).

TABLE IV.—The Pancreas

Case	Intracerebral Inoculation of Broth Extract	Intracerebral Inoculation of Jochmann Extract	Proteolytic Activity of Jochmann Extract on	
			Loeffler's Serum	Litmus Milk
1	Negative	Negative	Marked liquefaction	Slight action
2	"	"	"	"
3	"	"	Liquefaction	Marked action
4	"	"	Marked liquefaction	"
5	"	"	Liquefaction	"
6	"	"	Marked liquefaction	"
	"	"	"	"

Müller and Jochmann (1906) state that "pancreas induces the quickest and by far the most intensive digestive action on Loeffler's serum at 55° C." In our experiments with human pancreatic tissue we found that, while the Jochmann-Lockemann extracts induced active proteolysis on Loeffler's serum and in litmus milk, in no instance did the characteristic syndrome described by Gordon appear in the rabbits injected with this material. On several occasions when injections of 0.6 c.cm. were given intracerebrally, the animals died either immediately or within forty-eight hours with signs of paralysis which rapidly developed within five hours of the injection. Reduction of the amount injected to 0.3 to 0.4 c.cm. gave uniformly negative results, the animals remaining unaffected by such dosages. A 1 in 10 concentration of pepsin solution (B.P.) in 0.2 per cent. hydrochloric acid, and in neutral solution, was also injected intracerebrally into rabbits, but no significant changes were observed.

In Vitro Effect of Human Tissue Extracts on Rabbit Brain Tissues

Friedemann (1934) has drawn attention to the lytic action of Jochmann-Lockemann extracts of spleen and bone marrow on emulsions of rabbit brain as a possible interpretation of the occurrence of Gordon's syndrome in the rabbit. We have investigated this action in the following manner.

The brains of several rabbits were removed aseptically and ground up in an equal quantity of normal saline to make a dense uniform emulsion; 4.5 c.cm. amounts of this emulsion, together with 0.5 c.cm. of extracts of various tissues obtained by the Jochmann-Lockemann method, were placed in sterile stoppered tubes and incubated at 55° C. for three days. The control consisted of 4.5 c.cm. brain emulsion with 0.5 c.cm. normal saline.

TABLE V.—In Vitro Effect of Tissue Extracts on Rabbit Brain Tissue

Material	Result of Intracerebral Injection of Extract in Rabbit	Effect on Brain Emulsion after		
		24 Hours	48 Hours	72 Hours
Gland affected with lymphadenoma	Positive	Nil	Nil	Nil
Gland affected with lymphosarcoma	Negative	"	"	"
" " "	"	"	"	"
Normal pancreas ...	"	Progressive liquefaction		
" " "	"	"	"	"
Control	—	Nil	Nil	Nil

In those tubes containing pancreatic extract there was progressive digestion, whereas in the tubes containing extracts of lymphadenomatous and lymphosarcomatous glands and in the control tube no such change had occurred after incubation. This result shows quite clearly that gross proteolytic action on the brain plays no part in the effect produced by Gordon's agent.

Experiments with Lymphoprotease and Leucoprotease

It has been stated by Jochmann, and subsequently alluded to by Friedemann, that Jochmann's proteolytic enzyme is also to be found in the spleen, bone marrow, and leucocytes of normal dogs. Opie (1905-6-7) has studied the proteolytic property of dog tissues in great detail, and has shown that such activity is attributable to the presence of two proteolytic enzymes. The first, derived from the polymorphonuclear leucocytes and acting in an alkaline or neutral medium, he has called "leucoprotease." The second, which he termed "lymphoprotease," is derived from the mononuclear macrophages, and acts in an acid medium.

We have been able to confirm these observations, and have investigated the possible encephalitogenic action of the two enzymes when injected intracerebrally into the rabbit.

TECHNIQUE

Opie's method of obtaining the two enzymes has been used, and is briefly as follows. A healthy dog was injected intrapleurally on each side with 10 c.cm. of a 10 per cent. suspension (by weight) of aleuronat in a 3 per cent. solution of starch. An inflammatory reaction ensued and a pleural exudate formed. After two days part of this exudate was withdrawn with needle and syringe. At this stage the cellular content of the exudate consisted mainly of polymorphonuclear leucocytes, and, as Opie has shown, contained the enzyme leucoprotease. At the end of six days from the date of inoculation the dog was killed and the pleural exudate removed. Microscopically its cellular content consisted of a high percentage of large mononuclear cells. The lymph glands adjacent to the site of inflammation—for example, the substernal glands—were markedly enlarged, and microscopical examination showed that the sinuses were

crowded with large mononuclear cells. The cells of the pleural exudates were separated by centrifugation, washed with normal saline, and finally suspended in nine times their volume of 0.85 per cent. saline. The lymph glands were thoroughly minced and triturated with three times their volume of 0.85 per cent. saline. All three suspensions were placed at 55° C. for twenty-four hours, and then tested for their proteolytic and encephalitogenic activity.

TABLE VI.—Experiments with Tissue Enzymes from Dogs

Nature of Specimen	Encephal- itogenic Action in Rabbit	Proteolytic Action on				Litmus Milk
		Loeffler's Serum	Gelatin			
			0.2 % Acetic Acid	Neutral	0.2 % Na ₂ CO ₃	
Pleural exudate after 2 days	Nil	+++	Complete liquefac- tion	Complete liquefac- tion	Complete liquefac- tion	—
Pleural exudate after 6 days	"	+++	"	"	Slight liquefac- tion	—
Glands... ..	"	++	"	"	Complete liquefac- tion	—
Bone marrow I	"	++	—	++	—	++
Bone marrow II	"	++	—	++	—	++
Normal leuco- cytes	"	++	—	++	—	++

The exudates were found to be innocuous when introduced into the brain of the rabbit. In the case of the gland emulsion, however, results were obtained similar to those described with extracts of human pancreas (vide supra). Thus intracerebral injection with 0.5 c.cm. of a 1 in 3 concentration of gland emulsion caused paralysis and death within twenty-four hours, but if the dose were diminished to 0.25 c.cm. the rabbit remained unaffected. These results show, therefore, that no relation exists between the encephalitogenic agent under discussion and the enzymes leucoprotease and lymphoprotease described by Opie. Likewise, emulsions of spleen and bone marrow, and extracts of these tissues obtained by the Jochmann-Lockemann method, were actively proteolytic, but non-pathogenic to the rabbit.

Resistance Exhibited by Certain Bacteria to the Jochmann-Lockemann Method of Tissue Extraction

It has been stated by Friedemann (1934) that the procedure adopted for the extraction of Jochmann's enzyme from tissues was adequately rigorous for the destruction of most forms of micro-organic life which might have been present in tissues before treatment by this method of extraction. We have investigated the precise degree of lethal effect exerted by such extraction when applied to a variety of different bacteria, and have noted that certain organisms resist such treatment.

TECHNIQUE

Aerobic five-day cultures of *B. proteus*, *B. pyocyaneus*, *B. anthracis*, *B. anthracoides*, *B. mycoides*, *Streptococcus faecalis*, *Staphylococcus aureus* and *albus*, together with anaerobic cultures of *B. tetanus*, *B. botulinus*, *B. sporogenes*, and *Vibrio septique* were treated as follows: placed in a paraffin oven at 55° C. for three days; centrifugalized at 3,000 revolutions per minute for thirty minutes; the supernatant fluid pipetted off; 15 c.cm. of a mixture composed of 2 volumes alcohol to 1 volume ether added, and allowed to stand at room temperature for twenty-four hours; thereafter tubes were recentrifugalized; the alcohol-ether mixture discarded and replaced with a 50 per cent. aqueous solution of glycerin, which was next allowed to act for a further twenty-four hours at room temperature, and then removed by differential centrifugalization. The deposited organisms were again subjected to a final treatment by alcohol and ether, and then resuspended in saline solution.

The presence of living organisms in each case was next ascertained by inoculation of blood-agar plates, which

were incubated under appropriate conditions and then examined for the presence of typical colonies. The following results were obtained: *B. anthracis*, *B. anthracoides*, and *B. mycoides*, abundant growth after thirty-six hours; *B. sporogenes* and *B. botulinus*, numerous colonies at the end of seventy-two hours; *B. tetani*, *Vibrio septique*, *Enterococcus*, *Staphylococcus aureus*, and *Staphylococcus albus*, a few colonies after the same length of time. It should be mentioned in connexion with experiments on cocci that dense bacterial emulsions suspended in the minimum quantity of saline were used. A considerable amount of drying thus occurred during heating at 55° C. for forty-eight hours, similar to that which resulted when tissues were placed under identical conditions. As far as could be ascertained, therefore, from our observations, the survival of staphylococci following exposure to the Jochmann-Lockemann method of extraction was dependent on the degree of drying to which these organisms were initially exposed. *B. proteus* and *B. pyocyaneus* failed to withstand the extraction process, and no growth was obtained after attempted cultivation. Likewise, similar results were obtained with an active strain of vaccinia virus, which was inactivated by this treatment.

These findings indicate that the method of Jochmann-Lockemann extraction is not necessarily lethal to organismal life, certain bacteria, both sporing and non-sporing, being able to survive the procedure.

Discussion

Three different hypotheses have been advanced in an attempt to explain the production of the encephalitic syndrome in rabbits following the intracerebral inoculation of extracts of certain human tissues. Gordon (1933-4) has suggested that the pathogenic agent belongs to the virus category; van Rooyen (1934) that it is possibly a neurotoxic substance with special affinity for rabbit brain tissue; while Friedemann (1934) concluded that it is identical with Jochmann's proteolytic enzyme. The last-mentioned worker arrived at this conclusion after a study of its distribution in the human body and of its resistance to certain chemical substances. This theory has formed the subject of our investigations.

F. Müller (1888), and later E. Müller and Jochmann (1906), noted that a proteolytic enzyme was present in human leucocytes, spleen, bone marrow, and pancreas, and in pus. Jochmann (1906) further demonstrated that it was present in these organs in the monkey, and to a lesser degree in the dog. They found that it could be effectively demonstrated by its ability to liquefy coagulated serum, a drop of the enzyme solution placed on Loeffler's serum medium producing a crateriform depression after incubation at 55° C. From their studies on the comparative proteolytic properties of white blood cells derived from cases of lymphatic and myelogenous leukaemia they came to the conclusion that in so far as leucocytes were concerned the proteolytic action was due to a ferment produced by the polymorphonuclear leucocytes.

The enzymes derived from leucocytes have been carefully studied by Opie (1905-6) in the dog. He found that two distinct enzymes were present. The first of them he names "leucoprotease," and states that it acts in a neutral or alkaline medium. This appears to be the enzyme described by Jochmann *et al.*, as it can be obtained from pus and withstands dehydration by alcohol and ether and extraction by glycerin. It resembles trypsin, but is less active. The other enzyme, which he calls "lymphoprotease," was obtained from the large mononuclear cells of lymph glands and inflammatory exudates—the macrophages of Metchnikoff. This enzyme, unlike leucoprotease, digests with greatest activity in an

acid medium, being almost entirely inactive in neutral or alkaline media. It differs from leucoprotease further in that it is less resistant to the action of heat; being largely inactivated by temperatures between 60° and 70° C., and in being destroyed by drying after treatment with alcohol and ether. Hedin and Rowland (1901) demonstrated the presence of proteolytic substances in the spleen of various animals (ox, sheep, horse, and the pig); and Hedin (1904) succeeded in isolating two proteolytic enzymes: one, which acted in an alkaline medium, he termed "lieno- α -protease"; and the other, which acted in an acid medium, he called "lieno- β -protease."

The results recorded in this paper indicate, in the first place, that the encephalitogenic action of extracted leucocytes and other tissues on the brain of the rabbit is not due to proteolytic action *per se*, as extracts of human pancreas which were actively proteolytic were incapable of reproducing the characteristic syndrome in the rabbit. This result with pancreatic extracts further indicates that a lipolytic ferment plays no part in the production of the syndrome.

The question then arose as to whether the pathogenic action in the rabbit was due to a specific proteolytic enzyme, associated with the cells of the various tissues exhibiting encephalitogenic properties. A study of the various enzymes described by Müller and Jochmann, Opie, and Hedin in these tissues led us to the conclusion that none of these enzymes could be regarded as Gordon's encephalitogenic agent. For example, various Jochmann-Lockemann extracts of Hodgkin lymph glands, which yielded a strongly positive encephalitogenic action, were apparently devoid of proteolytic activity according to the standards adopted. The same was true of several specimens of bone marrow and spleen when tested in alkaline, acid, and neutral media respectively. On the other hand, several Jochmann-Lockemann extracts of pus which were actively proteolytic contained no encephalitogenic agent. Likewise, in material derived from the normal organs and inflammatory exudates of dogs, proteolytic enzymes, active in acid, neutral, and alkaline media, were obtained; but these proved to be incapable of producing the typical syndrome in the rabbit.

We do not propose in this communication to enter into a discussion of the neutralizing properties of normal serum on the enzymes and pathogenic agent under discussion. We would draw attention to the fact, however, that whereas Gordon's agent, derived from lymphadenomatous glands, does not appear to be readily neutralizable, the anti-ferment action of normal serum is well recognized. Opie (1907) has further shown that leucoprotease is easily neutralizable by such serum. This aspect of the subject is in course of investigation, and will be referred to at a later date. The question of the vital nature of the encephalitogenic agent in the light of its resistance to a somewhat drastic mode of treatment has also been investigated. Comparative experiments with a number of different bacteria have led us to the conclusion that its resistance to the Jochmann-Lockemann procedure does not necessarily exclude the possibility of its being a living organism.

After full consideration of the foregoing experimental data we have concluded that Gordon's agent is not identical with Jochmann's enzyme. Neither is it identical with the several proteolytic enzymes described by Opie, Hedin, and their co-workers. Further, it is of importance to note that the property of leucocytes to produce encephalitis in the rabbit appears to be limited to the human subject among various animal species which we have examined. It is still impossible, however, to define this unique pathogenic principle, and further work might appropriately be directed towards obtaining a more exact definition of its true nature and mode of action.

Summary and Conclusions

1. Gordon's encephalitogenic agent originally derived from lymphadenomatous tissue may occur in human bone marrow, spleen, and leucocytes (whether normal or pathological), but not in those of the dog or the rabbit.

2. This agent is capable of withstanding the Jochmann-Lockemann procedure for the extraction of proteolytic enzymes from tissues.

3. Extracts of certain tissues obtained by this procedure may be highly pathogenic to the rabbit on intracerebral inoculation (encephalitogenic) but devoid of proteolytic action.

4. Conversely, extracts of certain tissues may be markedly proteolytic, containing the so-called Jochmann ferment, but quite non-pathogenic to the rabbit.

5. Gordon's pathogenic agent found in lymphatic glands affected with Hodgkin's disease cannot, therefore, be identified with Jochmann's proteolytic enzyme.

6. Lymphatic glands affected with a variety of different pathological conditions have been subjected to the Jochmann-Lockemann method of extraction, but it has not been found possible to derive an encephalitogenic product from these glands which initially gave a negative Gordon reaction.

7. Experiments with certain bacteria have demonstrated that this method of chemical extraction does not destroy all forms of living organisms.

8. The enzymes lymphoprotease and leucoprotease obtained from the dog, and also lieno- α -protease and lieno- β -protease, do not appear to be concerned in the production of the encephalitic syndrome following the intracerebral inoculation of tissue extracts.

We desire to acknowledge our indebtedness to Professor T. J. Mackie for much constructive criticism and valuable advice received during the course of the investigations, also to Dr. Mervyn H. Gordon of St. Bartholomew's Hospital, London, for the interest which he has taken in our work.

BIBLIOGRAPHY

- Friedemann, U., and Elkeles, A.: *British Medical Journal*, 1933, ii, 1110.
 Friedemann, U.: *Ibid.*, 1934, i, 517.
 Gordon, M. H.: *Rose Research on Lymphadenoma*, Bristol, 1932.
 Idem: *British Medical Journal*, 1933, i, 641.
 Idem: *Proc. Roy. Soc. Med.*, 1934, xxvii, 1035.
 Gow, A. E.: *Ibid.*, 1934, xxvii, 1045.
 Hedin, S. G., and Rowland, S.: *Zeit. f. Physiol. Chem.*, 1901, xxxii, 341.
 Hedin, S. G.: *Journ. of Physiol.*, 1904, xxx, 194.
 Jochmann, G.: *Kolle Wassermann, Handb. d. path. Microörg.*, vol. ii, Part 2, p. 1301.
 Jochmann, G., and Lockemann, G.: *Hofmeister's Beitr.*, 1908, ii, 449.
 Müller, F.: Apparently unpublished observations cited by Kossel, *Zeit. f. klin. Med.*, 1888, xiii, 149.
 Müller, E., and Jochmann, G.: *Munch. med. Woch.*, 1906, liii, 1393, 1507, 2002.
 Ogilvie, R. F., and van Rooyen, C. E.: *Lancet*, 1933, ii, 641.
 Idem: *Journ. Amer. Med. Assoc.*, 1934, cii, 1842.
 Opie, E. L.: *Journ. Exper. Med.*, 1905, vii, 316; *ibid.*, 1906, viii, 410.
 Idem: *Physiol. Rev.*, 1922, ii, 552.
 Opie, E. L., and Barker, B. I.: *Journ. Exper. Med.*, 1907, ix, 207.
 van der Hoeden, J., and Hulst, L. A.: *Nederl. Tijdschr. v. Geneesk.*, 1934, lxxviii, No. 38, 4305.
 van Rooyen, C. E.: *British Medical Journal*, 1933, i, 50; 1933, i, 664; 1933, ii, 562; 1934, i, 519.

L. W. Sauer (*Amer. Journ. Dis. Child.*, January, 1935, p. 69) has treated 650 non-immune infants and young children by a vaccine which differed from all other pertussis vaccines in being made every few months from recently isolated haemolytic strains of the Bordet-Gengou bacillus. The total dosage after 6 months of age was 8 c.cm. Local and systemic reactions were negligible. Thirty-two children used as controls in twenty-five of the families contracted undoubted pertussis, whereas 170 of the injected children in these families who had been exposed escaped. No claims are made for the vaccine as a therapeutic agent, nor when exposure has already occurred or threatens, as the immunization takes about four months to develop.

SEROLOGICAL DIAGNOSIS OF WEIL'S DISEASE

BY

H. C. BROWN, C.I.E., M.B., D.T.M. AND H. C. CAMP.
 MAJOR I.M.S. (RET.)

WELLCOME BUREAU OF SCIENTIFIC RESEARCH, LONDON

Considerable attention has recently been paid to certain outbreaks and sporadic cases of Weil's disease in the British Isles. As the laboratory diagnosis by serological methods is somewhat specialized, and an early and definite diagnosis is of extreme importance, a description of the procedures may possibly be of use to those wishing to carry out serological tests for this disease.

Agglutination Test

I shall only refer briefly to this test, which has been perfected by Professor Schüffner (1934) and his co-workers in Amsterdam. It consists essentially of allowing varying concentrations of the patient's serum to interact for two hours at 32° C. with an equal volume of a young culture of *L. icterohaemorrhagiae*. After incubation small drops from the various dilutions are placed on a slide and are examined, without a cover-slip, by dark-ground illumination.

One factor which may make the results difficult to read when using living cultures is that lysis of the leptospira occurs in certain dilutions, and the result is consequently masked. In order to overcome this difficulty Professor Schüffner has, for the last two years, been using formalized cultures; with these no lysis occurs, but there is a tendency for acid agglutination to take place unless the strength of the pure formalin in the suspension is reduced to 0.5 per cent.

According to Postmus (1933) a positive agglutination reaction may be expected on the sixth day of disease. The reaction reaches its maximum on the twentieth day, when the titre of the serum may rise to 1 in 100,000. Control sera in Schüffner's hands have not shown positive agglutination even at a 1 in 10 dilution. Residual agglutinins may persist for several years: Postmus has demonstrated their presence 6,066 days after the onset of the disease, and in one of the cases recorded by Fairley (1934) agglutinins were shown to be present twelve and a half years after recovery.

The Adhesion Test

In many ways the results of this test are easier to read than the agglutination reaction, and it can be performed more rapidly. Its history is briefly as follows. A number of observers (Laveran and Mesnil, 1901; Rieckenberg, 1917; Brussin, 1925; and Kritschewski and Tscherikower, 1926) had drawn attention to the fact that particles such as bacteria or blood platelets would become adherent to trypanosomes and spirochaetes in the presence of immune sera. Brown and Davis (1927) applied this as a means of differentiating serological types of leptospira, and proposed the name of the adhesion test.

The questions now arise whether this test is specific and sensitive, and whether the reaction persists for some time after the disappearance of the symptoms of the disease. As regards the specificity of the test, we had no difficulty whatever in differentiating *L. biflexa* from *L. icterohaemorrhagiae* on the one hand, and *L. hebdomadis* on the other. We were also able to prove by means of this reaction that five strains of *L. icteroides* which Noguchi (1925) maintained were the cause of yellow fever and were morphologically and serologically distinct from *L. icterohaemorrhagiae*, were indistinguishable from the