THE NATURE OF ACIDOSIS

BY

O. C. M. DAVIS, D.Sc.LOND., M.R.C.P.
READER IN MEDICAL CHEMISTRY IN UNIVERSITY OF BRISTOL, AND
PHYSICIAN TO ROYAL HOSPITAL FOR SICK CHILDREN
AND WOMEN

AND

F. W. RIXON, M.Sc.VICT., Ph.D.GIESSEN
LECTURER IN CHEMISTRY IN UNIVERSITY OF BRISTOL

For a considerable number of years the word "acidosis" has been used in connexion with certain symptoms exhibited by children, and the question may well be asked whether this word connotes a disease, a symptom, or a chemical phenomenon.

In certain textbooks and writings the term is considered to be an undesirable one. We do not share this somewhat dogmatic view, but think it behoves practitioners to inquire what, precisely, is meant by the expression referred to. Since during the last fifteen years we have carried out a number of researches involving the nature of the condition sometimes labelled "acidosis," it may be helpful to discuss some of the theoretical and practical points bearing on this subject.

Definition

At the outset it is desirable to recognize that the meaning of many words commonly used in the English language has been arrived at by convention, and not solely from etymological considerations. If we assign a meaning to the word "acidosis" which is based entirely on such considerations the expression simply denotes "a state of acidity," and such a definition leaves much to be desired.

Unfortunately, there appears to be some confusion as to the meaning of this word and other related words commonly met with in medical literature. Thus in a recent textbook¹ the authors apparently consider "acidosis" and "acidaemia" to be synonymous terms (see index). On page 588 they make the statement that: "In acidaemia there is a tendency of the blood to become acid." On page 589 they refer to a compensated acidaemia. In view of the fact that there is always a normal tendency for blood to become acid, this seems to be an unsatisfactory definition. We prefer the definition given in another textbook²: "Acidaemia is an increased hydrogen-ion concentration of blood beyond the normal limits."

Wells³ writes:

"Acidosis, therefore, is a condition in which the essential feature is not an actual acidity of the blood, but the impoverishment of the blood in available bases, whereby there results a decreased capacity of the tissues to get rid of carbon dioxide and other acids formed in their metabolism."

He further mentions that the acid so formed is neutralized by ammonia, which would normally go to the formation of urea, and that the result is an increased concentration of ammonium salts in the urine.

In 1925⁴ we pointed out that the term "acidosis" had been employed to denote a variation in the acid-base equilibrium in the blood, causing a diminution of the alkali reserve. We further suggested that for clinical purposes this term might be taken to denote a condition in which the tendency for acid products of metabolism to accumulate in the blood and tissues exceeds the normal.

The term "acidosis" is not so inapposite as some writers consider it to be, since in cases where excessive production of acid occurs in the body there must, at one stage, be an actual but evanescent state of acidity previous to its neutralization by compensatory mechanisms.

Scope of the Term

While it is admitted that certain diseases—for example, chorea—are named after one prominent sign or symptom, most medical practitioners will probably agree that for clinical purposes it is unwise to diagnose any special disease on the evidence of one sign or symptom, and a special syndrome is usually sought for by the physician.

If this be admitted, acidosis can hardly be referred to as a special disease, since it is a biochemical phenomenon which may occur as one of the signs which, associated with other signs or symptoms, may constitute various syndromes, quite characteristic of definite but different diseases. It may, for example, be a prominent sign in the condition known as cyclic or recurrent vomiting, in the related toxæmic vomiting of children, in diabetes mellitus, tuberculous meningitis, and in a number of other disease conditions.

It is unfortunate that the condition known as acidosis is frequently confused with a related condition known as "ketosis," in which the urine contains acetone bodies. The two conditions may exist apart from one another, or they may be associated. Whereas ketosis may be readily demonstrated by a qualitative examination of the patient's urine, it is not possible to detect acidosis without the aid of quantitative analysis. It is surprising how often a qualitative examination of the urine of children will reveal the presence of traces of acetone bodies; this fact has probably given rise to the misdiagnosis of acidosis in some of these cases. When we remember that the very subtle equilibrium which we call health depends to a great extent on the relationships between various chemical and physical processes taking place within the body, it is not surprising that many causes, such as bacterial invasions, psychical trauma, and dietetic errors should upset this equilibrium.

There appears to have been some confusion as to the value of blood examination as a method of clinical diagnosis. Thus Wells,³ referring to blood, states:

"Very rarely is there more than the slightest deviation from the normal proportion of free H and OH ions, which is slightly on the alkaline side of exact neutrality. This neutrality is one of the most fixed of all the constants of the body."

Nevertheless, in dealing with ways of estimating acidosis, he gives as one method "direct estimation of the hydrogen-ion concentration of blood." He also considers the examination of urine to be unsatisfactory. We do not agree with these views.

Determination of R Factor

Since it is essential that the composition of blood should be maintained within narrow limits of variation, it is obvious that tendencies to upset this relative constancy can be best detected by an examination of variations in respiratory exchange, or, more readily, by a quantitative investigation of the urine.

In July, 1923, a paper was read by one of us⁵ on a possible mathematical relationship between urinary acidity and ammonia-combined acid in urine, as determined by Folin's method. We introduced a very simple factor "R," which denoted the ratio between the volume of alkaline solution required to neutralize a definite volume of urine and the volume of the same alkaline solution required to neutralize this previously neutralized urine after the addition of neutral solution of formaldehyde. Thus,

$$R = \frac{\text{titration II}}{\text{titration I}} \text{ or } \frac{\text{ammonia-combined acid}}{\text{free acid}}$$

Further papers were published by us in 1925⁴ and 1927,⁶ and in these we pointed out that the normal value of R, when determined with early morning specimens of

urine, appears to be round about 1, and not exceeding 1.4 in normal health, but considerably increased in various abnormal states of metabolism. At an early stage in our investigation we came across an interesting example of the phenomenon we define as acidosis.

In one healthy adult male the average value for R, determined with early morning specimens, was 1.14 (maximum, 1.35; minimum, 0.88). After he had taken 10 grains of aspirin on the previous night the following values for R were obtained: 8 a.m., 1.96; 11 a.m., 1.47; 3 p.m., 1.05.

It should be noted that the acidity titratable with standard alkali was diminished rather than increased, but the ammonia-combined acid greatly increased. As described in the papers referred to, we were able to determine the ratio R—later referred to in our work as R_A —in health and disease.

Its Practical Application

A paper published in 1928⁷ contained further notes on the subject. We examined a number of urines where acidosis existed in association with ketosis, and also urines in which one of the conditions was present without the other.

One case was that of a little girl, who was the subject of cyclic vomiting, the attacks of vomiting being initiated by various causes—for example, an attack followed an accidental blow with a spade on one occasion; on another occasion an attack supervened after a prolonged railway journey, and in other instances dietetic errors seemed to produce the vomiting and other symptoms.

In health the average value of R for early morning specimens of urine was 1.07, and during one attack on successive days this ratio was 1.8, 2.95, and 3.45, acetone bodies being also present. During another attack acetone bodies were present with a normal ratio—absence of acidosis—whereas at the end of the attack there was marked acidosis ($R = 3.25$), but no ketosis.

In a case of diabetes which terminated fatally R_A varied from 1.82 to 3.6, acetone bodies and sugar being also present. In one case of tuberculous meningitis acetone bodies were present, R_A being 3.

Summary

We submit:

1. That the term "acidosis" is a very convenient one, and should be used to denote a biochemical phenomenon, in which the normal tendency of the body to form acid products of metabolism is enhanced; and that the phenomenon is exactly comparable with such phenomena as elevated temperature, accelerated pulse, and various rashes which may form integral parts of quite distinct symptom-complexes. This definition postulates that there can be no "types" of acidosis.
2. That in acidosis the accumulation of acids in the blood is prevented by compensatory mechanisms. When these mechanisms fail acidæmia supervenes.
3. That the detection of acidosis by quantitative analysis of the early morning urine may furnish a valuable sign, and help in the construction of a definite disease syndrome.

REFERENCES

- ¹ Halliburton and McDowall: *Handbook of Physiology*, 1933.
- ² Samson Wright: *Applied Physiology*, 1926.
- ³ Wells: *Chemical Pathology*, Fourth Edition, p. 557.
- ⁴ *Clinical Journal*, February 11th and 18th, 1925.
- ⁵ *British Medical Journal*, 1923, ii, 21.
- ⁶ *Ibid.*, 1927, i, 758.
- ⁷ *Archives of Disease in Childhood*, February, 1928.

The Ministry of Health has issued in pamphlet form (H.M. Stationery Office; price 6d.) a revised list of sanatoria and other residential institutions approved by the Minister for the treatment of persons suffering from tuberculosis and resident in England and Wales, with the names of the administrative counties and county boroughs in which the institutions are situate.

MONILETHRIX

A SECOND GROUP OF FAMILIAL CASES

BY

J. GOODWIN TOMKINSON, M.D.

HONORARY CONSULTING PHYSICIAN, SKIN DEPARTMENT, WESTERN INFIRMARY, GLASGOW; DERMATOLOGIST, EDUCATION HEALTH SERVICE, HEALTH DEPARTMENT, GLASGOW; AND EDUCATION AUTHORITY, COUNTY OF DUMBARTON

In the Section of Dermatology of the Centenary Meeting of the British Medical Association, held in London in July, 1932, I read a paper on this condition based on twenty-two familial cases occurring in five generations in a mining district about eight miles east of Glasgow.*

On August 24th, 1933, I saw, in the Western Infirmary, Glasgow, a girl aged 8 years sent by Dr. W. H. Howat of Dalmellington, Ayrshire. She had just come under his care, and the condition for which she sought relief had previously been diagnosed as ringworm of the scalp.

First Case

On examination there was no mistaking the affection of the girl's scalp. The short hairs, some twisted, and exhibiting nodes, so distinct that careful inspection revealed them to the unaided eye, associated with the usual accompaniment of lichen pilaris (especially evident suboccipitally), presented an unmistakable picture of monilethrix. According to her mother lichen pilaris had been present from birth, and the hair had been abnormally short in infancy.

Her Family

I examined three other affected members of the family on February 3rd, 1934—namely, the mother herself and two boys, one aged 13 years, the other aged 3 years. In all, four persons are affected.

As regards the older boy, his mother said that the hair of the scalp up to the age of 9 or 10 years was very short at the back, but then began to grow somewhat better. This resembles, to some extent, the history of one of the group of twenty-two cases reported at the Centenary Meeting—namely, a youth whose hair at the age of 14 years showed signs of improved growth which, being coincident with puberty, might have been due to increased activity of the sebaceous glands. It is not improbable that the mother's statement as to age of improvement in hair growth of the boy now reported was not absolutely accurate, and that the improvement occurred when he was somewhat older. In his case she thought that the lichen pilaris began at about the age of 3 years. It was quite pronounced on the nape and occiput at the time of inspection.

The younger boy showed signs of the affection when a year old, and he has (November, 1934) well-marked lichen pilaris of the nape.

The mother, aged 32, stated that in her own case there was no improvement in growth with the advent of puberty, and that at one time "pimples" (? lichen pilaris) were present on the nape. These have now practically disappeared.

Microscopical examination revealed characteristic hairs in each subject. They all had pediculosis capitis.

Family History

There are three other female children of the family who are unaffected. One of them has flaxen hair, the others reddish-brown hair like the colour of that of those affected. In order of age the affected children come first, fourth, and sixth. The father, a bricklayer, is unaffected. His grandfather was employed as a forester on an estate at Stranraer, as also was his father, who was later employed in the same capacity on a Cumnock estate. The mother's father and grandfather were iron-workers at Dalmellington Iron Works. Her mother was the daughter of a weaver. She has a brother, a miner, who lives near to her, who is unaffected, and so are all his children. Her sister and family show no signs of

* *British Medical Journal*, 1932, ii, 1009.