15 to 25 years) recently seen at Victoria Park, and nearly all supplied some hint, not of what physical signs, but of what sputum test or skiagram would reveal.

Rejecting six cases which were apparently of at least six months' duration when I first saw the patients, I note that thirty-seven out of the remaining forty-four had symptoms suggestive of tuberculosis. Thus fourteen of these young persons complained of cough with expectoration, while no fewer than twenty-three admitted to haemoptysis ranging from stained sputum to frank haemorrhage of the classical type. Approximately 50 per cent. of the total number gave a family history of tuberculosis or admitted close contact with infection. A closer questioning would doubtless have revealed an even higher figure. Physical examination yielded suggestive signs in only half of the forty-four cases, while in the remainder physical signs were absent or equivocal on what one might call a surgery examination.

I believe that the somewhat discredited old term "the tuberculous diathesis" conveyed a measure of truth. The patient who looks the part is more likely to be tuberculous than the patient who does not. And if attacked the former is more likely to succumb than the latter. As middle life approaches, and in the male sex, the patient's appearance is less distinctive.

The plaintive whine of the patient seeking sympathy is seldom—I had almost said never—heard from the tuberculous. The tendency to minimize symptoms, return evasive answers, or offer explanations for any symptoms reluctantly admitted is so typical of the consumptive that such a disposition should at once suggest the possibility of this disease. This is especially noteworthy when the question of blood-spitting is raised. If the tuberculous patient has had no haemoptysis he denies in the tone of one who has been insulted by an offensive suggestion. In haemoptysis from other causes the patient may treasure up his expectoration, proudly unroll his handkerchief for your inspection, or triumphantly extract a little bottle of blood-stained sputum from his trouser pocket. He does not explain that he knows it only came from the back of his throat. "I had my tell-tale handkerchief, and would not even look at it myself because I wanted to believe that I had coughed up nothing but a little discoloured phlegm." Those are not the words of a factory hand, but of a highly trained scientific observer when describing a haemoptysis which occurred on the old underground railway in the days of his youth. The same "ostrich complex" may be reflected in the manner and supporting remarks of a parent.

The patient's appearance is worth more than a passing glance. The bright eyes and delicate complexion of the consumptive girl are familiar to us all. The smooth skin of the tuberculous was noted by Hippocrates. Even in middle age the skin of the thorax is often curiously smooth and fine in texture. Sweating axillae are more significant than a history of night sweats. The latter is too often obtained from all classes of patients to have much diagnostic significance.

Assessment of Symptoms and Signs

With regard to the physical examination, inspection of the form and movements of the chest are of little help in the detection of incipient tuberculosis. The same may be said of percussion, unless pleurisy with effusion or bronchopneumonic consolidation complicates the picture. But auscultation may be of more value, for a weakening of the breath sounds may be obvious over an area of early infiltration. Fine, moist crepitations—heard best in the inspiratory phase following a short cough—audible over the apex or subapex of an upper or lower lobe are almost diagnostic. But we must not expect to hear

them. They indicate very active disease and widespread involvement of the affected lobe. The apex of the lower lobe is favourably situated for auscultation. We are not likely to forget the "zone d'alarme" internal to the spine of the scapula when searching for crepitations, but we are apt to forget that a second "zone d'alarme" is situated internal to the angle of the same bone.

If the patient's symptoms, history, manner, appearance, if the physical signs or their significant absence, indicate the possibility of tuberculosis, the sputum test, repeated two or three times if necessary, will in the majority of cases prove the diagnosis. If the sputum test is negative, x-ray examination is essential before a final opinion can be given. And if this supplies evidence of tuberculosis x-ray control will thereafter be essential, for the progress of the case can only be adequately studied and its treatment properly conducted in the light of the information that only x rays provide.

Conclusion

Finally, I would stress the importance of radiological technique and interpretation. A poor skiagram is often worse than useless, and a good skiagram wrongly interpreted may lead to disastrous results. As an illustration of the difficulties that may attend the correct interpretation of a chest radiogram I might instance a case recently under my care. The patient had worked with silica for thirty years. The skiagram showed a small tuberculous cavity and a general miliary stippling of both lungs such as is commonly seen in pneumoconiosis. I jumped to the conclusion that the case was one of silicosis, but the skiagram might have been interpreted as indicating miliary tuberculosis. A few weeks later necropsy revealed miliary carcinomatosis and a single insignificant tuberculous cavity.

THE SPRINGS OF NEUROSIS*

BY

G. F. WALKER, M.D., M.R.C.P.

SUNDERLAND AND DARLINGTON

Those whom we loosely call "neurotic" live in a dismal twilight of ill-health, and drift from doctor to doctor or from cult to cult without receiving the study and understanding which we accord to other types of patients. Neurosis probably accounts for far more suffering and disability than tuberculosis, yet a comparison between the amount of study and research extended to each of these diseases reveals a serious deficiency in medical education. Asthmatics, diabetics, and dyspeptics receive careful handling, co-ordinated research, sympathetic reception, and rational treatment. Neurotics get short shrift. Indeed, it is not too much to say that the average doctor bristles with prejudice against those whom he calls neurotic, and against the word "neurosis" and everything about it. This prejudice is born of fear, which in turn is born of ignorance. For among the many anomalies of the existing system of medical education the absence of proper instruction in psychological medicine is perhaps the most glaring and chronic. Moreover, with one or two exceptions, the literature on psychological medicine is very dull and unhelpful. The psychologists were called "sad dogs" by Sir Clifford Allbutt when he discussed the confusing terminology of psychological medicine.

The object of this paper is to attempt to survey the whole range of those loosely termed "neurotic" in terms, not of formal psychology, but of practical clinical medicine.

I begin with a rough classification.

* A lecture given to the Darlington Division of the British Medical Association, November 22nd, 1934.

The Neurotic Hierarchy

- A. THE FALSE NEUROTICS are those in whom organic disease is present, but is either
Unrecognized (from not knowing), or
Unperceived (from not looking).
- B. THE ASTHENICS suffer from fatigue due to many causes, including:
Chronic pain or discomfort.
Chronic postural defect.
Endocrine disorder.
Toxaemia.
- C. THE ODD AND ECCENTRIC.
- D. THE NEURASTHENICS.
Sexual.
Environmental.
Traumatic.
- E. THE HYPOCHONDRIACS.
Senile.
Menopausal.
Vicarious.
- F. THE TRUE NEUROTICS, subdivided into several types, concerning:
Flight, or the seeking of refuge.
Aggression, or the seeking of power.
Imitation, or submission to suggestion.
Conflict between self and herd.
Repression of degraded instincts.
- G. THE HYSTERICS.
- H. THE PSYCHASTHENICS.
Obsessional.
Compulsive.
- I. THE PSYCHOTICS, including:
Manic-depressive types.
Schizophrenic types.
Paranoid types.
- J. DEMENTS IN THE INAUGURAL STAGES.
Late syphilis.
Arteriosclerosis.

My plan is to consider each of these classes briefly in turn, with the exception of I and J. The members of these two latter classes frequently masquerade as neurotic in the early stages, but that is as far as they concern us here. I merely want to emphasize that many patients loosely labelled "neurasthenic" are in reality suffering from manic-depressive insanity.

A.—False Neurotics

The false neurotics are those in whom we fail to detect organic disease, and I think this failure more often occurs from not looking than from not knowing.

The commonest maladies which are prone to mimic neurosis are suppuration in the accessory sinuses, dry bronchiectasis, disseminated sclerosis, and nodular fibrositis. The latter is a really common disease, and is the source of much ill-health and suffering. It can, further, be the source of great credit to him who knows of it, recognizes it, and cures it.

In similar fashion I think there is no doubt that phthisis, cerebral tumour, post-encephalitic states, and epilepsy may all mimic neurosis, but there are two organic diseases which I particularly wish to stress as very real, very distinct, very disabling, and very chronic. Each has become a rather dilapidated diagnosis, but each mimics neurosis very closely and very frequently. I refer first to gastritis and secondly to anaemia, and I go so far as to assert that no person should be dubbed neurotic, no matter what the temptation, until the physician is confident that gastric secretion and mobility are normal and the haemoglobin content well up to the normal range.

B.—Asthenics

The asthenics form a bulky and unwieldy class, but individually each member is worthy of very close scrutiny. There are innumerable causes of moderate pain, perhaps nothing more than occasional discomfort, which nevertheless in time wears down a person's stamina and fortitude. Among these causes might be mentioned at random cholecystitis, spinal arthritis, salpingitis, and pyelitis.

I do not know exactly how the assumption of personal untidiness affects the temperament, but I do know empirically that the surest way for a man to induce in himself a dingy neurosis is for him to allow physical slackness in himself. One might indeed say "Untidy man—untidy mind."

Endocrine dysfunction is one of the commonest causes of vague invalidism likely to be deemed neurotic. The most important type is called "chronic benign hypothyroidism." This condition is far short of classical myxoedema, but can often be recognized in tired middle-aged men. The patient complains of fatigue, flabbiness, somatic pains, headache, and pessimism. It is one of the few maladies in which an educated man will actually use the word pessimism in describing his symptoms. There is probably no condition where accurate diagnosis and appropriate therapy is more satisfactory to all concerned.

Other types of endocrine disorder in which apparently nervous invalidism may ensue include hypofunction of the adrenal cortex and hypofunction of the anterior pituitary. Extreme degrees of these are known respectively as Addison's disease and Simmond's disease, but milder syndromes corresponding to chronic benign hypothyroidism undoubtedly exist. Clinical research is here very backward. In connexion with endocrine disorders I have not forgotten Graves's disease, in which the emotional element is always present and frequently predominant.

Toxaemic asthenia is again a very vague and unwieldy group. I will hurry over it by stressing merely the general constitutional depression which may arise from pyorrhoea, from post-influenzal states, from cholecystitis, and from excess of nicotine and caffeine.

C.—The Odd and Eccentric

A consideration of those who are odd and eccentric involves the term "conditioned reflex." A "conditioned reflex" may be defined roughly as a habit which has been instilled by experience. According to one school of psychology associated with the name of Watson all temperament and all behaviour is made up of conditioned reflexes. In other words, a man is no more and no less than what he has been taught to be. The advanced students of Watson's notions of behaviour believe that there is no essential difference between conscious and unconscious, no volition, no moral law, no free will, and no mutual tenderness. A grown man, in other words, is merely a parcel of reflexes. I am not, of course, going to press this view, but I mention behaviourism because of two important aspects.

1. First, an enormous amount of neurotic invalidism springs from faulty education in home, church, and school. Some children are literally taught the technique of irrational fear or coached in the pernicious habits of evasion and compromise. "Safety first" is a thoroughly bad psychological principle. And a great many notions are given in the name of theological instruction which may or may not be sound theologically but which are certainly psychologically calamitous.

2. The second point about conditioned reflexes is that their interruption or eradication may lead to perplexity, bewilderment, anxiety, and jarring. In clinical terms, the break-up of lifelong habits may bring about a condition of neurosis far more severe than is ordinarily expected. Happy, therefore, is the man who prepares for his own retirement not only financially but also intellectually. There is no neurotic like an elderly neurotic.

D.—Neurasthenia

Neurasthenia is a well-defined and sharply limited malady in which the central feature is fatigue. Along with fatigue go obviously irresolution, indecision, and irascibility.

Sexual neurasthenia may spring up in one of several ways. One of the commonest types of sexual neurasthenia is that which arises from faulty technique in birth control. In this the autonomic nervous system of one partner is subjected to regular and repeated jarring by means of stimulation without adequate gratification. Fear and repugnance ensue instead of repose. There is

probably more misery and ill-health from this than from any other single cause in the whole range of medicine. Another and milder type of sexual neurasthenia is seen in young people who inflict on each other a prolonged and stagnant engagement period. Yet other types of sexual neurasthenia spring from fears, taboos, and repressions which I think are self-explanatory. And, finally, a very large amount of sexual neurasthenia springs from faulty education and confusion between the poetical and the ridiculous.

Environmental neurasthenia is fatigue caused by occupation which is too monotonous to be interesting and not monotonous enough to be capable of automatic performance. Weavers, machine-tenders, packers, and mass-production factory hands engage in work which is so automatic that they can indulge in relief by gossip or day-dreaming. By such relief fatigue is avoided. On the other hand, work which is monotonous and yet sufficiently catchy and varied to demand ceaseless attention often causes fatigue and neurasthenia. Accordingly neurasthenia is exceedingly common among bus-drivers, telegraphists, schoolmistresses, and housewives.

Traumatic neurasthenia is a very vague section of medicine. There is no doubt that a bodily shaking, coupled with the springing up of ideas of grievance, may undermine the most stalwart temperament. Exactly how this happens is not at all clear. In some cases there is a physical basis in the form of an actual bruise of the brain. I have personally seen such a bruise twice in cases of atypical traumatic neurasthenics who happened to die from other causes. I think a brain bruise may be a common lesion. Patients with traumatic neurasthenia drift from door to door, but rarely come to post-mortem.

Of course, traumatic neurasthenia is nearly always worsened by brooding, physical deterioration, misconceptions, and cupidity. The injured workman with an unsettled claim for compensation, who engages in endless pilgrimages to doctors' consulting rooms and solicitors' offices, and labours under the diagnosis of traumatic neurasthenia is a standing reproach and slur upon the professions of both law and medicine.

E.—Hypochondriasis

With the hypochondriacs or valetudinarians we enter the group of true neurotics. A true neurotic is one whose actions and temperament are coloured and flavoured by distorted instincts and emotions.

The hypochondriacs suffer from distortion of the instinct of self-preservation and the instinct of collecting and accumulating things. They are, as Hutchison has described, unreasonably and rather foolishly anxious about their personal health. They collect symptoms and doctors just like other people collect pictures, china, furniture, or coin of the realm. Fortunately most of a hypochondriac's collection consists of fakes and spurious symptoms, but inevitably he turns up sooner or later with a genuine article. No hypochondriac has a collection of symptoms that is wholly and permanently spurious.

Clinically hypochondriasis is very common indeed, and extreme degrees are frequently encountered. The late Sir Clifford Allbutt used to tell the story of an elderly hypochondriac whose study was decorated with numerous framed originals of specialists' reports, and Sir William Hale-White recently reported in the *Lancet* the case of the hypochondriac whose consumption of drugs was so great that the chemist called each day for orders.

Before leaving the hypochondriacs I should like to recall Hutchison's further observation—that hypochondriasis may be vicarious. In other words, a son or daughter may exhibit pathological solicitude about an elderly parent.

F.—True Neurotics

The true neurotics exhibit vague invalidism, intractable symptoms, and irresolute temperament because they suffer from distortion of primary instincts and emotions. It is alarming to discover how little one knows of the mechanism of this distortion when one comes to reflect seriously upon it. Factors of heredity, environment, and education all play a part, but there must be many other factors which are not properly charted.

The instincts which on distortion are most likely to be springs of neurosis are the instincts of self-preservation, the herd instinct, the love of power, the love of soft and warm security, and the instincts centred round sex. These cannot be considered separately because they act reciprocally and in highly complex fashion. Instead I suggest a consideration in terms of practical clinical medicine somewhat as follows.

1. Neurotic invalidism, especially of the submissive, resigned, placid type, may be the result of subconscious mechanism in a person who finds the world too hard a place and who desires security, refuge, and irresponsibility without loss of self-respect and without the stigma of incompetence or cowardice. A great deal of gastric, cardiac, and colonic illness springs from this source. Cardiac and gastric neurotics often enjoy quiet, comfortable, and respectable invalidism, which absolves them from personal responsibility and strife with the world.

2. Neurotic invalidism, especially of the flamboyant or ostentatious type may be the result of subconscious mechanism in a person who ardently desires power and priority. This phenomenon is frequently seen in children. Illness is a quick and certain means to tyranny and precedence in many a household. In certain types of persons the subconscious mind readily provides such means. Pseudo-angina, mucous colitis, and pyloric spasm are the best examples of this kind of flamboyant invalidism. There is a character in one of Shaw's plays who illustrates this type of invalidism. He was a repressed dissenting clergyman, who always referred to the Church of England as the "Scarlet Woman." He sought, and easily secured, immense power and prestige in his household and district by neurotic cardiac invalidism. For years he was a domestic tyrant of the worst type. By a grim irony he died of scarlet fever, and at the post-mortem the heart was found to be perfectly normal.

3. Neurotic invalidism, especially of the placid and peaceful types, may be nothing more than distortion of the instinct of submission, so that crude imitation takes place. A potentially neurotic person may copy the symptoms of one who is loved or feared. Aerophagy, habit-spasm, constipation, and sundry types of dyspepsia often arise in this way. The same subconscious mechanism operates by way of suggestion. A good deal of neurotic invalidism is inculcated in the timid, ill-informed, and debilitated by querulous relatives, newspaper advertisements, and injudicious "health" articles in weekly journals. Occasionally the same happens in our own consulting rooms, by what may be our most trivial and innocent remarks or our most natural diffidence.

4. Neurotic invalidism may also arise from conflict, which is the technical term roughly describable as subconscious perplexity. Such perplexity occurs when instincts of self-preservation clash with instincts concerned in the promotion of the welfare of the herd. A solution of the difficulty is furnished by subconscious mechanism in the form of safe, comfortable, and respectable illness, with preservation of self and escape from the sacrifices required in herd welfare or social service. Another form of conflict or subconscious perplexity arises when sexual instincts undergo regression or degradation. In such cases there is a serious clash between social and moral forces outside the individual and erupting forces within himself. Such a conflict is highly provocative of nervous restlessness, alternating depression and elation, superstition, selfishness, vanity, and all the hall-marks of neurotic invalidism.

G.—Hysteria

The hysterics are refugees from reality. Contrary to popular impression, the word hysteria does not imply turbulence, talk, and tantrums, but rather a kind of peaceful, pleasant pessimism.

The hysteric is one who, by subconscious mechanism, converts his emotional difficulties into visible outward physical expression. The hysteric's difficulties may spring from his own ideas of inferiority or superfluity, from a knowledge of incompetence or insufficiency, or from intolerable environmental difficulties. The commonest hysterical manifestations to-day are aphonia, sundry tics, tremors, spasms, epileptiform convulsions, and sundry

forms of muscular weakness. Each illness provides a solution of intolerable difficulty or a refuge from intolerable hardship. Unluckily for the hysteric, the subconscious solution or refuge is often highly troublesome and inconvenient. Hence the description of hysteria which I have just given—namely, pleasantly peaceful pessimism.

H.—Compulsive and Obsessional Neuroses

These are particularly severe and painful illnesses in which the patient labours under an imperious and overriding requirement to perform unnecessary and irrational acts, or to perform needless ritual or to think along ceaselessly recurring lines of thought. One of the

worst forms of compulsion neurosis is that in which the patient finds herself ceaselessly washing her hands. Milder forms of obsessional neurosis include sundry forms of explosive utterance or coprolalia, sundry odd habits, and mild forms of kleptomania. These forms of neurosis spring from hidden fears. The hand-washing neurosis, for instance, may spring from fear of contamination, real or metaphorical. Most of the fears concerned can be traced to fears about personal integrity.

The remaining classes (I and J) in the table are not strictly within the scope of this review. But members of these classes form a very large number of those commonly called neurotic or neurasthenic.

PULMONECTOMY

A PRELIMINARY REPORT ON TWO CASES IN WHICH AN ENTIRE LUNG HAS BEEN SUCCESSFULLY REMOVED

BY

GEORGE A. MASON, F.R.C.S.

SENIOR SURGICAL REGISTRAR, ROYAL VICTORIA INFIRMARY, NEWCASTLE-UPON-TYNE; THORACIC SURGEON, WOOLEY SANATORIUM, NORTHUMBERLAND

Total pulmonectomy, although occasionally attempted, was in no instance successful until 1931, when Nissen, working in Sauerbruch's clinic in Berlin, resected the entire left lung of a girl aged 12 years who was suffering from bronchiectasis secondary to a traumatic rupture of the left main bronchus.¹ In the following year Haight, working in Alexander's clinic in Michigan, repeated this success when he removed the entire left lung of a girl aged 13 years, in whom bronchiectasis had followed delayed removal of an aspirated foreign body.² The operation is probably more hazardous in cases of bronchiectasis than in neoplastic disease owing to the greater risk of sepsis; and although a few additional successful cases have been reported during the last two years in America—notably by Evarts Graham³ and Reinhoff⁴—these have been mostly for growths.

This is a brief preliminary report of two cases of unilateral bronchiectasis—one a girl and the other a boy—which have recovered after removal of an entire lung, the left and the right respectively. There was no degree of stenosis of the stem bronchus in either case, such as was present in those operated upon by Nissen and Haight. Mass ligatures were applied to the lung root in each case

to promote gangrene in the lung prior to its removal. This was the method employed by both Nissen and Haight, and appears to be the only one by which a bronchiectatic lung has been successfully removed.

First Case

N. W., a girl aged 13 years, was referred by Dr. Horsley Drummond. Since the age of 5 she had suffered from recurrent attacks of "pneumonia," each of which had been followed by profuse expectoration of foul sputum. In 1931, following such an attack, an empyema in connexion with the base of the left lung had been drained at another hospital. She was admitted to the Royal Victoria Infirmary, Newcastle-upon-Tyne, at Easter, 1934, suffering from a pneumonic attack. Subsequent lipiodol examination demonstrated bronchiectasis of the entire left lung. The sputum diminished and almost completely disappeared with general treatment, but marked pallor and clubbing of the fingers persisted. The chest wall was retracted on the left side and the sternum unduly prominent.

Operation: May 4th, 1934.—Anaesthesia was administered by Dr. W. J. Phillips. Preliminary basal narcosis with avertin (80 per cent. of the charted "Bayer" dose) was given, intratracheal insufflation of oxygen, warmed ether vapour, and a small quantity of nitrous oxide. A second intratracheal tube was also introduced to facilitate evacuation of pulmonary secretion. The patient was placed on her right side with a sandbag under the chest. A long incision was employed, extending from the spine behind to the sternum in front, and crossing the scapula about 3 cm. above its angle. The latissimus dorsi, etc., were divided in the line of the incision and the scapula retracted forwards and upwards. The fifth and sixth ribs were divided close to the spine, and their intercostal muscles divided throughout the length of the incision. These ribs were then widely separated with the help of a de Quervain retractor. The lung was completely adherent, but the two lobes were freed and mobilized with comparative ease. Two stout braided-silk ligatures

FIG. 1.

FIG. 2.

FIG. 3.