TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some of the major medical disorders met with in general practice

THE TREATMENT OF MEASLES

BY

WILLIAM GUNN, M.A., M.B., M.R.C.P., D.P.H.

The practitioner who has not already done so should occasionally visit the fever hospital to which he sends his measles patients. His aim should be to reproduce in the home, as far as practicable, the conditions which obtain in hospital; if they fall far short of this standard the patient is best removed to hospital forthwith. The sick-room should be large and airy, well ventilated, and warmed by a coal fire. Heating by radiators and hot-water pipes, or by gas and electricity, are equally unsuitable. The former interfere with the free circulation of the room atmosphere, leaving the patient in a warm, saturated air-pocket in which he must perforce remain, while the latter dry the air unduly, producing a state which is highly irritating to the inflamed respiratory tract. A temperature of 60° to 65° F. should be maintained; some variation within these limits is an advantage rather than otherwise. It is well to exclude direct sunlight from the eyes while photophobia is present.

All superfluous furniture, including carpets and curtains, should be removed, even the time-honoured disinfectant-saturated curtain over the door. The bed should be placed as near the centre of the room as possible. The bedclothes will be few and light; a single sheet and blanket should suffice, while the patient wears a single cotton nightshirt opening at the sides. Contrary to common opinion, the measles patient does not readily catch a chill during the pyrexial phase, and it is unnecessary to wrap him up tightly with blankets. Adequate warmth is readily provided by hot-water bottles. If it is found difficult to maintain the requisite room temperature and the patient feels cold, a loose woollen vest may be worn, but a Gamgee or pneumonia jacket is unnecessary.

General Hygiene and Nursing

The patient is washed completely once a day, and, in addition, tepid-sponged at intervals of from once daily to four-hourly. The temperature of the water and the frequency of sponging depend upon the amount of restlessness and insomnia or the degree of pyrexia. The buccal and upper respiratory passages must be cleansed gently and at frequent intervals, as secretions, epithelial debris, and food particles collect in various situations. The buccal cavity is usually best dealt with by inserting the index finger enveloped in cotton-wool moistened with warm sodium bicarbonate solution. In very young and refractory children this procedure is not free from the risk of damaging the delicate and inflamed mucous membranes; any breach of continuity of the mucosa might permit the ingress of pyogenic organisms, and the manipulation should be omitted if there is any possibility of damage being inflicted. The mere act of swallowing quantities of dilute fruit juice, especially lemon juice, at frequent intervals is by itself usually effective in keeping the oral cavity clean and sweet. This may be aided by a simple prescription containing a small amount of potassium chlorate taken internally for a few days. This substance appears to have a specific action in alleviating stomatitis and preventing ulceration of gums, tongue, and lips, and reaches all the affected parts only by swallowing.

An agreeable mixture—30 minims four-hourly, for a child of 1 year, for three or four days only—is:

R	Pot. chlor. ...	...	...	gr. x
	Ac. hydrochlor. dil. ...	...	...	m x
	Syrup. ...	...	...	3 j
	Aq. ...	...	...	ad 3 j

Any attempt at gargling is impossible in the great majority of cases. The glycerin preparations commonly prescribed should never be employed on account of their marked hygroscopic action on the inflamed epithelial tissues; they tend to produce or aggravate the ulcerations and fissures they are meant to prevent. The lips should be protected by the application of vaseline, alone or in equal proportions with ung. hydrarg. ammon. dil. The toilet of the nares and nasopharynx is equally important in the prevention of septic complications, including bronchopneumonia. It is important not to do too much; simple removal of discharges and cleansing of the anterior nares with alkaline solution, followed by application of the protective emollient mentioned above, usually suffice. The nasal douche should never be employed in the acute coryzal stage, and is preferably reserved for cases of persistent rhinitis with purulent discharge or crust formation. The solution should be at body temperature, and applied by gravity by means of the douche can, not more often than twice a day. A suitable prescription is: sodium chloride, borax, sodium bicarbonate—two grains of each to one ounce of water. Older children may profitably be taught to blow their noses, provided they do not do so too violently.

The patient should have few visitors. The ideal arrangement is that he is seen only by those actually tending him. Nobody with acute or chronic catarrhal infection should be allowed in the room, or else a mask should be worn. The utmost care should be taken that all articles of food and drink, and all utensils and clothing, should be prepared and handled in a cleanly manner. All discharges and soiled articles from the patient should be disinfected forthwith by burning, boiling, or immersion in disinfectant (carbolic 1 in 40 or lysol 1 in 20).

Conjunctivitis of some degree is invariably present in measles. The mildest grades require no special treatment, apart from application of ung. acid. boric. to the lids at night. In more severe grades irrigation with boric lotion (4 grains to 1 ounce) with the undine at intervals from twice a day to hourly, according to severity and amount of purulent discharge, with application of ung. acid. boric. to the lids after each irrigation, usually clears up the condition in two to three days. The more popular ung. hydrarg. ox. flav. should not be employed, or only at quarter strength, as it is usually too irritating for the conjunctivitis of measles. In more protracted or subacute infections irrigation with hydrarg. perchlor. 1 in 10,000, or with zinc sulphate 1/2 per cent., may prove more efficacious in promoting complete resolution.

The Diet

The onslaught of the measles virus falls mainly upon the respiratory and the gastro-intestinal tracts, and special attention to these passages is necessary in sickly or debilitated subjects in order to avert dangerous complications. During the winter months respiratory complications are most common, while in the spring and summer months enteritis is to be dreaded most.

Almost the whole gastro-intestinal tract is affected by the causative virus, initiating an inflammatory catarrh, which, if not promptly checked, may quickly lead to severe entero-colitis from invasion by secondary organisms. In the milder grades cows' milk, suitably diluted with water according to age, or with lime-water if there is diarrhoea, given at three- to four-hourly intervals and supplemented between feeds with bland fluids, usually corrects the condition. The usual vague instruction to give, at frequent intervals, small feeds of milk, presumably unaltered, is misleading, and calculated to provoke an intractable enteritis. The feeds should be given at the same intervals as prior to the onset of measles, and be adjusted to the body weight and individual tolerance of the patient. Between feeds the fluid intake is maintained by drinks of water, plain or with 2 to 5 per cent. of glucose, and flavoured with lemon or orange juice. The practice of giving plain water warmed to body temperature is apt to produce nausea and vomiting in many subjects, but this aversion is removed by flavouring the drink with fruit juice. A pinch of common salt is a useful adjuvant, stimulating both appetite and thirst. Many children have a rooted distaste to milk, plain or diluted, but will take it readily when flavoured with tea or coffee or when suitably modified—for example, Horlick's malted milk.

Beef tea and meat extracts are always best avoided in the first week, especially if there has been a tendency to diarrhoea, as they are apt to excite gastro-intestinal irritation. If milk is not well tolerated, as shown by the persistence of diarrhoea or the appearance of curds, it should be citrated or, preferably, peptonized; or it may even be omitted altogether for twenty-four hours, and measures directed towards rectifying the gastro-enteritis be applied forthwith. The great majority of patients, with a suitably adjusted dietary, rapidly regain their wonted appetite, and may be placed on full diet as soon as the temperature is normal. A generous diet is advisable, supplemented by cod- or halibut-liver oil for some months.

Specific Measures

So far we have discovered no specific of proved value in the treatment of measles. The number of so-called "specifics," some of which have enjoyed a reputable vogue, must be legion, but none of them has stood the test of time or of critical investigation. Various antipyretic drugs are still employed in many quarters, especially on the Continent and in America; of these amidopyrine has been the most popular in recent years. In full therapeutic doses (1 grain for each year of age to a maximum of 5 grains, at four-hourly intervals until defervescence), it certainly curtails the duration of the disease, or, at least, of the pyrexial phase; but it does not appear to have any mitigating effect on the severity of the infection or upon the incidence and gravity of complications. Nor should its risks be forgotten. Overdosage is likely to occur if the drug is invariably pushed until the temperature reaches normal, while in certain susceptible subjects a condition resembling or identical with agranulocytosis may be produced.

More success may reasonably be expected to attend the administration of specific immune sera. As the real danger of measles is from the complications which follow in its train, from invasion with pyogenic organisms, we should attempt either to mitigate the severity of the primary virus or to prevent or check the assault of the secondary invaders. The marked success in the attenuation and prevention of measles which followed inoculation with measles-immune sera (convalescent and adult) during the incubation period raised hopes that these measures would prove of value in treatment. In the majority of experiments recorded, 5 to 10 c.cm. of convalescent serum have been given intramuscularly on the fourth or fifth

day of the disease; the clinical results were uniformly disappointing. During a recent epidemic, larger doses of immune serum—20 to 50 c.cm.—have been administered to severe cases by the intravenous route, with demonstrable clinical improvement. The earlier the serum is given the most clear-cut are the therapeutic benefits. The therapeutic scope of measles-immune sera will doubtless be more thoroughly investigated in the near future, when more generous supplies are made available. The recent preparation of highly concentrated placental extracts, rich in measles-immune substances, which have already proved their worth in prophylaxis, give promise of a wider application in treatment.

Usually, patients come under the care of the practitioner at a stage when they are already virus-immune (sixth or seventh day of disease) and not, therefore, amenable to specific antiviral therapy; but secondary invasions are imminent or perhaps well established. Clinical experience has suggested that a potent anti-streptococcal (scarlatinal) serum, in doses of 20 to 50 c.cm. intramuscularly, is effective in lowering the incidence of faucial angina, ulcerative stomatitis, laryngitis, and even bronchopneumonia. Administration of polyvalent anti-pneumococcal serum, in similar circumstances, has proved consistently disappointing, while an effective antiserum for Pfeiffer's bacillus—the third secondary invader in order of frequency—has not been available in this country.

In view of the known fact that ill-nourished, rachitic subjects stand measles badly, early administration of vitamins A and D, in a highly concentrated state, has been advocated in the hope of reducing mortality and complications. On theoretical grounds the method is logical, and benefit actually appears to accrue in those cases in which vitamin deficiency exists, but it has been found difficult to demonstrate any consistently beneficial action in a series of measles patients admitted to hospital as compared with a control series under identical conditions except that the concentrate was withheld.

Complications: Prevention and Treatment

Under conditions of sound hygiene and nursing complications should be rare. They should be anticipated and averted or treated promptly and energetically as they appear. Too often, however, the practitioner is only called when dangerous complications have arisen, necessitating immediate removal to hospital for satisfactory treatment.

THE RESPIRATORY TRACT

Symptomatic treatment of bronchitis may help to ward off suffocative catarrh or pneumonic consolidation. Space forbids full discussion of the treatment of measles bronchopneumonia; it differs little from the treatment of pneumonia in children generally, except in its greater severity and higher mortality. Severe attacks of pneumonia are frequently accompanied by some degree of pleurisy, and should pus form diagnostic aspiration is necessary as a guide to subsequent treatment.

Laryngitis, appearing with the onset of the rash, or shortly thereafter, is a dangerous symptom, demanding prompt and energetic action. Anti-diphtheritic serum in doses of 16,000 to 24,000 units is injected intramuscularly without waiting for the result of bacteriological examination, and the steam jet is employed. The steam tent is not used nowadays, and some authorities have abandoned steam in all forms. It is probably wise to omit it if there is any likelihood of pneumonia ensuing, or if it is already manifest. Examination of the larynx and adjacent structures is made by the direct laryngoscope, and smears and cultures are taken. If the results of inspection and of bacteriological examination negative a diagnosis of diphtheria, anti-streptococcal (scarlatinal) serum is given in

doses of 20 to 30 c.cm. Culture plates nearly always reveal the presence of haemolytic streptococci, but their pathogenic role in a given case is not always easy to assess.

Should operative measures be demanded to relieve urgent obstructive symptoms aspiration of mucus or membrane through special tubes (Willard Parker tubes) is the operation of choice, but subsequent intubation is frequently necessary on account of laryngeal or subglottic inflammation and oedema, or when laryngeal spasm is pronounced. The intubation tube is most easily and safely introduced through the laryngoscope by means of a special introducer, such as the model made for me by the Medical Supply Association Ltd. Experience of the various methods has shown that diphtherial cases respond well to suction and intubation, but, when the haemolytic streptococcus is the causative organism, immediate tracheotomy is usually the most successful operation, on account of the special liability to laryngeal ulceration and consecutive bronchopneumonia. If the complication occurs with, or shortly after, the measles attack, the prognosis is grave, whatever method is employed.

GASTRO-INTESTINAL DISTURBANCES

The symptomatic gastro-enteritis characteristic of measles is usually transient, but may be quickly converted into a severe and rapidly fatal form should the warning diarrhoea be neglected. If the usual measures, such as diluting the milk with lime-water or altering it by peptonization, do not improve matters, although supplemented by hyd. cum cret. (1/2 grain for each year of age up to 3 grains, nocte) or by an astringent such as bismuth and chalk mixture, milk in any form must be discontinued. If there is no vomiting fluids should always be taken by mouth. For a child of 1 year two ounces of half-strength normal saline, with addition of 2 to 5 per cent. glucose, is given at hourly intervals for twelve to twenty-four hours; thereafter full-strength saline is substituted, with added glucose and fruit juice as desired for another twelve to twenty-four hours. If the general and local condition improves whey or secway may be tried cautiously, going on, in a day or two, to skimmed or half-cream Cow & Gate, or like proprietary dried milk. Some children do better on lactic acid feeds immediately after the saline course—either Cow & Gate "lacidac," or, if preferred, two drops of acid. lact. (B.P.) per ounce of skimmed cows' milk with added sugar. Small feeds must be taken at the start—for example, one to two ounces for a child of 1 year—but the feeding interval should be the same as before the attack of measles, and the fluid intake made up between feeds by bland fluids. Albumin water is useful to supplement the glucose, saline, whey, or lactic acid feeds.

Drugs have little place in the treatment of the severer gastro-intestinal disorders; opium is always contra-indicated, and even castor oil has gone out of fashion, but hyd. cum cret. is still used in suitable cases. In collapsed states a good brandy, in doses of 30 minims for a child of 1 year, is frequently invaluable, but should not be employed indiscriminately in gastro-enteritis. If vomiting complicates the picture, fluid must be administered rectally, subcutaneously, intraperitoneally, or intravenously, according to age and degree of intoxication or dehydration. In the worst cases it is best given intravenously by tying a cannula in a vein of the arm or the leg. Feeding by the stomach tube and rectal lavage may be necessary in obstinate cases, but should not be carried out too long.

OTHER COMPLICATIONS

Irrigation with boric acid lotion (5 grains to 1 ounce) suffices to cure the majority of cases of mild purulent conjunctivitis in the course of a few days, but severe or persistent infections require the addition of protargol

10 per cent., or argyrol 5 per cent., once a day, and full dilatation of the pupil, using, preferably, ung. atropin. sulph., 0.25 per cent. For the lids, ung. acid. boric., or quarter-strength ung. hydrarg. ox. flav., should be applied after each irrigation. The appearance of corneal ulceration is a serious complication, but most cases do well, and leave little permanent opacity. Should the ulcer tend to spread or resist the usual measures, the edges should be cauterized with pure carbolic acid or the electrocautery. Even in panophthalmia conservative measures are indicated, as the tendency to useful recovery is striking. Phlyctenular conjunctivitis is more prone to occur during convalescence; boric acid lotions, dilatation of the pupil, and eye-shade for photophobia, and administration of cod-liver oil and ultra-violet light usually lead to rapid cure.

The health standard of the modern child is probably the determining factor in the diminished incidence of severe buccal complications. The milder grades clear up rapidly with potassium chlorate mouth washes and general hygienic measures. For gangrenous stomatitis (cancrum oris) more heroic measures are imperative. The treatment of otitis media, mastoiditis, paranasal sinusitis, suppurative adenitis, cervical cellulitis, boils, paronychia, etc., differs in no essential respect from those conditions arising in other circumstances, except that it is always advisable to defer operative measures as long as possible, to enable the damaged tissues to regain their former vitality and powers of repair.

Convalescence and After-care

Convalescence is generally rapid in the absence of complications, but a generous dietary and abundance of fresh air and active exercise are always advisable. A course of ultra-violet ray treatment, or, better still, a holiday at the seaside or in the country, is helpful in preventing chronic pulmonary catarrh and tuberculosis. Various organizations, such as the Invalid Children's Aid Association, do a valuable service in this respect by providing convalescent and after-care facilities for children whose parents are unable to afford the expense.

Infectivity: Disinfection: Prophylaxis

Measles is infectious from the earliest appearance of catarrhal symptoms or the first rise of temperature, and infectivity is at its height immediately before the onset of the rash. As the rash fades, infectivity is markedly diminished, and is usually completely absent by the twelfth to fourteenth day from the onset of the disease, whether complicated or not. Children are safely allowed up by the tenth day, but, if there is any doubt, a few extra days in bed are advisable in view of the risk of pulmonary complications. Concurrent disinfection of all discharges and soiled articles must be rigorously enforced, as gross contamination may lead to spread of the disease, despite the low viability of the virus. Terminal disinfection may safely be limited to simple cleansing of the room and its contents with soap and water.

Experience has amply proved that we cannot hope to protect the individual or alter the course of measles epidemics by mere isolation of the acutely ill patient. Fortunately, in measles-immune serum we have a safe and reliable agent whereby temporary protection or a modified attack, according to circumstances, may be conferred on individual contacts or on the members of selected communities. The highly specialized technique of measles control is outside the scope of this article. Suffice it to state here that precise adjustment of dosage to body weight and due observance of the conditions governing success have placed in our hands the power, to use if we will, to cheat the measles virus of its prey.