the subject is more or less back in the condition that it was before 1914—another instance of that forward and backward movement that constitutes progress. The war familiarized British surgeons with blood transfusion. This procedure has now become a very useful part of our surgical armamentarium, made safe as it is by a knowledge of the blood groups. Gum saline was put forward as a substitute for blood transfusion and was extensively practised, with results that certainly satisfied those who used it: now for the time being it seems to be passing from popular favour.

Conclusion

If one tries to gauge the changes in surgery since the early years of the present century I think it is true to say that the period is marked more by an improvement in methods already in use than by any outstanding innovation. As to those in whose hands lies the art it can be said that the youthful surgeon is now better equipped to press forward, for he has a better comprehension of the ancillary sciences, in a knowledge of which lies our hope of progress.

PROGRESS IN OBSTETRICS AND GYNAECOLOGY DURING THE PAST TWENTY-FIVE YEARS

BY

JOHN S. FAIRBAIRN, M.A., M.B., B.Ch., F.R.C.P., F.R.C.S.

PRESIDENT, BRITISH COLLEGE OF OBSTETRICIANS AND GYNAECOLOGISTS

This brief retrospect calls for main characteristics rather than specific points, and the first to be mentioned, as the most impressive feature in the obstetrics of the last twenty-five years, is the change in the attitude of mind in which its practice is approached. Its outlook has moved from the pathological towards the physiological field. A like change has occurred throughout medicine, but obstetrics has, in its more limited sphere, been relatively more deeply permeated by the principles of constructive hygiene and preventive medicine.

Ante-natal Care

The spread of these principles throughout midwifery practice has been stimulated by the institution of ante-natal care. Once accepted as an integral part of obstetrics and its aims fully understood, it has tended to leaven the whole. Before its start, however, it was the custom in hospitals with an outdoor and indoor service to examine their patients in order to select those in whom abnormality was present, or suspected, for management in the hospital. The tendency, therefore, was to regard the new system as an extension of the old and concentrate on the detection and correction, when possible, of abnormalities during pregnancy and conditions that might cause complication in labour. The public and voluntary clinics independent of hospitals started as complementary to the older infant clinics, and followed their ideals by placing in the forefront the hygienic, educational, and social aspects. Thus the obstetrical work was more thorough in the one and the constructive hygiene more thorough in the other, but it was long ere the one learnt from the other.

By 1910 little more than a beginning had been made, but thereafter advance was rapid, and the stage reached in 1912-13 can be gathered from a report of an inspection of the London maternity hospitals and departments by Dr. Janet Lane-Claypon,¹ which states that most of them arrange for the medical examination and, if necessary, treatment of women in their first pregnancy or of multiparae with a history of difficult labour; in a few hospitals all women applying for aid were seen by a member of the medical staff. Clearly this was an elementary stage with little constructive hygiene. After the Maternity and Child Welfare Act of 1918 the number of clinics was greatly increased, and gradually it came to be more and more recognized that ante-natal care was primarily a system of constructive physiology for all patients, the discovery and remedying of abnormal conditions being secondary and required in but a small proportion of cases.

¹ Report of Chief Medical Officer to the Local Government Board for 1913-14, p. 31.

Steady progress has continued, but the possibilities of a rational application of physiological methods have not been fully explored. It is, for instance, but recently that it has been fully realized that disordered uterine function rather than mechanical difficulties is by far the commonest cause of delay in labour, and that efforts must be made by ante- and intra-natal management to promote and maintain normal uterine action.

Toxaemias of Pregnancy

The toxaemias of pregnancy still warrant their designation of "a disease of theories," and constructive physiology has its limitations where there is no adequate basis of knowledge of the conditions that favour and those that hinder normal function. Thus there is here no rational basis for constructive physiology, and reliance must be placed on the early discovery of the signs of disordered metabolism and restoration of the normal attempted by lowering metabolism by rest in bed and a low diet, in order to avoid the more serious consequences that may ensue. The failure of such measures entails termination of the pregnancy. The recognition of the liability to permanent renal damage and to recurrence or aggravation in subsequent pregnancies is an important advance. Active intervention in eclampsia has almost entirely given place to treatment on the lines of the Stroganoff or the Dublin school, though there still seems to be a place for Caesarean section in certain acute cases with clearly marked signs of impending eclampsia.

Management of Labour and Prevention of Puerperal Sepsis

The problem of adapting the methods of the operating theatre to the labour room or ward is by no means definitely settled. While there is a fairly general consensus of opinion among obstetricians in favour of an antiseptic and against an aseptic system, there is considerable variance of view between clinicians and laboratory workers regarding the most useful antiseptics for general use. The mercurial salts, which in the view of the former have stood the test of clinical experience, have recently been again denounced by chemists and pathologists as quickly rendered inert by admixture with the body fluids.

The most noteworthy contribution to our knowledge of the causation of puerperal sepsis during the period under review is the part played by spray from mouth and nose in infecting hands, instruments, and dressings, or even the vulval area. Conveyance by carriers of throat infections has been definitely proved, although an accurate assessment of the proportion of infections arising in this way is not yet possible. Some of the cases of infection after

natural expulsion and without internal examination may be thus explained, and face masks are now widely adopted to minimize this danger.

Management During the Puerperium

There is considerable scope for constructive methods to lessen the large amount of disability arising from incomplete involution and muscular laxity by physical treatment in the form of bed exercises to promote the recovery of the abdominal and pelvic floor musculature. The promotion of lactation in difficult cases is one of the triumphs of constructive physiology, and the restoration of the milk flow after weaning for six weeks or more is now possible by measures to stimulate secretion.

Post-natal care is usual in order to promote recovery of overstretched or injured muscles, maintain proper diet and hygiene for lactation, and treat any injuries, chronic infections, or other sequelae of child-bearing with a view to the prevention of later disability.

Stillbirth, Foetal Injuries, and Congenital Disease

An extensive investigation into the causes of dead birth and fatal birth-injuries was conducted under the Medical Research Council shortly after the war, and led to a clearer understanding of their production and nature. Attention has thus been drawn to the prevention of these obstetric disasters. By the intensive treatment of the syphilitic pregnant woman the high proportion of dead births in these cases has been greatly reduced, and if begun before the later months of pregnancy the birth of an unaffected child can reasonably be expected. Similarly, by active treatment of purulent discharge during pregnancy, careful prophylaxis of the infant's eyes during birth, and the notification and prompt treatment of all cases of discharge or inflammation of the eyes in the newborn, the incidence of ophthalmia neonatorum has been greatly reduced. It must be acknowledged, however, that the infantile mortality in the first month of life shows little reduction as compared with the marked diminution in that of the later months of the first year of life. Ante-natal care has not succeeded in lessening prematurity, the largest factor in the high neo-natal mortality.

Gynaecology

Representing the curative and reparative aspect, the gynaecological side of the subject is not affected to the same extent as the obstetrical by the principles of constructive hygiene and preventive medicine, but shows steady progress in technique and therapeutics.

The advances in radiotherapy are outstanding, both in malignant and in non-malignant conditions. Cancer of the cervix has proved particularly amenable to treatment by radium, which is gradually supplanting the radical operation as the results attained are at least equally good and without the risks and disabilities of so extensive an operation. Further, it offers a very considerable degree of relief in inoperable cases. The question of the best procedure in cases with glandular metastases is yet to be decided, and further experience must be awaited before the relative merits of radium applied to the pelvic glands through an abdominal incision and diffuse radiation through the abdominal wall by massive radium or deep x rays can be determined. The advances already made are a hopeful augury of further success.

In the treatment of metrorrhagia from chronic metritis or chronic subinvolution in patients near the age of the climacteric, radium has proved a most valuable and simple means of cure. For menorrhagia due to myomata x rays have also proved effective, and in properly selected cases should be preferred, the essential provisions being that there is no doubt of the nature of the tumour and the patient not of an age at which future child-bearing and

the conservation of the uterus by myomectomy are possible. Cases of tumours above the size of a four and a half months pregnancy are less favourable, and by many would be excluded, owing to the risk of degenerative changes.

Another noteworthy advance is the inflation of the uterine tubes from below to test their permeability in cases of sterility. There is some evidence that the inflation *per se* increases the chances of conception. The injection of lipiodol followed by x -ray examination is another means to the same end, and has the additional advantage of indicating the site and degree of any obstruction present.

Clinical Applications of Advances in Science

Fresh knowledge of the physiology of the ovary, its relation to other endocrine glands and to the menstrual cycle, has resulted in the excretion of anterior pituitary hormones in the urine of pregnant women being used as a test of pregnancy. A positive result is obtained by the stimulation of the ovaries by these hormones after the injection of a few cubic centimetres of the urine into immature mice (Aschheim-Zondek) or rabbits (Friedman). Some fallacies from other causes have been recognized, but the proportion of error is small—2 or 3 per cent.

Since the introduction of the Potter-Bucky diaphragm, a greatly extended use of x rays in obstetrical diagnosis has become possible. Pregnancy can be diagnosed from the fourth month onwards, also whether entopic or ectopic, single or multiple; foetal presentations and positions and certain foetal malformations can be recognized and the maternal pelvic capacity estimated.

Integration of Obstetrics and Gynaecology

The artificial divorce of obstetrics from gynaecology, which twenty-five years ago appeared a possibility on the narrow ground of convenience and differences in technique, faded as the infusion of the ideals of preventive medicine made clear the illogical separation of the physiological from the pathological, and the preventive from the remedial aspect. Provision for post-natal care definitely links together the two divisions of a subject that is scientifically and clinically one. The widespread feeling among those practising specially in this branch that gynaecology should not pass into the hands of those without a wide obstetrical experience was shown by the remarkable unanimity with which those not only in the British Isles but also in the Dominions threw in their lot with the British College of Obstetricians and Gynaecologists at its foundation in 1929. This college aspires to do for its branch of medicine what its older predecessors have done for theirs—to advance study and practice, maintain its standards, and be ready to advise and help in public matters of which it has expert knowledge.

Conclusion

Matters of public interest, such as maternal mortality and maternity services, are too far-reaching for survey here. There are too many factors impossible to evaluate, such, for instance, as the effect of the marked change in the attitude of mind of the whole community towards matters sexual and reproductive, so that the only guide is experience of actual working. A warning of the danger of preconceived ideas was given in the recent New York report, and another from a large English provincial centre, where the percentage of births in hospital has been steadily rising and the mortality rate increasing slightly also. But the plan of a direct attack upon the "black" areas is hopeful and the results from one are an indication that much may be done by an intensive effort over a wide front. Indeed, the need for experience of this kind suggests that plans for a maternity service had better await the results of local experiments.