Conclusion

The sporadic nature of the disease in this country suggests a wide dissemination of a virus of limited virulence, from which it follows that the majority of the adult population have acquired some immunity. This tends to contraindicate the appearance of an extensive epidemic. Small outbreaks in schools or institutions where children are collected are, however, always liable to appear. In handling these the various points I have mentioned should be enforced—namely, isolation of all cases or suspected cases, prophylactic inoculation of serum, and gargles of oxidizing substances for the contacts. In the treatment of cases diagnosed in the early stages serum therapy should be commenced without delay. In view of the crippling nature of the disease, although many cases will evidence little response, an improvement in a few cases is surely sufficient justification for serum therapy.

CORNEAL GRAFTS

WITH THE REPORT OF A CASE

BY

B. W. RYCROFT, M.D., D.O.M.S., F.R.C.S.

HUNTERIAN PROFESSOR AND ASSISTANT SURGEON,
ROYAL EYE HOSPITAL

AND

G. HANDELSMAN, L.M.S.S.A.

SENIOR HOUSE-SURGEON, ROYAL EYE HOSPITAL, LONDON

(With Special Plate)

Where blindness is entirely due to corneal opacity it has been proved that the operation of corneal grafting is of great value, and the recent investigations of Tudor Thomas have revived interest in a branch of surgery which was first successfully practised by von Hippel in 1888. From time to time the technique of such grafting has varied both in the shape of the graft and in the method of fixation.

On the one hand, Elschmig uses a circular trephine, whereas Castroviejo employs a twin-bladed knife which cuts a rectangular graft, and this he retains in position by means of a conjunctival flap. Tudor Thomas, basing his technique on several years of animal experimentation, obtains the graft by a circular trephine of 4 to 6 mm. in diameter, arguing that this is much more likely to "take" than that which has corners or sharp angles. He retains the graft in position by overlay sutures stitched into the cornea on either side of the graft, and lays great stress on the necessity of fashioning a shelving bed for the reception of the graft. In order to allow of subsequent swelling of the graft, this is cut smaller than the bed into which it is to be placed, and in transference the graft is protected by means of sterile olive oil. In the following case an endeavour was made to simplify technique and to combine the trephine method with a complete conjunctival flap, such as is occasionally employed for the protection of large gaping corneal wounds.

Several criteria must be satisfied if the graft is to be successful; they include complete sterility of the conjunctival sac, accurate projection of light, a pupillary aperture neither occluded nor secluded, and transparency of all ocular media other than that of the cornea. The tension of the eye must be within normal limits, and there must be no active inflammation. Cases which fulfil such conditions are: corneal opacification due to chemical burns,

large healed ulcers, and interstitial keratitis which has been quiet for at least twelve months. The scars of old perforating ulcers in which the iris is adherent are not suitable, since, owing to this adhesion, there is a predisposition to vascularity, and consequent opacification of the graft. In our opinion the degree of vision should not be better than hand movements to justify what is still experimental surgery.

The following case is that of a woman who for many years had kerato-iritis of both eyes. Secondary glaucoma supervened in the right eye, which was removed, and complete opacification of the remaining eye had reduced the vision to perception of light. A corneal graft restored the vision to 6/24, and allowed her to read ordinary manuscript.

Case Record

The patient was a married woman, aged 48. She attended the Royal Eye Hospital in July, 1920, and a diagnosis of tuberculous keratitis of the right eye was made. R.V., perception of light. The left eye at this time had fine corneal nebulae: L.V., 6/18. The usual local remedies, together with a course of tuberculin injections, were employed, and her attendance ceased in 1922.

In 1923 the vision was the same in each eye, and in 1924 the right eye was removed on account of secondary glaucoma and an absence of perception of light. In April, 1930, the left eye developed acute kerato-iritis, which subsided after treatment, leaving a vision of 6/60; at this time unit investigation was negative. In October, 1933, the vision was reduced to perception of light, and a blind certificate was issued.

In November, 1934, the patient came under our care, and it was decided to attempt keratoplasty. A complete investigation was carried out, and was negative apart from an increase of the blood sedimentation rate and evidence of old tuberculous cervical adenitis. The eye at this time showed grey infiltration of the whole of the cornea, with localized fasciculi of superficial conjunctival vessels and occasional evidence of deep vascular infiltration. It was not possible to discern the position of the pupil or iris pattern details either by diaphanoscopy or by direct slit-lamp investigation. The cornea was in the early stages of keratoglobus. Projection of light was accurate, and tension was 30 mm. Schiötz.

Preliminary treatment began with four-hourly irrigations of oxycyanide of mercury lotion 1 in 8,000, and local ultra-violet radiation of three minutes' duration by full mercury vapour spectrum of the conjunctival fornices and entire cornea. Subsequent cultures were negative on three occasions. Miosis was obtained by constant instillation of guttae eserine sulph., 1/2 per cent. A donor became available in a case where a young man had sustained a penetrating injury of the upper lid and globe, which left the cornea intact but which disintegrated the posterior half of the eye.

OPERATION

The preparation of the site in the host and the enucleation of the donor's eye were commenced simultaneously. In the host a complete conjunctival flap was prepared circumscribing the limbus, and being separated well back to the equator of the globe. A purse-string suture, No. 1 black silk, was inserted in such a way as to render the aperture eccentric when tightened. Under 4 per cent. cocaine anaesthesia and with seventh nerve novocain block (O'Brien) a 4 mm. circular bed was delineated slightly to the inner side of the centre of the cornea, so as to bring the proposed site of the graft below the margin of the upper lid. When the section had involved half the corneal thickness, the trephine was removed and the bed completed by scissors in such a manner as to produce a shelving edge towards the centre of the aperture. During this period the eye was constantly irrigated by normal saline at body temperature. When the corneal circle was removed the pupil could be seen firmly contracted, the centre of which coincided with the centre of the graft. This had been obtained purely by guesswork: Tudor Thomas has since elaborated a technique which enables the coincidence of the centre of the graft and the centre of the pupil to be accurately

estimated. Under constant saline irrigation a corresponding graft was cut with the same trephine and scissors through the cornea of the enucleated eye, which was held in Tudor Thomas's apparatus during this manoeuvre. The graft was transferred to the bed in a lens spoon, and manoeuvred into position by an iris repositr. At the same time gradual tension was put on the purse-string suture, and eventually this was tied so that the graft was maintained in position and entirely covered by conjunctiva. No part of the suture touched the graft, and no corneal stitches were used. Eserine was instilled, a retention suture fixed the upper lid to the cheek, and ordinary cataract routine was instituted.

Diagram of operation. A-A', Corneal graft. B, Conjunctival flap (note eccentric position of suture). C, Miotic pupillary aperture. D, Iris. E, Lens.

Three days later the graft could be seen in position through the widening conjunctival aperture, and was completely opaque. There was intense conjunctival engorgement, but no pain. At the end of the first week the conjunctival aperture had increased, and the graft was seen to be clear and to have the brightness of normal corneal epithelium. On the tenth day the stitch was removed, and thereafter dressings were continued until the end of a fortnight. Three weeks after the operation the appearance of the grafted eye was as in Fig. 2 (Special Plate). The central black area is the gap through the posterior layers of the graft, surrounded by a shelf, which in this case was slightly too excessive, and which has resulted in some limitation of the field. By slit-lamp examination the iris details can now be seen, and the pupil is found to be fixed by posterior synechiae; there has been an improvement in the opacity of the cornea outside the graft. Unaided vision at this time was 2/60. One month after the operation the unaided vision was 6/60, and the patient was able to walk about the ward, although on account of her limited field she had to exercise great care.

On January 7th, 1935, unaided vision was 6/36, and with the telescopic glasses this is improved to 6/24 and J.6. Judged by the rate at which the graft has cleared there may yet be further improvement. The patient is now able to read ordinary correspondence, and spends most of her time watching traffic through the hospital windows. It is possible that she may be removed from the blind register.

Postscript: March, 1935.—The vision is 6/24 and J.6: the patient now walks abroad unaided.

Commentary

In our opinion the important features of this technique lie in the use of a complete conjunctival flap with constant saline irrigation. It would have been better in this case to have used a larger graft or to have made the shelving edge less, although the stenopaic nature of the aperture undoubtedly improves her vision; in future cases we propose to adopt this technique.

We should like to thank Dr. Rushton of the Royal Eye Hospital for the photographs.

BIBLIOGRAPHY

- Castroviejo: *Amer. Journ. Ophthalmol.*, 1932, xv, 825, 905.
 Thomas: *Trans. Ophthalmol. Soc. U.K.*, 1930, l, 127; *ibid.*, 1931, li, 88, 96; *ibid.*, 1932, lii, 64; *Lancet*, 1931, i, 335; *Proc. Roy. Soc. Med.*, 1933, xxvi, 595 (Sect. Ophthalmol., 33); *Brit. Journ. Ophthalmol.*, 1934, xviii, 192; *Proc. Roy. Soc. Med.*, December, 1934, xxviii, No. 2, 195.
 von Hippel: *Arch. f. Ophthalmol.*, 1888, xxxiv, 108.

COLLOID CYST OF THIRD VENTRICLE OF THE BRAIN

REPORT OF A CASE OPERATED ON, WITH RECOVERY
BY

J. E. PATERSON, M.B., Ch.B., F.R.F.P.S.G.

NEUROLOGICAL SURGEON, VICTORIA INFIRMARY, GLASGOW; CONSULTING
NEUROLOGICAL SURGEON, GLASGOW EYE INFIRMARY, ETC.

AND

MARGARET LESLIE, L.R.C.P.

MUIRHEAD SCHOLAR, CLINICAL LABORATORY, VICTORIA INFIRMARY,
GLASGOW

(With Special Plate)

In the literature will be found a considerable series of reports of a peculiar cystic tumour of the third ventricle of the brain, with notes reflecting the interest which the discovery of this rare type of intracranial new growth seems invariably to have aroused in its finders. Until the appearance in 1933 of Dandy's monograph on *Benign Tumours in the Third Ventricle of the Brain*¹ such references were to findings at necropsy, and the publication of his series of five cases operated on with four recoveries suggested that neuro-surgeons in general had perhaps not been sufficiently alive to the importance of being on the look out for this rare but eminently operable variety of tumour.

In the spring of last year we were fresh from the study of Dandy's book when a case of brain tumour presented itself, unlocalizable by ordinary clinical methods, and having as its most characteristic feature the pronounced intermittency of its symptomatology. In the circumstances it was inevitable that the possibility of this being a case of third ventricle tumour should occur at once, and, by one of those coincidences which do happen sometimes, our suspicions were confirmed by ventriculography; operation followed, and a colloid cyst was successfully removed from the patient's third ventricle. With the exception of Dandy's four cases noted above, a study of the literature has failed to discover any reference to the operative removal, with recovery, of a tumour of this type, and our case would seem to be the first successful operation of the kind to be recorded in this country.

Patient's Previous History

The patient, a moulder aged 34, was admitted to the Victoria Infirmary on April 25th, 1934, having been sent in by Dr. Bell of Falkirk Infirmary with a diagnosis of brain tumour. About two and a half years before admission he began to suffer attacks of pain; these were distributed generally over the cranium, but tended to be most severe in the right side of the head. During the height of an attack there was frequently marked pain in the occipital region. These attacks, the onset of which was not associated with any special time of day or with any very obvious exciting cause, had at first a duration of a few hours. At first, too, a period of months used to elapse between the paroxysms, but as time went on the free intervals gradually shortened, until finally, just before admission, there was only a week or two of freedom, while the attacks themselves were of longer duration and were more severe.

A typical severe attack was characterized by a gradually increasing headache, which reached at length an extreme degree of severity, and on four occasions—the first time early in January, 1934—a period of unconsciousness of a few minutes' duration occurred at the height of the paroxysm. For the past six weeks, too, there had been vomiting on several occasions without any relief to the pain, and the patient's wife was emphatic in her statement that after these severe seizures there was complete failure of memory of the past attack, so that the patient himself had no recollection whatever of its characters. For three weeks before admission it had been noted that giddiness was present during the paroxysms. Even during the early mild seizures there was

R. W. FAIRBROTHER: SOME FACTORS IN THE CONTROL OF POLIOMYELITIS

FIG. 1.—Experimental poliomyelitis. High-power photomicrograph of spinal cord, showing vacuolation, degeneration, and neuronophagia of the nerve cells.

FIG. 2.—Experimental poliomyelitis. Low-power photomicrograph of spinal cord of recovered monkey with residual paralysis. Note absence of large neurons in anterior horn of one side.

B. W. RYCROFT AND G. HANDELSMAN: CORNEAL GRAFTS

FIG. 1.—Shows the eye before operation. The vacant staring look of a blind eye is apparent. The dark patch above the corneal centre is not the pupil, although it may appear to simulate it.

FIG. 2.—Shows the grafted eye twenty-four days after operation. Fixation is now present. The clear dark central gap is seen surrounded by a shelf, which in this case was too excessive. Absence of vascularity and reaction should be noted.