TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some of the major medical disorders met with in general practice

CONGESTIVE FAILURE

BY

SIR MAURICE CASSIDY, M.D., F.R.C.P.

General Principles

In the treatment of congestive failure absolute rest, mental and physical, is of fundamental importance. The patient must be strictly confined to bed and not allowed to make any unnecessary movement. Night and day nurses are essential. The patient should not be allowed to feed himself, and must be lifted on to and off the bed-pan with a minimal amount of disturbance. He should be discouraged from superfluous conversation, unnecessary visitors should be forbidden, and he should be shielded as far as possible from business or other worries. The bed should be neither too low nor too wide. The room should be adequately ventilated, but kept at a temperature of about 65° F. A dyspnoeic patient often demands that the windows should be thrown widely open, with the idea of alleviating his distress, and some nurses are inclined to be a little Spartan in their ideas of "fresh air." It must be remembered that cold is dangerous for these patients, and that it is sometimes wise to curb the propensity of either patient or nurse for free ventilation.

As to posture, the patient should be well propped up with numerous pillows, avoiding the common mistake of supporting the head and shoulders only, thus sharply flexing the spine. The whole spine from pelvis upwards should be supported and maintained at an angle of at least 45 degrees or, better still, at an angle approaching 90 degrees with the horizontal plane. The buttocks should rest on a biscuit-shaped air-ring underneath the draw-sheet. This posture may be maintained either by means of a knee pillow, or, if the patient is strong enough, with the help of an appliance known as the Hermit foot support—a padded board with an adjustable attachment to the foot of the bed. Owing to the weight of the oedematous feet and shins, the skin of the heels and ankles may become chafed and sore, and therefore may require protection. "Heart tables," such as Sister Rawson's, are often much appreciated by these patients, affording a well-padded support across the bed on which the patient may lean forward with folded arms. If such a table is provided with an adjustable padded support against which the patient can lean his forehead, so much the better. Some dyspnoeic patients find bed intolerable, and if so they may be allowed to spend the night well propped up in a really comfortable chair, preferably with the feet elevated. But they are best treated in bed altogether, and at all costs must be discouraged from frequent changes from bed to chair, and vice versa.

The diet should be simple and nourishing. The total amount of fluid in the twenty-four hours should not exceed 30 oz. Sugar is valuable, and may be given in the form of glucose lemonade (glucose 1/4 lb., two lemons sliced, water to the pint), or barley sugar may be sucked. Thin bread-and-butter, with jelly or honey, buttered or poached eggs, spinach, steamed fish, cream of chicken or of ham, and sweetbread are permissible, unless there is much gastro-intestinal disturbance. Weak China tea or cocoa may be taken, but not coffee. If the patient

is accustomed to take alcohol he should be given two to four ounces of whisky or brandy in the twenty-four hours. Port, champagne, and beer should not be allowed. Gin is sometimes useful by virtue of its diuretic property. Table salt should be avoided, and in chronic cases of cardiac dropsy a salt-free diet is sometimes beneficial.

Acute Congestive Failure

Acute congestive failure is almost always associated with auricular fibrillation, less commonly with auricular flutter or, rarely, with a paroxysm of paroxysmal tachycardia. The type associated with auricular fibrillation is the most amenable to treatment, this condition being the one which responds so dramatically to digitalis, given in adequate dosage. We are often told by the patient or his relatives that he cannot tolerate digitalis. As a matter of fact such an idiosyncrasy to digitalis is rare, though some tact may be necessary in order to persuade a patient to take digitalis when he has experienced toxic symptoms on previous occasions.

Now that the *B.P.* tincture of digitalis is pharmacologically standardized we can rely upon its potency, always provided that it has not been too long in bottle. Personally I now prefer to use digoxin, because this preparation is stable, and is quickly absorbed and quickly eliminated. Ventricular slowing begins within an hour of the administration of 1 mg. by mouth and is maximal within seven hours. It follows that intravenous administration is rarely necessary, but in very urgent cases 0.75 to 1 mg. of digoxin injected intravenously will produce ventricular slowing within ten minutes, which will be maximal within two hours. Great care must be taken not to inject this preparation extravascularly, as it is very irritant. If the ventricular rate is very rapid—150 or more—and if the patient is not already under the influence of digitalis, this drug may be given by the massive dosage method. To an adult weighing 140 lb. or more, 1.5 mg. of digoxin, or two drachms of the tincture, may be taken orally; for a lighter patient (for example, one of about 100 lb. weight) the dose would be 1.25 mg. of digoxin, or one drachm of the tincture. After six hours 0.25 mg. of digoxin, or 20 minims of the tincture, may be prescribed every six hours till the ventricular rate falls below 80. Digitalis must be stopped at once if the rate falls below 60; the appearance of coupling is an even more dangerous digitalis effect, for if digitalis therapy is continued when the rate is slow, with coupled beats, ventricular fibrillation and sudden death may supervene. Persistent vomiting or diarrhoea is also an indication for a rest from digitalis. When the ventricular rate has been controlled by digitalis, this drug must be persisted with in such dosage as will maintain the ventricular rate, by auscultation, at 70 to 80. This maintenance dose varies from patient to patient, and must be ascertained by experiment. Usually it is something like 10 minims of the tincture two or three times a day, or 0.25 mg. of digoxin once or twice daily.

If the patient is found by experiment to be really intolerant of digitalis, one of the strophanthus preparations may be employed. Of these the most reliable is ouabain, of which the average daily dosage is 0.25 mg. (1/240 grain) intravenously, or 2 to 4 mg. orally (in divided doses of 1 mg. each): 0.5 mg. of ouabain intravenously is roughly equivalent to 1 mg. of digoxin.

Venesection and Oxygen

If there is much engorgement of the neck veins, together with oedema of the lung bases and a large tender liver, venesection will give great relief. A French's bleeding needle may be inserted into one of the ante-cubital veins after the application of a sphygmomanometer armlet inflated to a pressure of 60 mm. Hg, and ten to twenty ounces of blood are withdrawn, according to the size of the patient. Personally, if they are obtainable, I prefer the old-fashioned leeches to venesection; by their use a more gradual and prolonged venous depletion is secured. It is usual to apply the leeches over the liver, and this is a convenient site, though no doubt they would be equally efficacious applied elsewhere. It is sometimes advisable to encourage free bleeding after the leeches have dropped off by applying hot fomentations to the punctures, and it is as well to warn the nurse on no account to attempt to stop the bleeding. I have even known a zealous house officer to apply adrenaline to, and actually to suture, the oozing leech punctures, thus attempting to defeat the therapeutic design. It is sometimes necessary to repeat a venesection or leeching in twenty-four to forty-eight hours' time, if signs of venous engorgement reappear.

Oxygen should be administered freely in most cases, and always if there is much dyspnoea. The degree of cyanosis is a less important indication for oxygen than is dyspnoea. The oxygen cylinder should be provided with an easily working reducing valve. The oxygen should be bubbled through warm water in a large flask, whence it can be delivered conveniently and efficiently to the patient by a tube held in his mouth. Better still, it can be administered via a nasal catheter. The warmed and lubricated catheter is passed till it nearly reaches the posterior wall of the nasopharynx. Its proximal end is lightly secured to the patient's forehead by a broad band of strapping. Through it oxygen can be administered continuously, or for periods of fifteen minutes hourly, the tube from the flask being disconnected between periods of oxygen administration. Pure oxygen should be given, and not a mixture of oxygen and CO₂. The administration of oxygen through the time-honoured tube and funnel is almost useless, and should not be allowed.

Symptomatic Treatment

Insomnia.—This is always troublesome in congestive failure, and measures should be taken to combat it. A few hours of peaceful sleep may effect a remarkable improvement in the patient's condition. For very restless and delirious subjects paraldehyde is far the safest and most effective hypnotic, and the patient will forgive its nauseating taste and odour if it brings him a good night's sleep. It is usually prescribed in hopelessly inadequate doses. It is best given by the rectum—in doses of 6 to 8 drachms well shaken up in four ounces of warm saline. If taken by the mouth, the simplest and most popular method of administration is to give 4 or 6 drachms of paraldehyde in one ounce of gin; alternatively, it may be prescribed in orange juice. For milder cases of insomnia a combination of bromide and chloral is indicated—for example:

R	Pot. brom. ...	...	...	gr. xx
	Chloral hydrat. ...	...	...	gr. xx
	Syr. zingib. ...	...	...	3 j
	Aq. ...	...	...	ad 3 j

Hypnotics of the barbitone group are not very satisfactory in this condition, and are, I think, more apt to produce mental confusion than is the above prescription. However, if bromide and chloral are not tolerated by the stomach, a tablet of medinal, 5 to 10 grains, may be taken, crushed and dissolved in a little hot water. It

has the advantage of acting rapidly. Morphine is not the hypnotic of choice in congestive, as it is in left ventricular, failure, but may be necessary if other sedatives fail, and particularly if pain is present, as, for example, in the event of pulmonary infarction. Sometimes quite a small dose (for example, 1/6 or even 1/8 grain) may be effective, especially if combined with chloral and bromide by the mouth. For a restless or maniacal patient who cannot, or will not, take paraldehyde, 2/3 grain of omnopon combined with 1/100 grain of scopolamine, hypodermically, would be indicated. Morphine should not be prescribed where there is much oedema of the lungs.

Cough.—This is inevitable in congestive failure, and the only really effective treatment of it is that designed to relieve the congestion. It is worse than futile to ply these waterlogged patients with expectorants, and thus increase the fluid which already clogs their bronchioles. Atropine is contraindicated because of its accelerating effect on the ventricular rate. For the dry tickling variety of cough which disturbs sleep, a codeine linctus is useful:

R	Syr. codein. phos. ...	...	...	m xxx
	Acid. citric. ...	...	...	gr. j
	Glycerin. ...	...	...	m x
	Spirit. chlorof. ...	...	...	m iij
	Mucilag. tragacanth. ...	...	...	ad 3 j

To be sipped in an ounce of hot water.

Regulation of the Bowels.—The bowels should be kept on the loose side by means of a morning dose of concentrated saline. This helps to relieve the portal congestion which is invariably present. One or two drachms of magnesium sulphate may be given in one ounce of hot water. If there is vomiting, one-half to one ounce of magnesium sulphate may be given in half a pint of warm water as an enema. It is important to avoid purgation, with its attendant disturbance. If constipation is troublesome, an enema is preferable to purges by the mouth. Diarrhoea, if present and exhausting the patient, should be checked by bismuth and opium:

R	Bismuth. carbonat. ...	...	...	gr. xl
	Tinct. opii ...	...	...	m x
	Pulv. tragacanth. co. ...	...	...	gr. v
	Aq. menth. pip. ...	...	...	ad 3 j

Pain.—This is not a common symptom of congestive failure, but it may be severe in the event of pulmonary or splenic infarction. Counter-irritation in the form of a mustard leaf, or dry-cupping, often leads to surprising relief, but morphine may be necessary. The pain of hepatic congestion usually yields to venesection or leeching.

Chronic Congestive Failure

Chronic congestive failure may follow acute failure with fibrillation. More commonly it is seen as a sequence to left ventricular failure with normal sinus rhythm—for instance, in hypertensive failure, or in the later stages of cardiovascular syphilis, or following a slow coronary occlusion, or in the late stages of emphysema or of pulmonary fibrosis.

The general principles of treatment have already been dealt with. If fibrillation is present, digitalis is prescribed (as previously indicated) in such dosage as will maintain the ventricular rate at 70 to 80. If, on the other hand, the heart is failing with a slow sinus rhythm, digitalis is not likely to be so effective, though worth a trial. It may be given in the form of Guy's pill:

R	Foliae digitalis ...	...	...	gr. j
	Pil. hydrarg. ...	...	...	gr. j
	Scillae ...	...	...	gr. j

One pill once or twice a day.

If a satisfactory diuresis is not thus obtained, 15 grains of diuretin (theobromine sodium salicylate) three times a day may be tried. This nauseating drug is best taken in the form of tablets, to be swallowed whole. It should not be given for more than three or four days at a time. If diuretin is not tolerated by the stomach, theocin sodium acetate, 3 grains dissolved in 1/2 oz. of peppermint water, may be successful, administered three times a day for three consecutive days weekly. This drug can also be injected subcutaneously, 3 grains dissolved in 15 minims of sterile water. But without a doubt the most valuable diuretic in these cases of chronic cardiac dropsy is either salyrgan or novurit, the introduction of which has revolutionized the treatment of this condition. The dose is 1 or 2 c.cm. intramuscularly, or preferably intravenously. The diuretic action of these drugs is enhanced by the administration of 90 to 120 grains of ammonium chloride in two or three ounces of water, given by the mouth half an hour before each injection of the salyrgan or novurit. The injection should be made in the morning, for if made in the evening the copious diuresis produced (which may be as much as five to ten pints in the twenty-four hours) interferes with sleep. Not more than two injections should be given each week, and if diarrhoea is produced this treatment should be discontinued, or a severe and possibly fatal enterocolitis may be produced.

It is important to restrict fluids to thirty or even twenty ounces in the twenty-four hours, and a strictly salt-free diet should be ordered. Gentle massage, and if the patient's condition warrants it, passive movements of the limbs, are also helpful.

Drainage of Oedema

If the oedema persists in spite of the above measures, an attempt should be made to get rid of it by drainage. The patients suitable for this form of treatment are those with tense but soft oedema, with shining skin. Long-standing cases with tough, indurated integuments seldom respond well to drainage, and if the skin is septic drainage is not safe. Nor should drainage be attempted with a restless or delirious patient. Before drainage, the patient should be kept sitting for some hours in a really comfortable armchair, with feet pendent. If he is not fit for this, the head of the bed may be raised on six-inch blocks. Southey's tubes may be used, at least four being inserted deep subcutaneously into the dorsum of each foot, or round the ankles. Care must be taken to ensure that the fine rubber tubing leading from each Southey's tube to the drainage receptacle is not blocked by clot; the nurse should be instructed to massage periodically each of these tubes downwards from tube to receptacle. With Southey's tubes there is less risk of cellulitis, and there is less attendant discomfort for the patient than is the case with drainage by means of multiple incisions, but there is no doubt that a more efficient and rapid drainage is obtained by the latter means.

To drain by incisions, six to twelve cruciform incisions are made into the deep subcutaneous tissues by a tenotomy knife. These incisions are made over the dorsum of each foot. This can be done painlessly under light freezing by an ethyl chloride spray. The incisions are covered by gauze soaked in glycerin of perchloride of mercury, which is useful for its hygroscopic as well as for its antiseptic action. The feet are placed in a large bowl if the patient is sitting; if he is in bed the fluid drained must be absorbed in voluminous wool dressings, frequently changed. If there is any considerable pleural effusion this should be aspirated, preferably combined with an air replacement by means of an artificial pneumothorax apparatus. It is not often necessary to tap the abdomen

for ascites in congestive failure, but this should be done if ascites is gross and does not quickly yield to diuretic treatment, combined perhaps with drainage of the legs. When so doing it is advisable to remove the fluid slowly through a fine cannula rather than to employ the large cannula usually provided for this purpose.

Insulin Treatment: Thyroidectomy

With the object of replenishing the store of glycogen in the myocardium, insulin, 5 units, may be prescribed two or three times a day, with a simultaneous dose of 10 to 15 grams of glucose by the mouth, in the form of barley sugar or as glucose lemonade. This form of treatment is, in my experience, beneficial in the more chronic cases, and is sometimes worth a trial even in acute cases.

During the last two years the operation of total thyroidectomy has been advised for the treatment of chronic congestive failure where other methods of therapy have failed. Dramatic results have been described, particularly in America. This method of treatment is still *sub judice*, and obviously it should not be employed until it seems certain that the patient is going downhill and has ceased to respond to medical treatment. It must be remembered that some patients with chronic congestive failure make, with medical treatment only, most remarkable and quite unexpected recoveries after many months of desperate illness. On the other hand, the operation risk of a total thyroidectomy in these cases is, in the hands of an expert surgeon, not as serious as might be anticipated, so that this treatment seems worthy of trial in carefully selected cases. At the same time I must confess that in my own very limited experience the results have not been very encouraging. When, however, congestive failure and auricular fibrillation are associated with a toxic goitre, the surgeon's aid can and should be invoked, usually with gratifying results. After the ventricular rate has been controlled by digitalis, Lugol's solution being administered simultaneously in a dosage of 10 to 15 minims, three times a day, a subtotal thyroidectomy is usually successful in restoring normal rhythm. If normal rhythm is not restored by the operation, a short course of quinidine after the operation rarely fails to bring this about. Not infrequently this treatment is capable of transforming an apparently hopelessly bed-ridden and almost moribund patient into a man or woman who is fit for quite a hard day's work.

After-care

Complete rest in bed should be enjoined for two or three months after all oedema and other evidences of congestive failure have disappeared. When the patient is allowed to move about in bed, and subsequently to get up, increase in the amount of effort permitted must be very cautious, and he must be treated as an invalid for at least six months.

Further information about the fourth International Hospital Congress at Rome from May 19th to 26th has now been received, and it should be noted that the last day for enrolling has been postponed until April 30th. A handsomely illustrated programme of the congress has been issued in various languages including English, the main topics to be discussed being the hospital in the public health scheme, general organization and equipment, hospital protection in war time, and the best disposition of wards and departments from the point of view of the patients as well as of administration. Tours of the chief Italian cities have been arranged for the periods before and after the congress sessions. All inquiries should be addressed to the general secretary of the congress, Ospedale di S. Spirito, Rome.