into the subclavian vein. Here a probe was passed with some difficulty; a nodule was felt at the junction, and on microscopical examination a collection of tumour cells surrounded and constricted the duct, and was projecting into the subclavian vein (Fig. 2 on Plate). A diagrammatic representation of the specimen is given below.

A, Thoracic duct. B, Growth. C, Subclavian vein.

Diagram of post-mortem specimen (see Fig. 2 on Plate).

The growth was found to be a retroperitoneal sarcoma of the small, round-celled type. It was growing densely in front of the lumbar vertebrae, invading the mesentery and the left kidney from below, and commencing to infiltrate the uterus and ovaries. There was no growth, either visible or microscopical, in the lungs or mediastinum.

A very small amount of fluid had collected in the abdominal cavity, though four days had elapsed since the abdomen had been tapped. This was to be contrasted with the condition found in the thorax, where, in spite of withdrawal of fluid on the day of death, about one and a half pints were present in each pleural cavity. This confirmed the belief that the main factor in determining the chylothorax had been the aspiration due to negative pressure in the thorax.

Summary

A case of bilateral chylothorax followed by chylous ascites is described. The symptoms corresponded with a condition arising from partial occlusion of the thoracic duct by a growth, and the diagnosis was confirmed at the post-mortem examination, when it was found that the junction of the thoracic duct with the subclavian vein was partially occluded by a metastatic deposit from a retroperitoneal sarcoma. The points of leakage in the chest and abdomen were believed to be small and valvular, but they were not definitely located at post-mortem examination either by dissection or by injection of lipiodol.

Grateful appreciation is herewith expressed to Dr. Hamilton Ross for permission to study this case, which was under his care. Dr. H. A. Cookson, pathologist, is thanked for providing laboratory facilities, and for his helpful advice and criticism. Dr. Sharp kindly provided clinical notes and the supply of fluid from numerous tapplings. Dr. Paige Arnold and his staff gave assistance by taking the x-ray and other photographs.

REFERENCES

- ¹ Wells, H. G.: *Chemical Pathology*, fifth edition, Philadelphia, 1925.
- ² Powell, R. D., and Hartley, P.: *Diseases of the Lungs and Pleura*, sixth edition, London, 1921.

An international medical week will be held at Montreux, Switzerland, from September 9th to 14th, to discuss subjects of practical interest to general medical practitioners. The discussions will be introduced by leading authorities from various countries. Sir Henry Dale will be one of the lecturers, the others including Professors L. Rajchman of the League of Nations, E. Bürgi of Berne, J. Holmgren of Stockholm, C. Nicolle of Tunis, Ximenez Diaz of Madrid, H. E. Sigerist of Baltimore, and M. Askanazy of Geneva. The subjects for discussion will be notified later. The fee for membership is 10 Swiss francs, and it will include payment for the report of the "Transactions." Further information may be obtained from the secretary of the International Week, Klosterberg 27, Basle.

TUBERCULIN IN THE TREATMENT OF CUTANEOUS TUBERCULOSIS

BY

H. S. BURNELL-JONES, D.P.H. Oxon

(With Special Plate)

Details are recorded in the *Proceedings* of the Royal Society of Medicine for January, 1934, and in the *British Journal of Dermatology and Syphilis* for February, 1934, of the treatment of three cases of cutaneous tuberculosis with tuberculin, the results of which encouraged continuation of the investigation. An account is presented here of three further cases similarly treated (out of a series referred from the skin department of the Royal Northern Hospital), for the purpose of encouraging further work and investigation.

Three tuberculins have been used—T.A.F., B.E., and T.R., obtained from Messrs. Bayer Ltd. In changing from one tuberculin to the other no particular rule has been followed, and no definite evidence has accumulated to indicate that one tuberculin is superior to another. Latterly I have been inclined to modify this statement in favour of T.A.F. and B.E. One fact that stands out pre-eminently—as pointed out by others—is that each case is a law unto itself with regard to intervals between injections, increase and decrease of dose, and the change of tuberculin, and that clinical insight and experience have to be depended upon.

Technique Followed

The site of injection is of importance; patients showing local reactions on the arm will often show no local reaction if injected between the shoulders. The leg appears to be as sensitive as the arm. After experimenting with increases varying from 5 to 50 per cent., results have led to the opinion that 10 per cent. (or thereabouts) is the most satisfactory increase for dosage. The effectiveness of this small increase in skin cases, as compared with that feasible in pulmonary cases, may be due to a smaller quantity of lysin present in the tissue. In contradistinction to pulmonary tissue, the skin is not well flushed with blood, which results in a smaller supply of oxygen and pabulum, and perhaps accounts for relatively slower growth of the bacillus in this situation.

For the small doses a Koch's syringe is most convenient. The varying doses can be measured at the time of injection from the 1 in 10, 1 in 100, and 1 in 1,000 dilutions with the ordinary 0.1 c.cm. pipette graduated in tenths. For doses of 0.2 c.cm. or over the ordinary 1 c.cm. tuberculin syringe, graduated in tenths, is quicker. The dose of tuberculin is first drawn in, followed by the diluting fluid. The diluent is normal saline with 0.2 per cent. phenol, not 0.5 per cent. phenol. To the tongue 0.5 per cent. is distinctly appreciable, and in patients hypersensitive to phenol may account for the doubtful positive reaction sometimes seen after an intradermal test, which is followed by negative subcutaneous test injections when a positive temperature reaction might have been expected. No cleansing of the skin is done and no dressing is applied. Distilled water is used in the small metal sterilizer to prevent deposition of calcium salts in the syringes, needles, and pipettes. The glass pipettes are sterilized in a naked flame after rinsing with the distilled water in the sterilizer. The water should not be ejected back into the sterilizer. From time to time it is useful to clear the pipettes with pure nitric acid.

Routine intradermal and subcutaneous test doses are given in all cases. Valuable information is obtained by these preliminary injections as to the degree of allergy and the approximate dose with which treatment should be

begun. The small doses used in the early stage of treatment are given twice weekly, but I have come to the conclusion that in treating cutaneous cases an interval of one week or more is advisable with doses smaller than those used for pulmonary cases. The patients take their own temperatures three times a day, allowing the thermometer to remain under the tongue not less than five minutes. Of existing charts Rigden's is the best (H. K. Lewis and Co.). This chart lasts two months, but a specially designed chart would be better.

If suspension of treatment is indicated, particularly in cases showing high temperature reactions not, so far as can be ascertained, due to tuberculosis—I have had two such cases—injections are still given, but diluting fluid only is used. This allows control of the patient, maintains encouragement, and also enables one to demonstrate whether the rise is associated with administration of tuberculin. Professor Jadassohn writes:

"The reason why the treatment with tuberculin has not yet given better results with tuberculoderms may be that the right method has not yet been found; and this is perhaps because as yet very little is known of the nature of the action of tuberculin. We must not forget that our knowledge concerning this question is still very incomplete. I must once more emphasize that here we have to deal with the very important unknown factors—namely, the reason why the tuberculin allergy develops in such different ways in different individuals. The difficulty of solving this question is enormous; nevertheless, through patient research and industrious observations we must do our best to find its answer."

Case I

Lupus Vulgaris of Face

F.M., a female aged 27, gave a history of fourteen years' duration. The mother first noticed a tiny pimple on the middle of the left cheek. Within a year it had spread over the cheek, notwithstanding medical treatment. She then had zinc ionization for two years, by which time the eruption had reached the left ala. The treatment was given at the School Clinic, Nottingham, every three or four weeks. Severe reactions occurred. She then went to different doctors for about three years. This was followed by the sun-ray and Kromayer lamp treatment at a general hospital, the patient reporting that "it thinned it down a lot, but it continued to spread." There had been continual progress of the disease when she came to the Royal Northern Hospital.

On February 23rd, 1934, a positive reaction followed intradermal injection of 0.0001 c.cm. T.A.F. There was no focal reaction. Treatment was begun with 0.0001 c.cm. B.E. on April 5th. Although this caused no temperature reaction, the patient's chart led me to reduce the dose to 0.00005 c.cm. on April 16th. I got back to 0.00011 c.cm. by June 7th. (Note on June 4th: "Margins still raised, no crusting now, some areas are normal in colour.")

This patient has always exhibited an irregular temperature chart, and on July 5th I changed the tuberculin from B.E. to T.R., giving the same dose (0.0001 c.cm.). The temperature went to 99.8° F. instead of 99.4° with the 0.0001 c.cm. of B.E. On July 13th the patient went to Nottingham for a change, and the treatment was not continued until August 16th, when 0.00001 c.cm. T.A.F. was given. On October 11th 0.00007 c.cm. T.A.F. caused a reaction to 100.6° F., with a very marked track reaction and burning of the face for four days. There was marked scaling. It must be mentioned that pyotropin was applied on September 5th for the first time. On January 24th, 1935, the note reads: "No sign of breaking down; scaling in the lower half of the eruption, where healing is not as advanced as in the upper half."

From September 6th, 1934, she has had weekly injections. Although the doses are so small the track reaction is always very marked. By April 11th, 1935, the dose was 0.000325 c.cm. T.A.F. It must be mentioned that this patient has also had Finsen light treatment since the latter part of October, 1934. The weight in March, 1934, was 8 st. 13 lb.; in April, 1935, 9 st. 8 lb. The skin condition is illustrated on the Special Plate.

Case II

Lupus Verrucosus of Left Hand

W. J. O., a male aged 58, was first seen in May, 1934. This patient had lesions over the metacarpo-phalangeal joint of the left index finger, on the thumb-joint, and on the skin between these areas. A thick, hard, infiltrated pad in the palm prevented approximation of the thumb and index finger. This condition had existed for twenty years, and dated from cutting his thumb with a knife. The patient had rheumatic fever at the age of 6 years, and the heart's action was weak.

Intradermally, 0.0001 c.cm. T.A.F. gave a weak positive reaction. Treatment started on May 24th, 1934, with 0.001 c.cm. B.E. By June 4th the dose had reached 0.003 c.cm. and the patient possessed much more energy. The lesions had paled by June 28th. On July 2nd he had 0.012 c.cm. B.E. The temperature excursions varied between 97° and 98.4°. Treatment has been interrupted in this case by catarrhal attacks. In 1932 the patient spent fifteen weeks in a sanatorium, and the sputum was positive. Notes on August 13th state: "Small nodule to ulnar side of the lesion, over the second metacarpo-phalangeal joint. Red area appeared over the first metacarpo-phalangeal joint. Not feeling well lately, pain over cardiac area with palpitation after food. Sleeps better since having injections, and has gained over 3 lb. in weight."

On August 27th, with the dose at 0.009 c.cm. T.A.F., the hand showed definite improvement. There was less subcutaneous infiltration, and the thumb and index finger could be approximated. This had previously been impossible for two and a half years. By September 20th there was considerable scaling (dose 0.04 c.cm.). On October 1st he reported that the catarrh and nervous condition had disappeared. When the dose reached 0.14 c.cm. T.A.F. the patient had Finsen light treatment in addition, the first application being given on October 22nd. Tuberculin injections were continued bi-weekly, and the temperature reactions remained between 97° and 98.6°, with very seldom any local reaction at the site of injection. Since the dose reached 0.25 c.cm. T.A.F. on November 5th, the injections have been given once a week. On December 20th he did some hard work without it causing swelling or aching of the hand, which was unusual (dose 0.375 c.cm. T.A.F.). On March 28th, 1935, the dose was 0.525 c.cm. T.A.F. The weight in May, 1934, was 9 st. 10½ lb.; in April, 1935, 10 st. 1½ lb.

Case III

Lupus Verrucosus of Both Hands

W. M., aged 57, an official killer of tuberculous cattle, stated that the condition complained of had started about August, 1932, on the dorsal aspect of the left hand, over the second metacarpal bone, as a "wart." When first seen, on February 22nd, 1934, the disease extended for two and a half inches above and below the wrist-joint, right across the hand, and around to the thenar eminence. On the right hand there was a wart-like area, two inches in extent, over the dorsal aspect of the fourth and fifth metacarpal bones. This started about June, 1933, and began to get rapidly worse seven months later.

On February 22nd a marked track reaction, raised, and three-quarters of an inch in diameter, followed an intradermal injection of 0.0001 c.cm. T.A.F. Treatment was started with 0.0001 c.cm. B.E. on February 26th; on March 12th, with 0.0009 c.cm., temperature reactions appeared with marked track reactions. By this date the left hand was drying and scaling; improved and painless movement was also possible. On April 30th 0.001925 c.cm. B.E. was given. The local reactions were now becoming less severe, and also the temperature excursions to 100° or over had steadied down to 98.2°. By May 24th the right hand had improved considerably. The eruption had flattened, although it was still warty in appearance. Two areas on the left hand exhibited marked improvement, but there were numerous soft areas with pus underneath, one-eighth to a quarter of an inch in diameter. The patient had been doing a lot of digging. On July 5th the dose was 0.011 c.cm. B.E., and at this time the left hand was more swollen and inflamed; there was also an area of extension the size of a penny around the inner side of the wrist. The lesion on the right hand was much reduced in thickness and was also dry.

About July 21st Dr. H. C. Semon scraped both hands, and on September 13th injections were again given, beginning with 0.005 c.cm. T.R., reaching 0.12 c.cm. by December 31st, without track reactions or temperature above 98.4°, and the left hand was much improved, there being no open wound and no pain. After 0.23 c.cm. T.R. on January 31st, 1935, the patient had 0.1 c.cm. T.A.F. on February 4th, followed by no track reaction, but a temperature of 100.2° and also slight general reaction. The dose on April 15th was 0.35 c.cm. T.A.F. The weight on February 22nd, 1934, was 12 st.; on December 17th, 12 st. 11 lb. The skin condition before and after treatment is illustrated on the Special Plate.

This patient reported a similar condition on the left arm of his son, aged 32, appearing just before his own eruption. It extended three inches across the outer side of the arm and three inches above the elbow-joint. It was not red and not raised. The veterinary surgeon treated it, but without change. The son was accidentally shot in September, 1932.

General Conclusion

The answer to the question, "Will tuberculin cure lupus?" is unknown, but the arrest of disease and the improvement seen in these cases are strong indications that the work should be continued. The fact should be mentioned that in one of the earlier cases (not recorded) seven months' treatment was given before any change in the lesions could be recognized. I have heard a doctor condemn tuberculin after using it on one case for three months only. All the cases treated were of long duration before they received tuberculin. It would be of inestimable value if cases could be treated in the early stage; we are more likely to learn from early cases than from advanced cases.

F. M. (Case I), notwithstanding the fact that allergy was so marked, continued to improve with small doses. The pertinent question asked me by an expert, "If you get improvement with small doses, why use large ones?" naturally comes into one's mind. I only proceed to large doses if the patient tolerates them without rise of temperature or unpleasant general reaction. This tolerance varies. One patient will sometimes feel ill after 0.1 c.cm., whereas another will feel no discomfort after 1.6 c.cm. I do not look upon any particular dose as the maximum for any tuberculin. The most sensitive case I have yet had has only recently come under treatment; the patient, a female aged 22, has extensive glandular enlargement in the neck and tuberculides on the hands. As little as 0.000005 c.cm. of tuberculin sent her temperature up to 101° F.

Increase of weight almost invariably accompanies administration of tuberculin. Sooner or later one finds the improvement is not as rapid as at first. I now look upon this as a sign for temporary suspension of treatment, or for extension of the interval: in any case it behoves caution. It does not occur when any particular dose has been reached.

Illustrative of the clinical lessons incidental to tuberculin treatment is a case reported in the issue of the *Proceedings* of the Royal Society of Medicine above mentioned, in which failure to respond to tuberculin in the usual way led to the correction of a wrong diagnosis of "tuberculosis" to the correct one of "syphilis."

Finally, I would call attention to the remarkable differences in the blood films one finds in the different types of cutaneous cases, and suggest that workers should do monthly differential blood counts on each case.

I am indebted to Miss Clark of the x-ray department, Royal Northern Hospital, for having the photographs taken, and to Dr. Robert Carswell for advice and assistance in the preparation of this article for publication. Attached is a list of authors whose work I have found useful.

BIBLIOGRAPHY

- Bandelier and Roepke: *Tuberculin in Diagnosis and Treatment*. Translation by Wilfred B. Christopherson, 1913.
 Bredeck, J. B.: "Tuberculin and the Schilling Blood Differential Count in the Diagnosis and Classification of Tuberculosis," *American Review of Tuberculosis*, April, 1933.
 Carswell Robert: Papers on tuberculin, 1933.
 Gillespie, John R.: *A Rational Method of Using Tuberculin*, 1918.
 Jadassohn, J.: "Remarks on Tuberculin in Dermatology," *British Journal of Dermatology and Syphilis*, 1929.
 Riviere and Morland: *Tuberculin Treatment*, 1912.
 Semon and Burnell-Jones: "Tuberculin in the Treatment of Cutaneous Tuberculosis," *British Journal of Dermatology and Syphilis*, 1934.
 Wilkinson, W. Camac: *The Principles of Immunity in Tuberculosis*, 1926; *Tuberculin, Vindicated by Technique*, 1933 (the table on page 61).

INTUSSUSCEPTION IN THE ADULT DUE TO TUMOURS

A NOTE ON FIVE CASES

BY

E. N. MACDERMOTT, B.Sc., M.D., F.R.C.S.I.

HONORARY SURGEON, CENTRAL HOSPITAL; PROFESSOR OF THERAPEUTICS,
UNIVERSITY COLLEGE, GALWAY

(With Special Plate)

Intussusception in the adult is of sufficient rarity to render reports of isolated cases interesting; the occurrence of five in the author's practice in the past year seems, therefore, worthy of report.

Case Reports

CASE I

A male, aged 28, was first seen by me on October 10th, 1933, complaining of abdominal pain, vomiting, and constipation for ten days; he stated that he had had similar attacks during the previous three and a half years, and that he had passed blood and "slimy stuff" at intervals. During the two days before I saw him he had received three enemata, with one constipated and two "coloured" results. On examination he had a temperature of 97.8° and a pulse of 100. His abdomen was distended and resonant to percussion; the distension made accurate palpation impossible, but there was a sensation of resistance in the right side. A tentative diagnosis of intestinal obstruction was made, and the abdomen was opened through a right paramedian incision under ether anaesthesia. On opening the peritoneum some blood-stained fluid escaped. An intussuscepted mass was found in the right hypochondrium, small gut entering large gut, and caecum and appendix not being visible. The small gut was enormously dilated, almost to the size of the colon, outside the intussusception. The latter was reduced, when it was found that an originally enteric intussusception had passed through the ileocaecal valve and, carrying this along with it, had intussuscepted the caecum and appendix into the ascending colon. The whole mass was about eight inches long. There was a polyp about nine inches above the ileocaecal junction; this polyp had occupied the apex of the intussusception. Above the polyp the bowel was dilated to about the diameter of three fingers, while below it was collapsed and contracted to the size of the little finger. As resection and anastomosis did not seem advisable, even if feasible, a McBurney incision was made on the right side, and the polyp, with about three inches of gut on each side of it, extruded and resected after suturing the two limbs to each other and to the peritoneum; clamps were left on the intestine until the following day. Before closing the abdomen one other polyp was located in the small gut and demonstrated to the assistant. The abdomen was then closed.

Post-operatively the patient did very well, but there was a tendency for the gut of the proximal segment to prolapse through the enterostomy opening, which interfered with spontaneous closure. In mid-November the redundant bowel was resected and the spur clamped. Faeces were passed per rectum on November 24th. On December 9th an attempt was made

H. S. BURNELL-JONES: TUBERCULIN IN THE TREATMENT OF CUTANEOUS TUBERCULOSIS

FIG. 1.—Case I (fourteen years' history). March 9th, 1934; before treatment.

FIG. 2.—Case I seven months later.

FIG. 3.—Case III (eighteen months' history). February 27th, 1934; before treatment.

FIG. 4.—Case III some eight months later.