Adherent Pericardium

Adherent pericardium hardly comes into our subject, as the deficiencies arising from this condition are those associated with its chronic effect on the heart. Many cases of pericarditis are followed by some degree of adherence between the visceral and parietal pericardium, but this is not enough to be very important in itself, or sufficiently widespread to prevent subsequent attacks of pericarditis and even of pericardial effusion. But the term is more often applied to the more extreme condition, where both layers of the pericardium are fixed to the surrounding mediastinum as the result of an old extensive mediastino-pericarditis. This adds a good deal to the work of the heart, and is known to be frequently associated with a very high degree of cardiac hypertrophy. No satisfactory local treatment has been devised, and the correct treatment is that of the underlying damaged heart muscle.

EVIPAN SODIUM IN GENERAL PRACTICE

BY

C. M. DOUGLAS, L.R.C.P. & S.

Notwithstanding the many excellent reports on the use of evipan sodium which have appeared in the medical press, I feel that a general practitioner's experience of just over 180 cases may be of some interest. My excuse is that the reports on its use mainly come from hospitals and clinics, where anaesthesia is induced under ideal conditions. The following notes are intended to supplement these reports from a general practitioner's point of view.

In the "little black bag" are carried a 10 c.cm. Record syringe and needles in a spirit-proof case, a rubber-capped bottle of doubly distilled sterile water, a thin rubber tube, with a pair of artery forceps, and, although I have never had occasion to use them, a full box of coramine and icoral.

I should like to criticize the warning which has appeared in so many reports that "evipan should never be administered single-handed except under exceptional circumstances." I imagine that this has discouraged its use by many country doctors. The above statement appeared in a very helpful report on 1,000 cases by Jarman and Abel, under the subtitle "Dangers—The Jaw." One would take it, then, that the danger of the jaw falling forward and interference with respiration resulting was the chief reason advanced against using the anaesthetic single-handed. I first used evipan in July, 1933, at a time when there was little literature published in this country, and, naturally, I went very cannily. Since I hoped for complete muscular relaxation, I expected to find that the jaw also became slack, and for my first fifty cases I always had a colleague with me, who paid particular attention to the jaw. Now I have no misgivings, whether I am alone or not. In my notes I find that in one-third of the cases the mouth remained tightly closed. These were patients who would have had gas and oxygen if it had been available—that is, only a very small dose was given. In about one hundred cases the jaw dropped enough to expose the teeth, and I found that it took a little pressure to depress the jaw enough to cause any interference with the breathing. Anyone who has given about 6 c.cm. for extraction of a few difficult teeth will bear me out when I say that the dentist has often asked whether the jaw could be opened any wider. In the remainder it has been enough to push the jaw up and hold it there for half a dozen breaths. In no case was there any sagging which caused me to ask anyone to devote his whole attention to the jaw. In my experience it requires a complete dose of 10.5 c.cm. to cause the masseters to relax to any extent.

In only one case did the condition of the patient give rise to any anxiety, either during or following the opera-

tion. This was in a woman of 26, who had 6 c.cm. for curettage. For three days she complained of her eyes feeling as though she had been staring at the sun. Nothing abnormal was found on examination; fundi and visual fields were normal, and the condition cleared without further treatment.

During the last two years I have used evipan in all types of cases which a general practitioner would meet in the everyday round. My youngest patient was a boy of 13, my oldest a woman of 63. I need not enlarge upon the number of times that a short anaesthetic is necessary during an average week in a practitioner's life. It often means one of two things—trying to get a colleague, who may be on his rounds, or giving ether and doing what is necessary with one eye and both ears on the patient. Single-handed one may have to deliver with low forceps, which often means a faulty anaesthetic, poor asepsis, and an empty feeling in the pit of the stomach. Now one scrubs up, gives the woman enough evipan, scrubs up again leisurely and thoroughly, and with the nurse, or sometimes a "friend" holding the leg in just the required position, delivers with ease and comfort just as one did when a hospital resident. The analgesia which persists for some time allows one to do any necessary perineal repair without hurrying. This, too, seems to me an important point in general practice. Faulty perineal repairs are more often due to dislike of keeping or putting the patient under an anaesthetic, again single-handed, than to lack of skill. I used to give chloroform and ether in practice with the hope that everything would be all right, but now I look forward to minor operations when using evipan. There is a lack of strain which has to be experienced to be believed. The time saved, too, is a great help to the busy practitioner. One does not like leaving a patient profoundly unconscious as, say, after an induction under ether, even in charge of a nurse, until she is partly round. Whereas, under evipan she will respond to a request to open her eyes before the instruments are washed and dried.

The pleasant and quick induction and the recovery, which compares at least favourably with ether, helps in one's relation with a nervous or apprehensive patient, and makes future treatment more trustfully received. I have no wish to cry the praises of evipan in an uncritical mood, but after two years' experience I feel that in most of the cases where one requires a short anaesthetic evipan holds a high place.

I should think that most people modify various techniques according to their own experience. I find that 1.5 to 2.5 c.cm. injected quickly, followed by a pause of forty seconds, gives a pleasant induction and a minimum of jerking. If one is working single-handed it is wiser to inject while kneeling on the patient's palm just in case there should be any jerking.

In my notes I have records of patients who showed a definite delay in becoming unconscious, from 5 to 10 c.cm. taking up to seven minutes to cause unconsciousness. It seemed to compare with what occasionally happens when using nembatal, the patient showing no response to several capsules, until suddenly they take effect—sometimes dramatically. In none of these cases, however, was the return to consciousness delayed beyond what I have come to consider as normal.

In thirty-one cases where the patient was young and healthy I have injected 10 c.cm., although the patient fell asleep with 2 to 3 c.cm. and the operation required a space of time which would have been attained by 5 to 6 c.cm. In nineteen of these the post-operative headache, so commonly complained of, was very slight, and the sensation which many patients describe as "that horrible feeling that you are awake yet can do nothing, not even open your eyes," was not so pronounced. It may be that a patient may, for a pleasant recovery, require more than the minimal dose which the operator requires to work in comfort. In fifteen cases it has been expedient to give a further injection, and I have found it necessary to inject very slowly, since the patient often goes very deep with a very small dose.

I have never used premedication—obviously in many cases it has been impossible. Injecting on a full stomach

seems to make no difference, although, if there is time, I feed the patients on tinned fruit juice and glucose, and only two of the 180 have vomited. One case had been dieted, the other had not.

One fact remains which I have not seen mentioned in reports: many patients, as soon as the injection is completed, adopt a peculiar position of the arms. These become stiff, flexed at the elbows and wrists, with the fingers extended and the thumb adducted across the palm.

Evipan is still in its 'teens, but from my own experience I feel that when it becomes more widely used there will be fewer grey heads among young general practitioners.

England and Wales

Red Cross Clinic for Rheumatism

The report of the medical board for the year 1934 shows that the activities of the British Red Cross Society Clinic for Rheumatism, in Peto Place, N.W.1, have been fully maintained. The range of routine investigations and treatments has continued to expand, and researches are being made in several directions in the attempt to solve some of the problems presented by the different rheumatic diseases. Three morning, five afternoon, and four evening clinics have continued to be held weekly throughout the year by the visiting physicians, in addition to those held by the specialist staff. There were 89,755 attendances and a total of 132,619 treatments in the general out-patients' department, and 8,901 attendances and 16,354 treatments in the private patients' department. There are still a considerable proportion of advanced cases sent up to the clinic, although the importance of referring early cases is becoming more and more appreciated. There is a very real need for adequate facilities for in-patient treatment under the supervision of the visiting physicians at a site, if possible, near the clinic, where any special or detailed investigation as well as any form of physical or other type of treatment that would be contraindicated in an out-patient department can be carried out. The lack of these facilities is proving a definite handicap in the full and effective treatment of the more serious and chronic types of the rheumatic diseases. The provision of a certain number of beds would also obviate the delay that now occurs before patients are able to have the operations advised. There are many patients, especially those suffering from the early manifestations of rheumatoid arthritis, who would derive incalculable benefit if they were lifted out of the depression of their home surroundings and the strain of their everyday life and sent to the country for a lengthy period of convalescence. The clinic tries to take every advantage of the facilities offered by the different convalescent homes, but it is felt that the equipment of a building within reasonable distance of London, of suitable aspect and elevation and accommodating twenty patients and the necessary staff, with grounds large enough for extension, would prove an inestimable boon.

Coroners' Inquests in London

The number of deaths reported to coroners in 1934 in the County of London was 8,548, as against 8,533 in the previous year. Of these, 2,351 took place in hospitals and other institutions, and 721 in mental hospitals. The coroners decided to hold inquests in 3,455 cases, as against 3,543 in 1933. A post-mortem examination by direction of the coroner was carried out in 82 per cent. of the inquest cases, and in 64 per cent. of the other cases in which no public inquiry was necessary. A verdict of murder was returned in eleven cases; four inquests were held in connexion with executions, and there were 690 suicides, a decrease of seventy on the previous year. The number of people who met their death by accident was 1,925, as against 1,965 the year before. There were fifty deaths by drowning. Verdicts of "death from natural causes" were returned in 543 cases, and in seven the verdict was "cause of death unknown." Inquests on

newly born children increased from forty-four to forty-eight; in seventeen instances it was decided that the children were stillborn. Excessive drinking accounted for fifty deaths, as against fifty-four in 1933.

Central Midwives Board

At the May meeting of the Central Midwives Board for England and Wales, the application of the public assistance officer of Essex and the matron of the Old-church Hospital, Romford, for permission to train fourteen pupils per annum instead of ten was granted, but it was decided to point out that when the training rules, which are at present under revision, come into operation, it may be necessary for the Board to reconsider its decision. A letter was read from the Clerk of the London County Council setting forth the results of the Council's experience of the treatment by midwives of cases of asphyxia by the "sparklet" carbon dioxide apparatus. The following resolution was passed at the meeting, and a copy is to be forwarded to the Ministry of Health:

"That the Board, after consideration of Recommendations 49 (b) and (c), page 78, of the Departmental Committee Report on the Training and Employment of Midwives (1929), and in the light of further experience in the hearing of cases of alleged breach of its rules by midwives, is of opinion that all new appointments to non-medical positions for routine inspection of midwives should be filled by midwives who are State-registered trained nurses with wide experience of practical midwifery."

Scotland

Rheumatism in Children: Investigation at Edinburgh

Speaking at the annual meeting of the Royal Hospital for Sick Children at Edinburgh, Mr. Colin Black, chairman of directors, said that a biochemical department had been erected in the grounds of the hospital, and a consulting biochemist and trained assistant appointed. An investigation had been carried out in regard to the serious and crippling effects of rheumatism. The importance of this subject might be appreciated from the fact that over 25 per cent. of the young children who developed acute rheumatic infection did not survive to adult life, while a further 30 per cent. were permanently invalidated as a result of damage to the heart. The staff was still working at this investigation, which consisted of an analysis of cases treated in the wards of the hospital during the past fifteen years, with a "follow-up" to discover the late effects of acute rheumatism. The medical staff had recommended that vigorous measures should be taken to combat rheumatism, and had suggested that any scheme to deal with this problem in children should have as its three main objects: (1) to prevent rheumatic disease; (2) to prevent recurrence in children already affected; and (3) to care for the children already crippled by disease of the heart. By a gift of Lord Forteviot it had become possible for the directors of the hospital to institute a convalescent home in addition to the one at Gullane, and it was hoped to inaugurate this new home during the present year.

Plaques at Edinburgh Mental Hospital

Six bronze plaques in memory of pioneers in the humane treatment of mental disease were unveiled recently by Sir Arthur Rose, chairman of the General Board of Control for Scotland, at the entrance to the Royal Edinburgh Hospital for Mental and Nervous Disorders. The idea of placing these commemorative plaques was conceived by the late Professor Robertson, who in 1930 caused a bust of Philippe Pinel to be unveiled there by Monsieur de Fleuriau, the French ambassador. The present six plaques are arranged in two groups of three flanking this bust. They represent Dr. William Tuke, Dr. Andrew Duncan, Florence Nightingale, Dorothea Dix, Dr. Gardiner Hill, and Dr. Campbell Clark, and have been presented by various groups of persons in Britain and America who are