TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some of the major medical disorders met with in general practice.

THYROTOXIC HEART DISEASE

BY

GEOFFREY BOURNE, M.D., F.R.C.P.

Cardiac symptoms, slight or severe, are present in every case of thyrotoxicosis; indeed, cardiac complications constitute the gravest dangers of the disease. Before discussing the details of treatment, it will be helpful to describe briefly the manner in which the heart is affected.

Two main pathogenic factors are at work: persistent overdriving and toxæmia. The manner in which these produce their effects is briefly as follows. The hyperthyroidism, or dysthyroidism, results in symptoms of sympathetic stimulation, one manifestation of which is the increased basal metabolic rate. The tissues therefore need more oxygen, the circulation rate is much accelerated, and tachycardia is produced: ultimately some generalized cardiac hypertrophy also develops. At the same time the excessive or abnormal thyroid secretion has a toxic action on the heart muscle, and this becomes dilated to a greater or less extent and sooner or later shows signs of failure. A secondary effect of this toxæmia upon the heart muscle is an increased excitability, which is seen by a tendency to premature beats, auricular fibrillation, and, more rarely, auricular paroxysmal tachycardia or flutter.

The degree of failure varies greatly in different cases. It tends to be more severe in the secondary type of Graves's disease, or toxic adenoma of the thyroid, as it is sometimes called. But this is not always true, and no useful therapeutic purpose is served by regarding primary and secondary Graves's disease as two distinct entities. The fundamental basis of therapy is the same in both cases. The first step in successful treatment is careful evaluation of the degree of failure. In slight cases tachycardia is the only sign, and palpitation and dyspnoea the only symptoms present. In more severe cases the heart is enlarged, the degree of enlargement being proportional to the degree of failure. In the most severe cases auricular fibrillation is generally present. This irregularity is not only symptomatic; it adds mechanically to the severity of the heart failure.

When the source of the thyrotoxicosis has been removed the cardiac insufficiency disappears in a remarkable way. The best means of removing the source of poisoning is a surgical operation. X-ray treatment is less satisfactory in severe cases owing to the greater difficulty of estimating exactly how much of the gland has been destroyed. In slight cases it is a valuable method of treatment. Should surgery be decided upon, the medical preparation of the patient becomes of the utmost importance. Only if this is seriously and carefully undertaken can the mechanical removal of the gland be carried out with the minimum of risk.

Outline of Treatment

From the medical or preparatory point of view the treatment of thyrotoxic heart disease falls under four main heads: these are, rest, diet, iodine, and, in cases with auricular fibrillation, digitalis. Treatment by all of these methods should be so synchronized that the optimum result to be expected from each will be obtained

simultaneously. It is clear that relief to the patient will be at its maximum if treatment by each method can be so timed that its greatest effect will coincide with the greatest effect produced by each of the other three methods. The necessity for thus carefully planning the treatment will be clearly understood when it is remembered that treatment with iodine, if the usual full doses are given, leads to a marked improvement in patients with hyperthyroidism in about fourteen days; and that after this time, although the drug be continued in the same dosage, a partial relapse occurs. On the other hand, rest and diet may have to be persisted in several weeks longer before full benefit is seen. Again, digitalis takes a variable time to yield its maximum effect. Therefore, the best results can only be got if iodine and digitalis are administered in such a way as to secure the most satisfactory therapeutic response at the time when rest and diet shall have done their utmost in improving the patient's condition. When this has been accomplished the patient is ready for thyroidectomy.

Rest in Bed.—This must be complete. The patient should be isolated from all domestic, financial, or business worries. A quiet room must be chosen, and visitors should be strictly limited in numbers. A bedside telephone is not permitted. To assist mental rest sedative drug treatment is wise in severe cases; sodium luminal, 1/2 grain twice or three times a day during the stage of more severe nervous excitement, is probably the most efficacious. The services of a trained nurse are essential in all pre-operative cases. The patient should be kept strictly in bed during the whole course of medical treatment. The length of time which this requires averages about six weeks, but may vary between four weeks and three months.

Diet.—A high calorie value is essential, and the food should consist of normal solid meals. The calorie value of the daily diet should be between 3,000 and 3,500, and the patient should be encouraged to eat well. The meals should consist, as far as possible, of food which the patient is accustomed to, and should include in addition three pints of milk, taken as such, or in junket, cocoa, coffee, Horlick's, or ovaltine. Two ounces of cream and four to six ounces of lactose should also be added. The effect of the high calorie diet, together with the rest in bed, should be to enable the patient, first to make good the wastage caused by the increased basal metabolic rate, and, secondly, to put on weight.

Iodine Therapy.—Iodine should be prescribed with great care, particular notice being taken of its hardening effect on the thyroid gland. The maximal hardening of the gland can be obtained within a relatively short time (about fourteen days) if full doses of iodine are given, but when this point has been reached the drug appears to be without further avail, and the patient once more loses ground. Iodine, therefore, should be taken in smaller doses, which are enough to cause the gland to become fairly firm. It is kept in this state until the general condition of the patient has sufficiently improved to warrant operation. Three or four days before the date fixed upon for thyroidectomy full doses of iodine are administered. The dose at the beginning of treatment is 2 to 5 minims of Lugol's solution three times a day in milk. As soon as the gland starts to become

somewhat firm the dose is reduced to 1 to 2 minims twice or three times a day, enough being given in any particular case to keep the gland firm and elastic, but not hard. This dose is persisted with until three to five days before operation, when it is increased to 10 minims three times daily after food. In primary Graves's disease (primary toxic goitre) the true consistency of the gland is readily determinable. In secondary toxic goitre (toxic adenoma of the thyroid), where hard nodules exist already in the gland, palpatory information is not so easy to assess; in these cases attention should be directed to the feel of the gland away from the hard adenomatous areas.

Digitalis Therapy.—Digitalis is useful only in cases with auricular fibrillation. Employment of this drug in cases of thyrotoxic tachycardia with normal rhythm is a waste of time. This statement applies to all preparations of digitalis and strophanthus, both pharmacopoeial and proprietary. In the presence of auricular fibrillation digitalis may be used. The best results are obtained from it when oedema of the feet and other signs of severe congestive failure are present. In cases not thus complicated it will probably be found that rest, diet, and iodine, as described above, will slow the heart rate more than will digitalis. Patients with normal rhythm need no digitalis. Patients with fibrillation of the auricles but with no signs of failure should be given 40 minims of tinct. digitalis or 4 grains of digitalis folia (B.D.H.) a day for three days before operation. Patients with auricular fibrillation and signs of congestive failure should receive 40 to 60 minims of tinct. digitalis, or 4 to 6 grains of digitalis folia a day for four to six days at the commencement of the period of medical treatment. If symptoms of anorexia or nausea appear, the dose is reduced to 30 minims or 3 grains a day; in the absence of such symptoms of overdose the drug is continued in the larger doses as long as signs of failure persist. When the degree of heart failure is especially severe the following measures may be undertaken. Digitalis should be pushed if auricular fibrillation is present. When severe and protracted attacks of vomiting complicate the heart failure the drug may be administered per rectum in the form of the tincture. In previously untreated cases and as an emergency measure 3 drachms in 4 ounces of water may be injected per rectum, three such doses being given at four-hourly intervals. A maintenance dose would be 1 drachm in water twice or three times a day.

Thyroidectomy.—The medical treatment outlined above will lead in most cases to a diminution in the signs of hyperthyroidism, an increase in weight, and a fall in the pulse rate. So long as improvement, as judged by these criteria, continues medical treatment should be persisted in. When the improvement becomes considerably less rapid, or when it appears that further improvement is not to be obtained by these measures, surgery should be considered, and is in most cases necessary. Thyroidectomy is contraindicated, as a rule, if weight is being continually lost, if the heart rate is above 120, and if there are signs of gross failure. Exceptions to these rules are met with, but judgement as to the safety of operation in them is a skilled matter, and the risk is increased. Should severe symptoms persist ligation of the thyroid arteries should be carried out preliminary to thyroidectomy.

Post-operative Failure

Post-operative acute failure is to be treated by the following measures.

1. Continuous oxygen administration, by oxygen tent or nasal catheter.
2. Continuous administration, per rectum or intravenously, of saline containing 5 per cent. glucose.

3. Sugar should also be taken by mouth, and insulin in the proportion of one unit for every 2 grams of sugar ingested should be injected intramuscularly.

4. Sodium iodide, 20 to 40 grains in 10 c.cm. of sterile water, should be given intravenously twice daily.

5. Lugol's iodine should be prescribed in milk in doses of 30 minims four-hourly, or per rectum in doses of 1 drachm four-hourly.

Treatment of Auricular Fibrillation

A high percentage of those who before operation had auricular fibrillation lose the irregularity spontaneously after thyroidectomy. Should this not occur within three or four weeks it is desirable to restore the normal rhythm by means of quinidine sulphate. This drug is avoided by many practitioners because its use has occasionally been followed by a fatal accident. The same result follows many simple surgical operations and in about the same percentage of cases; the risk involved should be made clear to the patient or the relatives. The second point to be remembered is that quinidine produces a definitely depressant action on nearly all cardiac functions; therefore a patient who is being treated with it must be nursed flat, as though suffering from a myocarditis; he must be fed and washed, must not sit up, and, *a fortiori*, must never be allowed to get up out of bed. Thirdly, it must be remembered that the drug is excreted very rapidly, and that doses should be given two-hourly if a reasonably constant quantity is to remain in solution in the blood stream. Better results are obtained if this is done. My usual routine is as follows:

1. Digitalis is prescribed in reasonably full doses for fourteen days, the amount being 45 minims of the tincture a day, or 4 grains of the folia. Previous digitalization improves the therapeutic effect of quinidine.

2. For the first day of quinidine treatment 1 grain of quinidine sulphate is taken three times a day to eliminate the possibility of the patient's having a quinine idiosyncrasy.

3. For the second day quinidine sulphate is given in 2-grain doses two-hourly for eight doses.

4. For the third day the dose is 3 grains two-hourly, and for the fourth and fifth days 4 grains and 5 grains respectively are administered two-hourly for eight doses. The drug is prescribed thus:

R Quinidin. sulph. gr. ij
Acid. sulph. dil. m xv
Aq. chlorof. ad 3 ss

(The sulphuric acid dissolves the otherwise somewhat insoluble drug.)

If 30 to 40 grains of quinidine have not produced the desired effect, it is wiser to desist, unless the physician has seen a sufficient number of cases to feel confident that a further increase in dose in the particular case is a safe measure. Signs indicating that a cessation of quinidine treatment is desirable are buzzing in the ears and deafness; feelings of faintness or dizziness; any considerable rise in the ventricular rate.

No case of thyrotoxic heart disease with auricular fibrillation should be under the care of a nurse who cannot accurately count the heart rate at the apex with a stethoscope. If the heart rates recorded on the chart cannot be relied upon by the physician, he is not in a position to estimate with full accuracy the clinical position. This is equally true whether he is treating the auricular fibrillation with digitalis or with quinidine.

Prognosis

After thyroidectomy the cardiac insufficiency disappears in most cases in a remarkable manner. It is my experience that the prognosis depends upon whether any disease additional to the thyrotoxicosis, such as rheumatism or arteriosclerosis, has affected the heart. In a careful follow-up at St. Bartholomew's Hospital,¹ it was found

¹ Bach, F., and Bourne, G.: *Quart. Journ. Med.*, 1932, xxv, 579.

that the cases fell into two groups—namely, those who still had evidence of heart disease, although the hyperthyroidism had been cured by thyroidectomy; and those with normal hearts. In the former group nearly every case had evidence of rheumatic or arteriosclerotic heart disease. In the latter group no evidence of heart disease was discovered. It would seem that the permanence of any cardiac incapacity after hyperthyroidism and thyroidectomy depends, more than on any other factor, on whether the patient's heart has been previously diseased. Thyrotoxicosis may accentuate and may perpetuate the effects of rheumatic or degenerative cardiac disease, but it leaves a previously healthy heart in a normal condition.

Another abnormality which tends to follow thyrotoxicosis is hyperpiesis. This is not present in all patients who have suffered and recovered from Graves's disease, but the incidence is certainly higher than normal in these cases. The condition may result either directly from the thyrotoxaemia or from the wear and tear produced by the long period of increased basal metabolic rate. It is my opinion that the latter is the more probable hypothesis, and that the thyrotoxicosis accelerates the onset of a condition which would in any event have occurred later on.

Noëba et Vetera

HISTORY OF CANCER OF THE BREAST

Sir D'Arcy Power tells us that when he was asked to deliver the sixteenth William Mitchell Banks Memorial Lecture at the University of Liverpool,¹ he had little hesitation in choosing a subject, for, although Banks was interested in most surgical topics, his work on the operative treatment of cancer of the female breast stands forth pre-eminent among all his surgical studies.

Sir D'Arcy Power traced the history of cancer of the mamma and its treatment from the earliest records of surgical procedure. In this, as in so much else, we first find something approaching firm ground from which to make a survey in the works of Galen. To the ancient and mediaeval surgeon the term "cancer" was clearly associated with the existence of an open and spreading sore, and malignant disease affecting the female breast, but not yet involving the skin, was scarcely recognized as cancer. For many years attempts to cure by means of caustics were the only means in vogue with reputable practitioners and quacks. When, after horrible and prolonged suffering, the superficial portions of a cancerous mass exfoliated, hopes of cure were entertained, to be all too soon followed by recurrence and, if the patient was fortunate, by speedy metastasis and death. Jean Louis Petit, director of the French Academy of Surgery and a Foreign Fellow of our Royal Society, who lived from 1674 to 1750, made a great advance in pathology and treatment, although his views were not published till twenty years after his death. To him Sir D'Arcy Power ascribes the credit of recognizing and acting on the principle that the "roots of a cancer were the enlarged lymphatic glands: that the glands should be looked for and removed and that the pectoral fascia and even some fibres of the muscle itself should be dissected away rather than leave any doubtful tissue. The mammary gland, too," he said, "should not be cut into during the operation."

It is tantalizing to read of such great strides forward as those of Petit and yet to find that he was not equally enlightened as to the true line of advance in some important respects. Some of his principles were to be entertained as new truths a century later by his spiritual heirs, while he was unfortunately so mistaken as to try and preserve the nipple and to avoid excision of skin. Petit's teaching was known outside France before his death and recommended in an English translation of Lorenz Heister's "Surgery."

¹ *Liverpool Medico-Chirurgical Journal*, 1934, xlii, 29.

The teaching of Benjamin Bell of Edinburgh showed a considerable advance in theory and practice. He taught that "when practitioners have an opportunity of removing a cancerous breast early, always embrace it." His teaching in other respects was advanced as to the treatment of glands and the parts adjacent, and he had a great and lasting influence upon British practice for many years.

If it may be conceded that the radical operation of amputation of the cancerous breast as at present practised is that devised by W. S. Halsted of Johns Hopkins University at Baltimore, U.S.A., it is equally necessary to give credit to those American surgeons who paved the way for the advance. Of these Pancoast and Samuel W. Gross, both of Philadelphia, were his precursors. American surgery, looking more to Germany than to the rest of Europe for surgical inspiration, had based itself on scientific pathology, and it was with the aid of such foundations and his own investigations and deductions that Halsted based his proceedings, culminating in an operation which was intended to remove all infected tissue with a wide margin. American surgeons were ready and eager to accept Halsted's teaching, while in Great Britain the response to new ideas was nugatory or at best slow.

In 1867 C. H. Moore of the Middlesex Hospital contributed a remarkable paper to the Royal Medical and Chirurgical Society of London. In this paper, as Sir D'Arcy Power says, is the germ of all the modern surgery in connexion with cancer of the breast, and it is noteworthy that among the cases cited in support of his thesis Moore describes the putting of a drainage tube through the back of the armpit—and this was in the year 1858. No report of this paper appeared in the *Lancet* or the *British Medical Journal*, but a short summary of the subsequent discussion was given in the *Medical Times and Gazette*. Its reception was distinctly unfavourable. None of the speakers can have realized the immense importance of Moore's contribution. He does not seem to have pressed the subject on his colleagues' attention.

The epoch-making paper of W. Mitchell Banks was read by him before the Lancashire and Cheshire Branch of the British Medical Association in 1877. It was entitled "A Plea for the More Free Removal of Cancerous Growths." It was reprinted from the *Liverpool and Manchester Surgical Reports*, which had a limited circulation locally; but no notice of it appeared in the *British Medical Journal*, the *Lancet*, or the *Medical Times and Gazette*. Banks continued his attack on the subject in 1882 and 1883, basing his work almost exclusively on clinical observations, and not, like Halsted, on micro-pathology. In 1887 he read a paper in London before the Harveian Society on "Extirpation of the Axillary Glands, a Necessary Accompaniment of Removal of the Breast for Cancer." Halsted's first communication on the subject was made in the *Johns Hopkins Hospital Reports* for 1895. His work was original, and he evidently knew nothing of the work of Moore and Mitchell Banks.

Sir D'Arcy Power's paper is meant not to detract from Halsted's fame but to show that others, by different methods, had arrived before him at similar results.

The annual general meeting of the Union Thérapeutique will be held on Wednesday, October 9th, at the Faculté de Médecine de Paris. The subjects of discussion are: "Nos Connaissances Actuelles sur l'Hormone Mâle," by Professor A. Guy Laroche; "La Chirurgie du Splanchnique dans l'Hypertension Artérielle," by Professor Ag. René Fontaine (Strasbourg); "Les Composés Antimoniaux en Thérapeutique," by M. J. la Barre (Brussels). In the afternoon: "L'Équilibre Physico-chimique et Glandulaire dans le Traitement des Spasmes Viscéraux," by Professor Pende (Genes); "Les Substances Spasmodiques et leurs Antagonistes," by Professor Burgi (Berne); "La Médication Nerveuse Sédatrice des Spasmes," by Dr. J. Decourt; "La Physiothérapie des Spasmes et des Algies Viscérales," by Dr. Delherm. Anyone wishing to attend the meeting should write direct to Dr. G. Leven, 24, Rue Téhéran, Paris (8^e).