TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some of the major medical disorders met with in general practice.

FUNCTIONAL DISORDERS OF THE HEART

BY

A. G. GIBSON, M.D., F.R.C.P.

Functional disorders of the heart are seen in those who are organically sound as well as in those with disease of the heart or vessels. In the latter case, however, the functional element is so overshadowed by the organic as to be of minor practical interest. The remarks that follow are mainly concerned with hearts that are organically sound.

The symptoms of functional disease bulk considerably in ordinary practice, and unless treated upon right lines may result in a condition of chronic ill-health. It is presumed that every patient who is suspected of having any functional ailment will be examined by all the usual clinical methods. It is especially necessary to be clear as to the presence or absence of organic murmurs, enlargement of the heart, the systolic and diastolic blood pressure, and the cardiac reserve as defined by the patient's reaction to ordinary exertions in the course of his life and to simple tests which may be done during an examination.

Palpitation

The normal heart in a normal body gives no sensation or awareness of beating, and it is very rare to hear a complaint from a patient with gross heart disease of discomfort from that source, even though the whole chest is shaken at each beat. But patients frequently complain of this awareness of the beat, and an inability to sleep on the left side owing to this sensation keeping them awake. Examination of the cardiovascular system may sometimes reveal a raised blood pressure from early arterial change, sometimes also the signs of hyperthyroidism. But in other patients the cardiovascular system to clinical examination is normal and the exercise tolerance is not diminished. The cause may be that the patient has recently passed through an acute infection and has resumed work too soon. In other patients it is necessary to search for the effects of a toxin such as alcohol or for a focus of infection, or to make inquiry as to whether the patient has recently been subject to any mental or physical strain. The treatment in any case should be directed towards eliminating undue stresses or toxins, by correcting any faulty habits, and by giving bromides (pot. brom. 15 grains thrice daily). Treatment by cardiac tonics must be avoided, in order that the patient's idea of cardiac disease, which he may secretly cherish, may be removed; also for the reason that cardiac tonics such as digitalis do no good.

Extrasystoles

Extrasystoles may occur without demonstrable cardiac disease or as the result of myocardial affections. As purely functional features they disappear when the cardiac rate is raised by exercise, and they are not accompanied by pain, embarrassment, or breathlessness, through exertion. They are sometimes extremely uncomfortable to the patient, and unless the condition is explained to him as being of no consequence they may repeatedly remind him of a heart of which he was previously

unaware. The source of such extrasystoles is frequently extracardial—a septic focus such as enlarged tonsils, and the result of irritation of the gastric mucosa, as, for instance, in chronic alcoholism, in which the liver is beginning to be enlarged. Occasionally they may give warning of more serious myocardial conditions in middle-aged people, such as the approach of disease of the coronary arteries. Under these circumstances a more careful examination of the heart should be made, including an electrocardiogram.

The main element in treatment when there is no sign of incipient cardiac disease or insufficiency is to assure the patient that the symptoms are not serious, to make a complete overhaul of the other systems, and to inquire into the daily life of the patient so that the necessary corrections may be applied. Sir James Mackenzie used to recommend as the best treatment 20 grains of ammonium bromide three times a day for two days, to be repeated again after an interval.

Simple Tachycardia

The treatment of a tachycardia which may be producing discomfort or symptoms depends on having a clear notion of its cause, for it is seldom amenable to efficient treatment by drugs which act on the heart. Tachycardia which depends on a recent attack of some infective disease such as pneumonia, diphtheria, or influenza is best treated by a prolonged convalescence and rest. Acute rheumatism which has caused a tachycardia would seldom be without some other feature either in the history or state which would give the clue. An anaemia would have its appropriate remedy according to its type, though even severe anaemias do not always quicken the pulse at rest.

There are three common types of tachycardia for which the remedies are clear.

1. The first is that dependent upon abnormality of thyroid action, whether as in exophthalmic goitre or from what is commonly called toxic goitre. In the latter type there may be few of the ordinary signs of thyroïdal disease, and yet the excision of the larger part of the thyroid may bring great amelioration in the condition of the patient. It occasionally happens that a patient may secretly be taking thyroid for "slimming" or some other purpose.

2. Chronic infection of the throat from a haemolytic streptococcus sometimes gives rise to a persistent tachycardia which may be very difficult to distinguish from early hyperthyroidism. The throat may show no more than a granular pharyngitis, such as may be seen in many persons during the winter months. The treatment should be local and general. Astringent preparations such as glycerin of tannin or glycerin of alum painted on the throat two or three times daily tend to heal at least those parts of the throat that can be reached by a brush; vaccine therapy is also of value, but if an autogenous vaccine is used the doses should be minute at first—for example, one million streptococci or less—because these patients may be unduly sensitive, and the reaction, both local and general, may be severe.

3. In some patients coming with a trivial complaint the only physical sign may be tachycardia; in a case of this nature it is well to suspect some emotional cause.

such as an anxiety neurosis. The rate of the heart in this type of case tends to become less as the confidence of the patient is gained, because many of his fears are not necessarily deep-seated. The fear of cancer or of pregnancy in women are common examples. If these patients are put to some simple exercise test the pulse is frequently the same or less rapid after the test; it thus becomes clear that the rapid pulse is not an indication of diminution of cardiac reserve. In such a patient it is best to explain that no disease has been found, and with tact to inquire if there has been any psychological difficulties. When the confidence of the patient has been gained it is frequently possible without offence to ask direct questions as to possible causes of anxiety. Drug treatment is best avoided altogether.

Precordial Pain

A pain on the left side of the chest may have its origin in a pleurisy, intercostal neuritis, or some other non-cardiac ailment. Apart from disease of neighbouring organs, pain in the region of the heart is the unmasking by some abnormal stress of a normal reflex mechanism. The pain of "stitch" that is complained of by school-boys and untrained athletes at the limits of their tolerance sometimes has the distribution of the pain in angina pectoris. Pains that occur in young persons and in women in this region are less likely to be cardiac or serious than pain which occurs in middle-aged men, and the first thing to be satisfied about is whether the heart is normal to clinical tests and has a normal exercise tolerance. When necessary an electrocardiogram should be taken, because it might reveal something hitherto unsuspected.

The treatment of these precordial pains, as of the other functional disturbances that have been considered, depends on relieving the patient of some stress that has been causing the pain. Rarely a careful history may reveal some toxic cause. Alcoholic patients frequently refer their abnormal sensations to the heart. Excessive smoking should be inquired into, and though we cannot often be certain that tobacco is the cause of pain in the heart an obvious excess should be cut down. Most patients resent a suggestion that they should give up tobacco completely, but are ready to accept the advice that they should cut it down by half.

Among the reflex causes of precordial pain is an irritation from an inflamed gall-bladder or a duodenal ulcer. Abnormalities of the oesophagus, such as a chronic oesophagitis, may also simulate a cardiac pain. Abnormalities of this nature should be corrected. An anxiety neurosis is also a frequent cause of precordial pain, and the treatment should include a strongly worded assurance of the absence of disease. Associated with the anxiety neurosis in women there may be a mild degree of hypochromic anaemia, and the anxiety may be based on an inability, through the anaemia, to do what is demanded by the exigencies of normal life. The treatment of the anaemia is to give *ferri et ammon. cit.* 30 grains thrice daily in a mixture. All nervous stress and fatigue may accentuate pain of this nature. Drug treatment of the heart would be a confession of failure, and the method is to allay the excitability of the cardiac reflex previously referred to by getting the patient to pay less attention to the heart, and by giving bromides combined or not, as thought necessary, with hypnotics.

It is best in dealing with these patients to avoid the use of the words angina or anginal pain, which are now employed for clearly defined cardiac disease. The term "precordial pain" has no such ominous significance. It merely states the position in which the patient feels discomfort and does not necessarily imply the existence of organic disease.

The Effort Syndrome

The effort syndrome is now rare, and though more common in men is not confined to them. Repeated clinical examinations are necessary to exclude organic heart disease, hyperthyroidism, and chronic infections, especially from faulty teeth and an anxiety neurosis. It is wise to have an electrocardiogram done in case of subsequent developments, for though with lapse of time and general treatment with this patients tend to recover, it may be the beginning of slow coronary change, which may ultimately show itself in sudden death. The patients must be put to a healthy life with a regular amount of exercise, which should be taken at their own pace, short of great pain or fatigue. The most useful drug is bromide.

The Athlete's Heart

Athletic strain is a potential source of temporary cardiac disability; but in well-trained persons with a normal physique it would appear to be excessively rare. Minor changes, however, are not infrequently seen in those who engage in athletics without proper training or with a recent history of some mild infection. Nevertheless, young men engage in competitive athletics a week or so after a febrile attack without ill effect.

The changes that are most likely to be met with are a pulse that is more rapid than normal, a slightly misplaced apex beat, a systolic murmur at the apex, and very rarely a diastolic at the base. When any such abnormality is found it is best to restrict exercise to walking. It is seldom necessary to put such a patient to bed unless there is a suspicion of rheumatic endocarditis. Temporary disturbances of this kind, if due mainly to strain, settle down in a week or two, and subsequent athletics can be undertaken gradually with proper training, preferably under supervision. It is necessary to be sure that what may be suspected to be a cardiac murmur is not a cardio-respiratory murmur, that the misplaced apex beat is really in the fifth and not in the fourth space—where it might have no significance—and that the rapid pulse is not due to some psychological cause. If there is no adequate reason for the cardiac condition the patient should be examined for a cryptic infection, and inquiry should be made into his habits, especially in respect of alcohol.

O. Lassen (*Hospitalstidende*, May 21st, 1935) believes, after many years' experience in the tuberculosis dispensary of a large Danish town, that it is through this institution that further progress is to be sought in the reduction of the tuberculosis mortality. But if the dispensary is to serve its purpose, its staff must go to the patient and his contacts, and not wait to be visited by them. The ideal is the visiting and examination of every person in the household of a new case. In 1933, in the town in question, fifty-four new open cases of pulmonary tuberculosis were notified. They belonged to fifty-two households with a total membership of 152 apart from the patients. As many as 124 of these 152 contacts were examined, and in thirty households all the contacts were examined. The author believes that it should soon be possible for such home medical examinations to be accepted by every member of the household as a matter of course; at present it is often the peccant source of infection who refuses to be examined. Between 1920 and 1933 the deaths from pulmonary tuberculosis per 10,000 inhabitants in this town (Aarhus) have fallen from between eight and nine to between three and four. In the same period the percentage of tuberculosis deaths occurring in hospital or sanatorium has risen from between thirty and forty to between eighty and ninety. This growing tendency for patients to die in an institution instead of at home is most desirable in the opinion of the author, who sees in the isolation of infectious cases one of the best guarantees for stamping out tuberculosis. Home conditions are often unknown to the medical authorities.