BRITISH MEDICAL JOURNAL

LONDON: SATURDAY, JULY 13th, 1935

THE INJECTION TREATMENT OF VARICOSE VEINS

Some Late Results and the Question of Recurrence

BY

G. H. COLT, M.B., B.Ch.CANTAB., F.R.C.S.

ISOBEL S. W. RAMSAY, M.B., CH.B.ABERD.

AND

MARGARET M. M. MORRISON, M.B., CH.B.ABERD.

(With Special Plate)

The present position of the injection treatment of varicose veins should properly be determined by the late results. Sufficient time has not elapsed in this country for the really late results to be available in large numbers, and, as the variables in assessing the results are numerous, it is difficult to come to definite conclusions. This difficulty is so great that some observers have returned to ligature treatment in numerous cases. Most of these cases appear to us to be curable without ligation if the injections are given at longer intervals. Such factors as the tendency to the disease, the difficulty of classification, and the variety of solutions employed all enter into the problem.

Types of Cases

It is well known that in any clinic in which the cases are followed carefully for some years a residual number become chronics. This type (A) presents a multiple spongework of relatively small varicosities, and although the patients have been greatly improved symptomatically by injection treatment alone, numerous varicosities, having a large total blood volume, remain. In some cases this type presents in the late stages one main trunk, which can be either injected or ligated, but in most instances this fortunate termination does not occur. It would appear that this is the one variety in which, could it be diagnosed early, a preliminary ligation of the internal or external saphena, or both, should be practised and in some cases combined with injection. As a guide we have the good results of Babcock's extraction operation, combined with local excisions, in the case of soldiers on active service. Followed over a period of five years the cures are in the region of 80 per cent.¹ Jeannel² gives 60 per cent. for multiple resections and 95 per cent. for total saphenectomy, the latter in seventy-seven limbs.

The other chief types which present themselves are: (B) The single varicosity, generally with a positive Trendelenburg sign. In this almost any variety of sclerosing solution injected into the empty vein will give a good and lasting result. It is unnecessary to ligate the vein.

(C) The more extensive type, with positive Trendelenburg sign and perhaps also evidence of a greater or less degree of deep reflux, may be treated most effectively by using the more caustic varieties of solutions. In these types (B) and (C) the degree of varicosity and the number of injections of salicylate (40 per cent.)-saline (10 per cent.) that were required have been given before³ as:

Very severe in	5 limbs, requiring a total of	24 injections—Av.	4.8
Severe	31	152	4.9
Marked	98	284	2.9
Medium	82	174	2.1
Slight	27	27	1.0
	243	661	

or an average of 2.7 injections per limb in 150 patients. This finding has held in cases done later. What has been named the "stiction" effect, in which the walls of the vein collapse and stick together almost at once after the injection, and so remain, occurred in fifteen of these cases. Continued observation indicates that this very happy state of things remains permanent for six years, and is presumably a cure of the particular veins involved. It is to be noted in this connexion that the best and most lasting results obtained by Linser and Zirn⁴ and by Fischer⁵ were procured by the use of mercuric chloride before this solution was abandoned on account of its danger to the general health.

William M. Cooper⁶ based a study on 3,164 cases requiring 35,000 injections of various solutions, and 239 cases of extensive and recurrent varicose veins treated by preliminary ambulatory ligation and subsequent injection, a proportion of about one to thirteen. He finds that the majority may be treated successfully by injection alone. No definite figures are given for cases treated by any single method, neither is the number of limbs stated. If each case represents one limb the average number of injections per limb was eleven, or more than four times those required in our series. He points out that some patients cannot be treated successfully in this way without preliminary ligation, which in a valveless vein protects the thrombus from recanalizing. He considers that the first thing to do is to classify the patient by means of the Trendelenburg test. Now this test, which was described by Brodie⁷ fifty years before Trendelenburg, has, when positive, a perfectly definite meaning, and used to have an equally definite one when it was negative. It seems to us that some writers confuse the issue by adding to the sign when it is negative the positive evidence of a reflux of blood from the deep communicating veins. This has nothing to do with Trendelenburg's sign as formerly understood. It may occur whether the sign is positive or negative. These further additions to the nomenclature are called "Trendelenburg double" and "Trendelenburg nil." When double the veins fill both by superior and by inferior reflux, and when nil they do not fill at all. When the test is positive Cooper is in favour of high ligation of the internal saphena, but, as we can show (Tables II and III), there are many cases in which injection alone answers equally well and the patient is saved an operation. Our own figures for ligation are three limbs out of 952. When the reverse flow is through the incompetent communicating veins Cooper would proceed directly with injection, but if the veins are large a preliminary ligation is performed at the highest point at which a reverse flow can be demonstrated. Our experience has frequently been that all such

veins can be obliterated by injections alone, except at the actual points where the perforating veins emerge and join the superficial circulation. Turner Warwick⁸ considers that ligation below the deep fascia is safer than injection for the remaining cul-de-sac of a perforating vein. "In the past three years I have given up preliminary ligation, and in all cases have begun with injection. In very few cases has any resort to operation been necessary. . . ." Our experience strongly confirms this view. In the Trendelenburg double cases Cooper performs multiple ligations at all the demonstrable points of back flow. As already mentioned, a five-year interval shows that such a plan, when combined with total saphenectomy, is followed by partial failure. Our experience of injections in old operation cases shows that the results are excellent, presumably because the flow of blood in the varices injected is sluggish. This, however, is not an argument in favour of preliminary ligation, for equally good, if not better, results can be obtained by primary injection. Cooper would limit injection without ligation to patients who present either large or small varices with no evidence of *marked reverse flow*.

Patients, then, who presented very large and extensive varices with marked reverse flow, as well as those with recurrent veins, were selected by him for preliminary ambulatory ligation. It must be admitted that such a course is likely to yield good results, but only the future can determine whether it is better than injection without ligation and whether it is devoid of risk.

Risk of Ambulatory Ligation

The actual risk of ambulatory ligation appears to be a small one if the danger spot near the junction of the saphena and femoral is avoided. Moszkowicz⁹ states that ligation here is so dangerous that it should be abandoned, and that the curative results are unreliable. M. Hanselman¹⁰ states:

"... the longer the proximal stump the more apt is the proximal clot to appear . . . ligation close to the sapheno-femoral junction is the best measure to avoid large clots in the proximal stump. Thus, aside from the ambulatory operation, the shortness of the stump is another safeguard against massive embolism."

This statement is true theoretically, but practically the injury to the intima at the ligature, especially when the vein is severed, as it usually is, must produce a tangential thrombus. One of us (G. H. C.) can recall the case of a woman of 37 who was operated on for a left femoral hernia in 1912. On the fourteenth day of recumbency the right internal saphena was ligated for a varicose cyst in its upper part close to the sapheno-femoral junction. Fatal embolism occurred on the tenth day. The wound was aseptic. There was no necropsy. If the ligature is placed below the lowest competent valve, as has been done by de Takáts¹¹ to ensure a better protection against the superior reflux, and so prevent proximal dilatation, the thrombus caused by the injury to the intima will be shut off by this valve from the superior circulation. No embolism should occur if the valve remains closed until sufficient organization has taken place. One of his patients showed a sacular dilatation with impulse between the sapheno-femoral junction and the site of ligation. The piece of vein was removed up to the junction. "The final analysis of these recurrences indicates that high ligations should be done in every case. . . . Ambulatory management and shortness of the proximal stump are important in the prevention of embolism."

The explanation of such diverse views on the risk of ambulatory ligation appears to us to lie in the flow of blood as found in the upper end of the saphena. When the ligature is placed flush with the femoral, so that no pouch is left, the thrombus at the division of the intima

is femoral and tangential. It is washed by a rapid stream of blood. But when the ligature is placed lower down the saphena the thrombus lies secure in a tube of vein, nourished by a slowly moving stream of blood contributed by the superficial epigastric, circumflex iliac, and external pudic veins. These conditions are admirably suited to the production of a firmly adherent and organized thrombus which is unlikely to be detached. Proof of this explanation will no doubt be obtained as experience is gained and records are published stating the length of the proximal stump of vein and giving the number of tributaries it receives. In the Babcock extraction series, and in many cases done since then, the saphena was divided well below the entrance of the veins referred to. No embolism occurred, either during recumbency or later. In that series 103 ligations were performed and some seventy-eight yards of vein extracted. Ambulatory treatment alone, therefore, cannot be the factor which renders ligation of the saphena safe, though it has often been given the credit for doing so.

Recurrence of Varicosity

We come to the problem of so-called recurrence. It is clear that there are three chief factors involved in this. (1) The natural course of the affection in its various manifestations; (2) the degree of completeness of the thrombosis in the particular venous area under observation; and (3) the varicosities observed in the injected and adjacent areas. The following extracts are not meant to be regarded as a complete summary of the published observations of others, and are only given as bearing on our own observations.

Linser and Zirn¹² noticed many cases of complete obliteration of the veins extending over several years, but collateral varicosity appeared some three to four years after the original perchloride of mercury injections. In Fischer's¹³ patients the veins were impervious cords three to five years after the use of perchloride. Bazelis,¹⁴ using salicylate, had no recurrence of veins or of ulcers after three years' trial, but this seems to be too short a period, as our cases show. Hemple¹⁵ found collateral varicose development in 12 per cent. of his cases. Howard, Jackson, and Mahon,¹⁶ in recording recurrence of varicosis and figuring the process of recanalization, say: "It is of interest that in patients who had competent valves of the perforating set, but incompetent valves of the saphenous vein, recurrence developed with as great a regularity and within the same space of time as in those who had incompetent valves in both perforating and saphenous veins." This would point to the superior reflux having been much the more important factor. Of sixty-five cases fifty-two showed recurrence in a year or longer. The injections were made with 20 per cent. NaCl at intervals of one to two days, and then at weekly intervals, apparent complete thrombosis being obtained. The intervals between injections seem to us to be too short to allow of accurate observation of the blood paths in a process which takes at least seven weeks to harden up after its initiation. One case is mentioned with massive complete thrombosis on both sides, which showed recurrence at eight months and was reinjected. Recurrence, which was presumably recanalization, again took place. Wm. M. Cooper¹⁷ a year ago found that the deep reflux was chiefly responsible for a 15 per cent. recurrence in cases of extensive varicosity injected with various solutions. Preliminary ligation of the saphena reduced the number of injections required to that of one-third to one-fifth, and symptomatic relief was immediate in nearly all cases. Four per cent. showed complete occlusion after ligation and required no injection. Patey and Tatham¹⁸ found recanalization to be a frequent event in fifty cases called for examination two or more years after quinine injection. Forestier¹⁹ says: "In 15 per cent. of the cases recurrences are noticed, not on the obliterated varices, but on varicose sections appearing after the injection. As we know that the varicose condition is a diathesis, we are not surprised by such recurrences. In such cases a few supplementary injections after one or two years are likely again to bring out favourable results."

The subject is difficult to investigate accurately, especially in an out-patient clinic, where the patients are not always seen by the same observer. It was our advantage to be able to use part of a surgical ward. Apart from a good descriptive note and a diagram a clear mental picture of the condition has to be carried over a period of years. Infra-red photographs²⁰ are a help, but scarcely practicable as a routine, on the grounds of time, labour, and cost. They show up small varicosities and collaterals which can neither be seen nor felt, and which may become varicose later. An intelligent patient, anxious to help in the cure of his trouble, will give valuable information about the appearance of new varicosities, the reappearance of older ones, and the question of recanalization. One patient said he could feel blood trickling past a stoppage (thrombosis) in a vein at his knee and, as he thought, opening up the vein again. Five weeks later this vein looked and felt as if it had never been injected. All the patients have been under the continuous observation of one of us from June, 1927, to June, 1934.

The classified facts are shown in Table I. The numbers scarcely warrant a percentage basis, but in round figures 29 per cent. of limbs showed no new varicosities, 19 per cent. did show new varicosities, and 40 per cent. showed either incomplete thrombosis or recanalization. In the majority of the latter the matter was a small one, and easily rectified by a further one or two injections. The result of ten cases out of twenty-six of diffuse varicosity showing no new veins is a good test of the method employed; the remaining sixteen were all of them greatly improved. Turner Warwick²¹ says of the diffuse type: "When the varicosis involves not only the main superficial veins and chief perforating channels but also innumerable small superficial and undescribed perforating veins, complete success cannot be ensured. Relapse is frequent, and repeated courses of injection are often necessary. In some of these extensive cases the deep veins are varicose, and then the permanent cure of the superficial varices is still more doubtful."

TABLE I.—*Recanalization and New Varicosities*

Facts Noted	Enormous	Large	Marked	Medium	Small	Diffuse	Total
No new varicosity	2	2	3	16	13	10	46
Perforating varicosity present	1	1	3	4	2	0	11
Incomplete thrombosis	0	18	17	6	5	6	52
Recanalization ...	1	6	2	3	0	1	13
New varicosities	1	4	5	6	6	7	29
Uncertain...	0	2	0	5	0	2	9
Total limbs ...	5	33	30	40	26	26	160

All the limbs in Table I were examined three years after the last injection.

Methods Employed

With the occasional exception of the trial of a different solution in well-defined cases the salicylate (40 per cent.—sometimes 30 per cent.)—saline (10 per cent.) solution of V. Meisen²² has been used throughout. It is the most reliable of all the non-toxic, guidable²³ solutions and rarely fails, though sometimes the reaction is excessive if the blood stream is sluggish and the dose has not been adjusted accordingly. The dose has varied from 12 c.cm. of 40/10 to 1 c.cm. of 30/10 according to the blood volume of the vein and the speed of blood flow. In general, the injections have been given into the empty vein from below upwards, and have been guided into

the tributary areas as required. Injections in the standing or sitting position have seldom been employed. When so used the diluting blood volume has been small. The injections have usually been given alternately to each leg at monthly intervals, sometimes fortnightly, when the varicose areas have not been closely adjacent. Such a system allows the patient one leg to walk on should the reaction happen to be too severe, and enables the surgeon to exercise the optimum time of a month for observing the thromboses and their effects in diverting the circulation. Even one small injection may cause a wide alteration in blood flow, and it is only in the most extensive cases, or in widely separated varices, that more frequent injections are called for. When we hear of patients with enormous varices, with the breathing reflux at the knee visible from afar, being injected with mild sclerosing solutions given in the standing or sitting positions, and repeated two or three times a week, it is evident that there is a lack of appreciation of the principles which should guide our efforts. Such procedures may perhaps account for some of the enormous differences in the published results of injection treatment; but as no satisfactory way has been found of classifying the cases so as to be able to obtain a comparison in any two lengths of vein, apparently similar, in patients who are in good general health, it is difficult to be certain. The mere size of the vein does not affect the cure. In one case a superficial anastomotic vein between the external and internal saphenous veins was noticed to have a specially rapid current of blood, but it was readily obliterated. Great tortuosity tends to help thrombosis by allowing the solutions longer time to act. In such cases the reaction is sometimes excessive. Veins containing phleboliths, as evidence of stasis in the bays, react well. Previous phlebitis, which has cleared up, undoubtedly aids the induction of sclerosis. One patient showed ordinary varicose phlebitis in her left thigh and leg and an apparently comparable arrangement of non-phlebitic varicosities on the right side. The right side was injected with salicylate saline. Six months later nature's result was just a little better than man's aided by nature, but both were good and persisted for three years.

Lightly applied thin elastic-webbing bandages have been worn night and day throughout the course to diminish the blood flow and to prevent excessively large clots from forming. Theis²⁴ condemns bandaging and reclining as being causes of red instead of white thrombus formation, but our experience is that the results are good and that pain is relieved.

Necrosis

A useful technical observation must be added here. The recognized causes of necrosis are: (1) injection outside the vein, (2) the bursting of a tributary junction, and (3) injection into the substance of the vein wall. All these causes having been largely remedied by practice and care, it was puzzling to find that necrosis still occurred, though rarely. The explanation seems to lie in the loss of elasticity in the wall of the vein following a previous injection, so that it does not seal itself and the puncture leaks. By waiting longer than usual before withdrawing the needle the solution will become more dilute and innocuous.

Results of Treatment

The classified results of unselected cases seen one and three years after injection are given in Tables II, III, and IV, and have been marked down rather than up. It is interesting to note that the largest number of poor results occur in the "marked" size of vein, and not, as might be expected, in the diffuse type. Broadly speaking, the results are good in 75 per cent. and poor in

25 per cent. There is little doubt that bad nutrition accounts for many of the latter, the process of thrombus formation being slowed down and incomplete. Over-tight bandaging may also have been a factor. When under-nourishment or anaemia is suspected advice is given as to diet. The figures for the heredity factor are given in the hope that they may help to swell the totals of other collectors. The numbers are too small for statistical treatment.

TABLE II

Condition	Perfect	Excellent	Very Good	Good	Poor	Total
Very severe ...	2	0	0	0	0	2
Severe ...	5	8	3	1	1	18
Marked ...	16	20	5	6	6	53
Medium ...	17	12	6	3	3	41
Slight ...	10	10	1	1	1	23
Total limbs ...	50	50	15	11	11	137

All the limbs in Table II were examined one year after the last injection and midway along the seven years that the clinic was operative. At this time 629 limbs had been treated.

TABLE III

Size of Vein	Perfect	Excellent	Very Good	Good	Fair	Poor	Total
Enormous	0	1	0	2	1	1	5
Large ...	2	9	6	9	5	2	33
Marked ...	1	4	5	5	4	11	30
Medium ...	8	11	9	8	3	1	40
Small ...	8	7	2	3	4	2	26
Diffuse ...	2	6	8	4	0	6	26
Total limbs	21	38	30	31	17	23	160

TABLE IV.—Heredity Factor

Result	Present	Marked	Absent	Not Noted	Total
Good ...	39	27	48	6	120
Poor ...	13	15	9	3	40
Total limbs ...	52	42	57	9	160

All the limbs in Tables III and IV were examined three years after the last injection.

A test of the effects of different solutions in a patient with multiple cavernous angiomas brought out the fact that sodium morrhuate 10 per cent. in a stagnating blood stream may cause necrosis. Salicylate was too caustic, but saline 20 per cent. was satisfactory. Many of the naevi were destroyed by diathermy coagulation. The results were not so good as by injection. The scars were dark purple, raised, and indurated with small telangiectases round them, whereas those left after injection were pale and supple with no telangiectases. Sodium morrhuate 10 per cent. produced thrombosis in a vein proof against salicylate-saline after sufficient time in months had elapsed to render it unlikely that the previous injection had influenced the process. We have had no experience of the sugar or quinine solutions, as they do not appear to cause sufficient pain to make them guidable. Sodium morrhuate in 10 per cent. solution appears to cause a slight degree of pain if the solution is cold, and it is possible that heat and cold of moderate degree might impart a guidable property to the solutions used. An investigation into the perception of pain in veins would yield fruitful results. It is an almost untouched field.

We cannot agree that recurrence is unlikely to be related to the solution used merely because the general mode of action of all "sclerosing" solutions is probably similar. Enormous clinical differences seem to occur in reported cases, as judged by the number of injections required and the number of so-called recurrences, especially with sodium morrhuate and the sugar solutions. And throughout we have to bear in mind that the apparent absence of effect of any solution tried on any piece of vein may always be that there is some anatomical peculiarity present whereby the solutions are directed quickly into the deep circulation and thereby rapidly diluted. Success from injection must depend on the closeness with which the chemosis approaches the veins of the deep circulation without actually affecting them.

Failure of Obliteration

Patey,²⁵ in his Hunterian Lecture, figures the section of an excised varicose vein which had been injected at weekly intervals for several months without success, and states that there is evidence that a condition of the vein refractory to thrombosis may arise. The special plate shows a similar condition in a slightly anaemic woman of 54. Large-volume varicosities with numerous anastomoses and impulse to the ankle were present on each side. Eight injections of salicylate-saline cured the left side, but nine failed on the right. Four of these were given at intervals of four and a half, one, two and a half, and four months into the cyst. A thickening and shrinkage of the thigh saphena was produced, but no actual obliteration of it or of the cyst. Both were removed through small multiple incisions from the groin to below the knee. The tributaries did not bleed at all freely. The vein was not adherent to the adjacent tissues. The cyst was supplied by one small radicle only. No competent valves were noticed in the thigh saphena. The result six months later was excellent.

Here, therefore, was a cyst of the saphena supplied by one small radicle and injected many times with strong salicylate-saline solution without showing any obliteration either of the cyst or of the main vein. It was unfortunate that one at least of these injections was not made with a different solution. The sections made of the cyst and of the saphena are shown in Figs. 1 to 5 and the limbs in Fig. 6. (See Plate.)

The following instance, however, shows how difficult it is to come to a correct decision.

A healthy girl of 13½ years, who had not menstruated, had a large varicosity treated. Eight large injections with salicylate (40 per cent.)-saline (10 per cent.) were spread over the ensuing six months. The only effect seemed to be that all the veins became a little thicker in their walls, and here and there the skin over them showed a brown discoloration. The note reads: "The intima of the veins or the blood must be abnormal." Four months after the last injection the result was excellent. The sclerosing process had evidently continued gradually, and what was apparently going to be a case requiring multiple ligation became a good result. Menstruation began a year after the last injection had been given.

In collections of small varices and telangiectases thrombosis may sometimes be followed as a slowly extending inflammation resembling thrombo-phlebitis migrans, except that it does not begin in embolic fashion. It is an excellent reason for giving the first two or three floodings at long intervals, such as three or four months for each limb, and allowing a still longer interval to elapse between inspections, cautioning the patient to report should there be any rapid increase in the size of the veins. But whether this method would give good results when small injections are used may be left for the opinions of those who have had a larger experience of them.

Treatment of Ulcer Cases

Adamson²⁶ has drawn attention to the difficulty of continually maintaining pressure by means of elastoplast applied once a week. Practitioners and patients have mentioned this from time to time at the clinic. We have had little experience of the method. Our view is that the old-fashioned Unna's paste casing, applied in the old way and from toes to knee, is hard to beat. This method has been used with great success throughout the seven years that the clinic was operative. The oedema and blood are drained by elevation of the limb for twenty minutes, while the patient lies supine. The carbolic gauze, cut without selvedge, is applied, as far as possible with even tension on each of its edges, for two or three layers, and the paste worked in with a brush. Two more layers and paste, and a bandage or stocking over all, complete the casing, which is renewed each month. The casing is semi-rigid and *not* elastic. A useful adjunct is to cover it with a non-perforated Martin's bandage, or with a sheet rubber stocking at certain times, so that the patient may lie in a bath without dissolving the casing. At the stage of induration the method is highly efficient, and should be continued for at least three months after amelioration. Injection is performed as and when required. Used in this way the injury to or irritation of eczematous skin which often occurs with other preparations and bandages²⁷ is avoided; neither is it necessary to *forcibly* reduce the oedema. The saving in labour and in initial cost is considerable compared with other methods. Later, at the stage when a simple elastic support is required, a light, porous, elastic webbing bandage, worn if necessary over a thin stocking, will act as the transition stage to light flannel or other support, and is one that the patient sometimes prefers to continue.

Recent Additional Facts and Physical Signs

The practical value of Kosinski's²⁸ painstaking dissection of the veins in 124 limbs lies in his establishing the fact that the small saphena is generally subfascial in the upper half of the leg. The course of the main vein can be traced beneath the deep fascia by the percussion method of Barber,²⁹ and it is then possible to inject it. This frequently gave a good result in our cases where, up till then, there had been failure. In one case a hidden external saphena revealed itself by becoming hard and tender.

The breathing test described by Barber,³⁰ where the changes of pressure caused by the abdominal respiratory movements are seen in the outlines of the limb veins, sometimes as far as the ankle, is useful. It indicates an entire absence of competent valvular mechanism in the affected veins.

In describing the percussion test as venous shock Barber³¹ draws attention to its value in determining the range of thrombosis, the thickening of the vein wall, and the occasional phenomenon of recanalization. He points out (p. 70) that it is due to a wave engendered in the blood by percussion and perceived either visually or by palpation. One of its chief uses is in tracing patent veins when they lie beneath indurated tissue or beneath the deep fascia. MacPheeters,³² who described the test independently some two years later, calls it the "percussion pulse transmission."

X-Ray Examination

Further to the stereoscopic x-ray observations of Professor Sicard and Drs. Forestier and Gaugier in 1928,³³ x-ray screen examinations and the examination of films, after the intravaricose injection of opaque solutions^{34 35 36} have shown the possibility of increased accuracy in diagnosis in these conditions—for example, the extent of the

deep varicosities in early cases. Such an examination would be of especial value in those cases in which curative injections fail either wholly or in part. The published observations are not yet numerous. It would appear that in attempted injection of the whole superficial varicose system of the lower limb from above, with the patient standing and keeping his muscles inactive, the solution is likely to remain in longer and be less quickly diluted by the blood in the superficial than in the communicating and deep veins. If such a patient tilts himself into the horizontal position and the limb is raised, the solution tends to travel by the great and small saphenae. If, however, the injection is made when the patient is lying down it is likely to pass more quickly into the deep circulation. It is also shown that the sclerosing solution reaches the deep veins in concentration. It follows that the increased volume of blood there and its velocity are the factors which save the sclerosing method from causing disaster from deep thrombosis. It has also been shown that muscular movement of the foot may not only aspirate the blood from the superficial to the deep circulation, but when the valving is inefficient may force it in the reverse direction. The observations of the effects of inspiration and expiration are as yet inconclusive for the superficial veins, but expiration tends to aspirate the blood from the deep veins towards the trunk. This confirmation of facts formerly ascertained by laborious dissection, or observed clinically, places the whole subject on a surer foundation. Clinically one of the chief reasons why the massive flooding method of injection from above downwards sometimes fails is that one main area may be completely missed. This is presumably due to the degree of obliquity of the tributary junctions or to their valvular sufficiency there. While it is sometimes possible to negotiate the difficulty with the use of guidable solutions by rapid changes of position, it will be much to our advantage to know beforehand how and where to commence the attack.

The application of the cineradiographic methods recently demonstrated by Dr. Russell J. Reynolds^{37 38 39} promises in the near future a solution of many of these difficulties. The selection of cases suitable for primary ligation might be readily controlled in this way and operation avoided in those judged curable by injection. It is probable that at first the method will be reserved for the investigation of cases which have given imperfect results by the combined methods of ligation and injection, and later on will be used as a preliminary in cases obviously difficult to assess.

The Cause of Varicosity

While stating that his series is not large enough to be conclusive, Turner Warwick⁴⁰ gives it that inadequacy or absence of valves in the perforating, or occasionally in the deep veins, is a factor commonly underlying the hereditary tendency to varicosis. The valvular defect is apparently the primary lesion; there is no evidence of weakness of the wall. This point lies at the root of the whole matter, and in the past a large amount of work has been done to try to settle it.^{41 42} While the basic facts become more and more clear, the cause of them still remains elusive.

Conclusions

From an analysis of the current state of the subject, and from the evidence available in this series of cases, the following conclusions seem to be justified:

1. The injection treatment of varicose veins with salicylate-saline solution is safe and satisfactory in two chief types.
2. A third type of varicosity, when treated in this manner, gives poor results. Although great amelioration takes place a cure does not follow. It is as yet uncertain from the published results if more than amelioration can

be promised by operation, or by operation combined with injection, in this type of case.

3. In a very small proportion of cases injection alone is not entirely successful, and ligation of the internal saphena vein below its upper end is indicated as an adjuvant; but no evidence sufficiently strong has yet been published to justify primary saphena ligation in preference to primary injection with salicylate-saline solution.

4. The classified results of other methods of injection after a three-year interval would, if published, advance the subject considerably. The results given by the non-guidable solutions—for example, quinine, sodium morrhuate, and the sugars—are especially called for.

5. Experience tends to show that almost all cases which have failed to react to other solutions can be successfully treated with salicylate-saline solution.

6. The converse is not true, but sodium morrhuate appears to answer well in short lengths of vein when salicylate fails.

7. Longer intervals than those commonly given between injections are evidently necessary if the accepted pathology of sclerosis is to be the guiding principle in treatment.

8. Improvements in the pathological diagnosis of cases unsuccessfully injected may be expected from the application of the newer radiological methods of investigation.

9. Unna's paste bandage applied in the hydrostatic manner gives much better results than are obtained by the modern elastic-pressure applications.

Our thanks are due to the members of the staff of the Aberdeen Royal Infirmary for their help on many occasions.

REFERENCES

- ¹ Colt, G. H.: *Brit. Journ. Surg.*, 1921, viii, 486.
- ² Jeannel: Rapport à Congrès français de Chirurgie, 1910.
- ³ Colt, G. H.: *British Medical Journal*, November 9th, 1929.
- ⁴ Linser, P., and Zirm, C.: *Derm. Zeit.*, 1916, xxiii, 650; *Munch. med. Woch.*, 1919, lxxvi, 382; *Derm. Zeit.*, 1925, xlv, 22; *Die Therapie der Gegenwart*, 1925.
- ⁵ Fischer, Carl: *Wien. klin. Woch.*, 1918, xxxi, 295; and 1922, xxxv, 11.
- ⁶ Cooper, William M.: *Ann. of Surg.*, May, 1934, p. 799.
- ⁷ Brodie: *Lectures on Pathology and Surgery*, 1846.
- ⁸ Warwick, W. Turner: *The Rational Treatment of Varicose Veins and Varicocele*, London, 1931.
- ⁹ Moszkowicz: *Wien. med. Woch.*, October 20th, 1934, p. 1156.
- ¹⁰ Hanselman, Margrit: *Inaug. Diss. Zurich*, 1926.
- ¹¹ De Takáts, Géza: *Journ. Amer. Med. Assoc.*, 1930, xciv, 1194.
- ¹² Linser, P., and Zirm, C.: *Loc. cit.*
- ¹³ Fischer, C.: *Loc. cit.*
- ¹⁴ Bazelis: *Presse Méd.*, 1924, xxxii, 358.
- ¹⁵ Hemple: *Munch. med. Woch.*, 1924, lxxi, 900.
- ¹⁶ Howard, Jackson, and Mahon: *Arch. of Surg.*, 1931, xxii, 353.
- ¹⁷ Cooper, W. M.: *Amer. Journ. Surg.*, April, 1934, p. 159.
- ¹⁸ Patey and Tatham: *British Medical Journal*, November 11th, 1933.
- ¹⁹ Forestier, J.: *Journ. Amer. Med. Assoc.*, 1928, xc, 1932.
- ²⁰ Payne, R. T.: *Lancet*, 1934, i, 235.
- ²¹ Warwick, W. Turner: *Loc. cit.*
- ²² Meisen, V.: *Acta Chir. Scand.*, 1926, lx, 435; 1927, lxii, 1.
- ²³ Colt, G. H.: *Loc. cit.*
- ²⁴ Theis, Frank V.: *Ann. of Surg.*, 1933, xcvi, 82.
- ²⁵ Patey, D. H.: *Lancet*, August 8th, 1931.
- ²⁶ Adamson, H. G.: *British Medical Journal*, 1931, i, 284.
- ²⁷ Cokkinis, A. J.: *Lancet*, 1933, i, 1168.
- ²⁸ Kosinski, C.: *Journ. of Anat.*, 1926, lx, 131.
- ²⁹ Barber, T. H., de T.: *The Treatment of Varicose Veins by Injection*, Bristol, 1929, p. 70.
- ³⁰ Idem: *British Medical Journal*, 1928, i, 594.
- ³¹ *Loc. cit.* (reference 29).
- ³² MacPheeters, H. O.: *Journ. Amer. Med. Assoc.*, 1931, xcvi, 1114.
- ³³ *Beit. z. klin. Chir.*, 1929, cxxxiii.
- ³⁴ MacPheeters, H. O., and Rice, C. O.: *Surg., Gynecol. and Obstet.*, July, 1929, p. 29.
- ³⁵ Barber, T. H., de T., and Orley, A.: *Lancet*, 1932, ii, 175.
- ³⁶ Patey, D. H., Tatham, R. C., and Nicholas, F. G.: *Lancet*, 1933, ii, 1309.
- ³⁷ *Proc. Roy. Soc. Med.*, June, 1934, xxxvii, 1.
- ³⁸ *Brit. Journ. Radiol.*, July, 1934, vii, New Series, No. 79.
- ³⁹ *Ibid.*, February, 1935, viii, New Series, No. 86.
- ⁴⁰ *Loc. cit.*
- ⁴¹ Gay: *Varicose Disease*, London, 1867.
- ⁴² Briquet: *Thèse de Paris*, 1824.

The dietetic department of the London Hospital is to be enlarged, and the new accommodation will include nine beds for patients and four private rooms, a laboratory, a special kitchen, and a rest room for patients not entirely confined to bed. There are already over 9,000 patients on the books under treatment in this department of the hospital.

THE AFTER-RESULTS OF PSYCHOTHERAPY IN 500 ADULT CASES

BY

MARY C. LUFF, M.D., D.P.M.

ASSISTANT MEDICAL DIRECTOR TO THE INSTITUTE OF MEDICAL
PSYCHOLOGY

AND

MARJORIE GARROD, M.R.C.S., L.R.C.P.

REGISTRAR TO THE INSTITUTE OF MEDICAL PSYCHOLOGY

The attention of the medical profession has, during recent years, been increasingly directed towards psychotherapy, but the claims for this method of treatment are sometimes criticized on the ground that they are too often based on successes in individual cases. This branch of medicine, however, is of such recent growth that it is only now becoming possible to publish series of case results that are at all comparable with those recorded from other medical fields.

Since the foundation of the Institute of Medical Psychology in 1920 as the Tavistock Clinic, the number of patients treated there each year has steadily increased. It has therefore been possible to collect from the Institute's records a considerable volume of material by which the effectiveness of psychotherapy as a remedial measure may be judged. In this paper the results of this method of treatment as applied to 500 adult cases at the Institute are set out in tabulated form, together with explanatory remarks and comments.

The results of psychotherapy cannot fairly be judged by the patient's condition at the termination of treatment. He may then appear very much improved and his symptoms may have completely disappeared, but he has still to prove his capacity to withstand without further breakdown the stresses that life will inevitably bring. Only time can show whether treatment has given him this capacity. Further, a serious criticism made of psychotherapy is that, although benefit may be apparent, the improvement is really due to the patient's reliance and dependence on the physician, and is not likely to be lasting. In fact, it has been suggested that psychoneurotic symptoms indicate an inherent personality defect which cannot be "cured." These criticisms can only be conclusively answered by collecting on a large scale information concerning the remote, as well as the immediate, results of psychotherapy. In this paper a step is made towards the provision of such information. The valuable statistics recently published by Neustatter from the Maudsley Hospital (*Lancet*, April 6th, 1935) give results at the end of treatment only, and it is hoped that the after-results in his group of cases may later become available.

With these considerations in mind we have followed up as a routine all patients discharged from treatment at the Institute since 1930. This "follow-up" covers a period of three years after discharge, and during this time at least three reports on the patient's condition have been collected. The information is usually obtained by letter from the patient himself or from his private doctor. In some cases, however, a return visit is made to the physician who treated him at the Institute, and in others a social worker calls at his home. Whenever possible information from several of these sources is considered. If the patient himself, his relatives and friends, or the Institute physician give continuously favourable reports it is fair to judge the result of treatment as successful.

Assessment of Results

At the termination of treatment a summary of the case is made by the physician, and on his judgement the patient is placed in one or other of four groups, under

JULY 13, 1935

THE BRITISH
MEDICAL JOURNAL

G. H. COLT, ISOBEL S. W. RAMSEY, AND MARGARET M. M. MORRISON: INJECTION TREATMENT OF VARICOSE VEINS

FIG. 1.—Shows fibrotic thickening of the inner coat affecting chiefly one side and encroaching on the lumen.

FIG. 3.—Shows some thickening of inner coat of vein at middle of thigh.

FIG. 5.—Shows some thickening of inner coat of vein below knee.

FIG. 2.—Shows a cystlike bulging at one side of wall of vein, the coats being thinned out at this part. Remainder of vein shows hyaline thickening chiefly affecting inner coat of vessel. Lumen dilated and filled with blood clot.

FIG. 4.—Shows some thickening of inner coat of vein in lower part of thigh.

FIG. 6.—The limbs after injection and before operation. Infra-red plate.