TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some of the major medical disorders met with in general practice.

TREATMENT OF HEART-BLOCK

BY

ROBERT MARSHALL, M.D., F.R.C.P.I.

"Before entering upon the consideration of the treatment of affections of the heart it is necessary to have in mind a clear idea of what we aim at in treating any given individual . . . there are many obscure phenomena which nevertheless are often the object towards which treatment is directed, though in themselves these phenomena are not the signs of disease nor suitable indications for treatment."—Sir James Mackenzie.

In applying these principles to the treatment of heart-block we must first locate the site of the block: this can in most cases be easily done by electrocardiography; but if this is not available some forms of block can be determined by careful clinical examination. Secondly, we must discover the cause of the block. In congenital cases there is frequently, but not invariably, demonstrable evidence of patent interventricular septum. The toxins of rheumatism, diphtheria, influenza, enteric fever, scarlet fever, and syphilis are frequent causes in youth and early adult life; in later years degenerative and coronary vascular changes are most commonly found; tumours, usually secondary, are a rare cause at any age. Thirdly, it must be decided whether the block is adversely affecting the efficiency of the patient's circulatory system; and, fourthly, whether we have any weapons that are likely to be effective in removing the cause of the block or in ameliorating its effects.

It is customary to classify heart-block on an anatomical basis, according to the portion of the junctional tissue of the heart involved, into (1) sino-auricular block; (2) auriculo-ventricular block; (3) bundle-branch block; and (4) arborization block.

Sino-auricular Block

In this form the impulses from the sino-auricular node fail at times to provoke an auricular contraction, and give rise to arrhythmia characterized by long pauses between the pulse beats, during which no heart sound is heard. Its cause is frequently obscure. It is not very rare in young people, is not of serious significance, and may be regarded as "an obscure phenomenon . . . which is not a suitable indication for treatment."

Partial Auriculo-ventricular Block

Prolonged Auriculo-ventricular Interval.—This cannot be detected without polygraphic or electrocardiographic tracings. It is a common finding in acute rheumatic and other infective carditis and is always significant of severe infection. It is relatively rare in the ambulant rheumatic children attending an out-patient clinic. When the apparent cause is a septic focus this should be dealt with, and if a general anaesthetic is necessary gas and oxygen should be used. When the cause is acute rheumatism or other infectious fever the patient should be kept absolutely in bed until the delay in conduction has disappeared or until it seems inevitable that the delay is permanent.

Intermittent Failure of Ventricular Response.—In this form the auricular impulse may occasionally fail to reach the ventricle, giving rise to a dropped beat. This can be distinguished clinically from extrasystole by the fact that the heart is silent when the beat is dropped; if the apparently dropped beat is due to an extrasystolic contraction too feeble to create a pulse beat at the wrist, the sound of that extrasystole may be heard on auscultation over the heart. The auricular impulse may regularly fail to reach the ventricle, giving rise to 2:1 or 3:1 heart-block. In 2:1 heart-block the ventricular rate is frequently 40 to 50, but rises abruptly to double the original rate on exercise. The administration of atropine (1/200 to 1/100 grain) hypodermically, the inhalation of amyl nitrite (m iiij to v), or the ingestion of ephedrine (1/3 to 1/2 grain), may also temporarily double the ventricular rate, but this experiment is sometimes accompanied by considerable distress to the patient, and treatment is better directed to his general condition and the underlying cause. It is important to note that in this and other forms of heart-block the best results are likely to be obtained in syphilitic cases. Potassium iodide is the most valuable drug in such cases, and may be prescribed as follows:

R	Pot. iod.	...	...	...	gr. xv
	Spirit. ammon. aromat.	...	...	...	℥ xx
	Spirit. chlorof.	...	...	...	℥ v
	Infus. gent. co.	...	...	...	ad 3 ss
				t.d.s., p.c., ex. q.	

Mercury is best administered by inunction. One drachm of ung. hydrarg. is rubbed into each of six areas of the body in turn for six successive nights with one night's interval each week for six weeks; less efficacious is its administration by the mouth thrice daily.

R	Hydrarg. cum cret.	...	...	...	} āā gr. j
	Pulv. ipecac. co.	...	...	...	

Bismuth is easily administered in the form of bismostab (0.2 to 0.5 gram) intramuscularly at weekly intervals. It has the advantage that the site of injection is less painful than when mercury has been injected. Intravenous injections of arsenic should not be given when there is anginal pain. (For further details in the treatment of cardiac syphilis the reader is referred to Dr. D. Evan Bedford's article in this series in the *Journal* of June 8th.)

Complete Auriculo-ventricular Block

Complete heart-block is a relatively rare phenomenon. Gilchrist found it in only 31 of 5,000 consecutive electrocardiograms recorded at the Edinburgh Royal Infirmary, representing an incidence in 24,000 patients of 0.57 per cent.; nevertheless, its recognition and significance are important because the symptoms to which it gives rise may constitute a serious medical emergency. The condition may be transient, due to toxic effects, or permanent, due to gross lesion of the auriculo-ventricular bundle. It is sometimes difficult to be certain that the block is permanent. The presence of complete heart-block can usually be determined by bedside examination, and can always be confirmed by graphic methods. In simple bradycardia the ventricular rate at rest is rarely less than 50 per minute, and rises quickly with excitement, exercise, the injection of atropine, or the inhala-

tion of amyl nitrite. The bradycardia of heart-block is unlikely to be influenced by the inhalation of amyl nitrite or by the injection of small doses of atropine, although Gilchrist has shown that 1/30 grain of atropine injected intravenously will raise the ventricular rate in varying degree. These drugs will cause the bradycardia of 2:1 heart-block abruptly to double in rate, and, when their effect has passed, to regain the former slow rate with equal suddenness. The apparently slow pulse which is found as a result of digitalis overdosage when extrasystoles alternate with normal beats, the extrasystoles failing to cause an appreciable pulse wave at the wrist, can always be distinguished by careful auscultation from the much more serious pulsus alternans. In the former the muffled sound of the early beat may be heard, followed by its compensatory pause; in the latter the cardiac cycles sound alike and there is no compensatory pause.

In complete heart-block the heart sounds vary in intensity, and sometimes the dissociated auricular sounds may be heard over the right auricle or at the apex. Inspection of the cervical veins may show the auricular waves occurring independently of the apex beat. Complete heart-block due to congenital causes is rare: Joseph Lewis found approximately forty cases reported in the literature. It is common, but not invariable, to find evidence of patent interventricular septum; there is a history that bradycardia was observed in infancy, and that frequently there were syncopal attacks in infancy or childhood. The ventricular rate is higher in congenital cases, averaging 43 per minute. I have seen a transient idioventricular rate of 72 per minute in a child with dextrocardia.

Acquired complete block is usually degenerative rather than inflammatory in origin, is commoner in later life, and is to be regarded as indicative of a widespread affection of the heart muscle with special involvement of the neuro-muscular mechanism. Therefore there is little to be gained by ill-considered efforts to interfere with it. Where there is evidence of toxic cause this should be treated, and complete rest be insisted upon until it becomes apparent that the block is permanent. A Wassermann reaction should be obtained in every case in the hope that the cause is syphilis and that it may yield to treatment. It is during the stage of onset that the patient is most prone to symptoms of the Adams-Stokes syndrome, for which immediate treatment is imperative.

The Adams-Stokes Syndrome

The syndrome varies in intensity from transient vertigo to syncopal attacks without or with convulsions. Vertigo is the mildest form, and this may recur at varying intervals. The syncope is abrupt in onset. The face is at first pale as the patient loses consciousness and reddens as he "wakens" with a long-drawn sigh. Convulsions resembling those of epilepsy may arise, and are of still more serious significance. The symptoms are dependent on the slow ventricular rate and bear a direct relation to it. Sir Thomas Lewis states that

"cessation of the heart-beat for a single period of five to ten seconds is sufficient to produce unconsciousness; arrest lasting from fifteen to twenty seconds causes obvious distension of the veins, occasionally cyanosis, and deep breathing; a little twitching of the face and upper limbs is added in pauses of longer duration... death results when the ventricular arrest lasts from one to two minutes."

The treatment of Adams-Stokes attack is the immediate hypodermic injection of 0.25 c.cm. of adrenaline. It is important that the preparation should be fresh (discoloration is evidence of oxidation). The effect of the injection may be increased by massage over its site. If there is no immediate response, and if convulsions occur, 0.5 c.cm. should be injected directly into the heart. Oral

administration of ephedrine should be commenced: a dose of 1/3 grain every eight hours may be increased to 1/2 grain if symptoms recur; the aim should be to give the minimal dose required to increase the ventricular rate. It is noteworthy that treatment by adrenaline and ephedrine will in most cases prevent the recurrence of Adams-Stokes attacks, but does not remove the block.

Atropine is a less reliable drug in these cases. It differs from adrenaline in that it produces a greater acceleration in ventricular rate when the initial ventricular rate is high, whereas adrenaline causes proportionately greater acceleration when the initial rate is slow (Gilchrist). It should be administered intravenously and in doses of 1/50 to 1/30 grain. It is more likely to be effective when vagal overaction is a factor in slowing the ventricular rate. Barium chloride, in doses varying from 1/2 grain to 8 grains a day, has been prescribed, and one has seen the pulse rate rise to normal following its use. It appears to be very doubtful whether the block in such cases has not been transient in nature and the increased rate independent of the drug.

There is probably no condition in which success depends more on the temperament and on the co-operation of the patient. If he is of a calm and unhurried habit he can be "taught to live with his lesion." It should be explained to him that his heart is now acting on a slower gear, to which his circulatory system will become accustomed, and that he must regulate his life accordingly. During the period when the altered rate is recent, and especially if the Adams-Stokes syndrome is present, he must have a prolonged rest in bed. Return to limited activity must be gradual, and if there is any tendency to vertigo or syncope he should be warned to avoid the risk of attack in traffic or in hot and stuffy atmospheres.

Digitalis is definitely contraindicated when heart-block is incomplete or transient, but in established cases, when signs of congestive heart failure arise, it may be given, and its excellent results suggest that these are the effect of its direct action on the ventricle itself.

Complete heart-block is compatible with life, and cases are recorded where it has been present for ten, twenty, or even, as in Harris's case, for twenty-eight years. The presence of syncopal attacks affects the prognosis very materially for the worse.

Bundle-Branch Block and Arborization Block

Bundle-branch block and arborization block cannot be recognized without electrocardiography. As Crichton Bramwell has recently pointed out, gallop rhythm is not directly associated with such lesions. The correct localization of the block as to right or left branch is apparently unsettled, but certainly the type formerly designated as right bundle-branch block is the commoner. The prognosis is bad in both types, except perhaps in those of syphilitic origin, and is worse in those formerly regarded as left branch lesions. Occasionally both types of electrocardiographic complex may occur either in long sequences or, as in K. D. Wilkinson's fatal case of diphtheria, in alternate beats. The treatment is that of chronic myocarditis.

M. Braschi, writing in *La Pediatria* (April 1st, 1935) on leucocytosis in whooping-cough, reports as follows from observations on twelve cases: Leucocytosis may be slight or entirely absent, and when it does exist, is subject to wide oscillations which are quite independent of the stage of the disease. The paroxysms of coughing do not appear to cause any remarkable increase in leucocytosis, whereas a definite increase may be caused by parenteral administration of adrenaline or nucleinate of soda. Braschi comes to the conclusion that leucocytosis in whooping-cough is mainly due to the action of the infective agent upon the organs.