AN ATYPICAL FORM OF TUBEROUS
SCLEROSIS

REPORT OF A CASE

BY

R. M. STEWART, M.D., F.R.C.P.ED.

MEDICAL SUPERINTENDENT, LEAVESDEN MENTAL HOSPITAL

(With Special Plate)

Multiple tumours in the skin, epileptiform seizures, and retarded mental development form a triad of signs sufficiently arresting to make the diagnosis of tuberous sclerosis a comparatively easy task for the clinician, but in the so-called *formes frustes*, in which one or more of the cardinal features are absent, the underlying pathological process may easily escape recognition, and in those cases in which epilepsy is the sole manifestation a correct diagnosis may be impossible.

The following case is of interest, not only because the patient was a married woman of normal intelligence, but also by reason of the strict limitation of the pathological process to the central nervous system, the skin and visceral organs being entirely free from tumour formation. Both for the clinical notes and for the opportunity of examining the nervous system I am indebted to Dr. T. W. S. Hills of Watford.

History of Case

In May, 1932, a married woman, aged 21, whose previous history and upbringing, save for the usual ailments of childhood, had been in every way normal, fell suddenly while in the act of advancing to the pay desk of a hairdresser's shop. At the inquest it was disclosed that the hair wave she had received had involved nothing more than the application of an ordinary gas-heated iron to the hair. The witness who described the seizure stated that her client suddenly looked up in a surprised way as though something had happened, and immediately afterwards fell on the floor, "making a terrible noise in the throat as if she were trying to breathe." The pulse, after being fluttering, became imperceptible, and artificial respiration was tried without result.

At the post-mortem examination made by Dr. Hills a persistent and enlarged thymus gland was found; the heart, kidneys, and other organs were entirely normal, but the brain contained a number of firm nodules in the substance of the cerebral convolutions. The view was advanced at the inquest that the deceased had died in an epileptic fit, which was regarded as being the first and only manifestation of tuberous sclerosis. A verdict of death from natural causes was returned.

Examination of the Nervous System

The brain, which weighed 1,350 grams, was large, symmetrical, with well-formed convolutions of normal pattern, and on cursory examination showed no distortion of the cerebral gyri. To the touch, however, it was obvious that embedded in the grey matter of both cerebral hemispheres there were a number of small discrete tumours of a hard pea-like consistence. Only in one or two situations was there naked-eye evidence of nodules in the shape of slight broadening and pallor of the gyri where these overlaid the tumours. On the lateral aspect of the right cerebral hemisphere three nodules were found in the parieto-occipital area; on the mesial surface, four nodules; on the lateral aspect of the left cerebral hemisphere, six nodules; and on its mesial aspect, four nodules; a total of seventeen tumour masses. When cut into the masses showed pallor of the affected grey matter, so that in their neighbourhood the normal sharp transition between grey and white matter was lost.

Horizontal section of the right cerebral hemisphere in Marie's plane exposed a hard, oval-shaped mass, white in colour, and about the size of a pea, lying in the anterior end of the globus pallidus; part of the tumour projected as a small bead-like body into the cavity of the lateral ventricle. A similar, but smaller, mass was also found in the wall of the

lateral ventricle of the left cerebral hemisphere below the head of the caudate nucleus. The cerebellum was relatively large, but appeared normal, as did the pons and medulla oblongata.

In the immediate neighbourhood of the tubera the microscopical appearances were quite characteristic, the cyto-architectonic arrangement being markedly disturbed, many of the pyramidal cells occupying abnormal positions. The myelin sheaths were scanty and poorly stained, and throughout the sclerotic areas a dense felt-work of glial fibres was present.

Undifferentiated "large cells," many of grotesque shape and with numerous coarse processes, were found in groups in the sclerotic nodules; in their morphological characters they resembled giant glial cells rather than deformed ganglion cells. Calcified capillaries and large deposits of calcium salts in the form of rounded concretions were present in several of the tumours.

Sections of the small tumour projecting from the wall of the right lateral ventricle revealed the presence of a dense gliosis, made up of fibrillary strands in which a certain number of spindle-shaped cells were present. In other cortical areas where no nodules existed the histological appearances showed no obvious departure from the normal.

Discussion

Epilepsy and mental deficiency occur in combination in so many different forms of nervous disorder that unless there are present tumours of the skin or the rare phacomatous tumour of the retina, it may be quite impossible to establish the diagnosis of tuberous sclerosis during life. Even more interesting are those cases in which the intelligence is unimpaired. Schuster,¹ and more recently Critchley and Earl,² have reported examples of this atypical form, but in the case reported above not only was there no evidence of mental defect, but the patient was entirely free from cutaneous manifestations, and until the attack which heralded her death had never suffered from any form of epilepsy. Still more remarkable was the complete absence of visceral tumours. The possibility of their having been overlooked at the post-mortem examination can be eliminated in this case, since the physician who conducted it was familiar with the condition, and was therefore particularly careful to examine the heart, kidneys, and other visceral organs for their presence.

The condition being essentially a congenital one, it is interesting to note that in this case the presence of nearly a score of small tumours in the brain was yet compatible with normal mental development. Their small size and the almost entire absence of diffuse morbid changes, which can usually be found in areas remote from the immediate neighbourhood of the sclerosed cortical zones, may perhaps afford an explanation for the *formes frustes* in which no evidence of mental inferiority can be found.

REFERENCES

- ¹ Schuster, P.: *Derm. Woch.*, 1913, lvii, 1475.
- ² Critchley, M., and Earl, C. J. C.: *Brain*, 1932, lv, 311.

In connexion with a report of two cases of suicidal thallium poisoning, E. Unseld (*Med. Welt*, April 6th, 1935, p. 487) remarks that in Germany thallium-containing rat or mouse poisons are easily procured—their only recommendation to the intending suicide, for thallium poisoning has often a long period of latency, and always has a protracted clinical course of two to three weeks in lethal cases, or a convalescence taking many months. The outstanding clinical feature is alopecia occurring after three weeks—assuring the diagnosis. Others are: extremely severe pain from toxic polyneuritis with much muscular atrophy (the legs being chiefly affected) and ataxia; morbid psychic conditions with delirium; severe colic and constipation; and dry tongue and an anidrosis. Thallium is demonstrable in the urine for many weeks. Treatment is purely symptomatic.

R. M. STEWART: AN ATYPICAL FORM OF TUBEROUS SCLEROSIS

FIG. 1.—Three tumour masses are indicated by crosses.

FIG. 2.—Four large undifferentiated cells from an area of cerebral gliosis.

FIG. 3.—Large abnormal cells in the deeper layers of the cerebral cortex.