TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some diseases of the nervous system met with in general practice.

SYDENHAM'S CHOREA

BY

C. M. HINDS HOWELL, M.D., F.R.C.P.

It may not be out of place to preface what I have to say about the treatment of this condition by referring briefly to the diagnosis. There are a number of diverse conditions which give rise to involuntary movements, and some of them may cause difficulty in diagnosis. It is important to remember that, as well as by movements, chorea is characterized by weakness and diminished tone in the muscles of the affected limbs. In some cases this weakness may be much more obvious than involuntary movement. The distribution may be hemiplegic, and, unless the child is closely watched, such involuntary movements as there are may escape observation. In cases of this type it is not difficult at first to make an error in diagnosis. The weakness in these—and also in those, by far the commoner, in which involuntary movements occur—is shown by lack of sustained effort, but in some instances it may be so pronounced as to suggest actual paralysis.

Chorea occurs much more frequently in girls than in boys, and is rarely met with after the age of 20, except in association with pregnancy (chorea gravidarum), when it is more likely to arise in first than in subsequent pregnancies, and is more common in the unmarried woman.

Tics consist of involuntary movements, but should not give rise to confusion, as the movements are co-ordinated and purposive and there is no loss of power, in marked contrast to what obtains in chorea. Hysteria may give more trouble in diagnosis, but in this disorder the movements are more stereotyped than in chorea, and other manifestations of hysteria are usually to be found (anaesthesia, etc.). In both chorea and hysteria the movements increase under observation. Choreic movements may be observed during the acute phase of encephalitis lethargica, but the ocular palsies so common in the latter are unknown in the former. Post-hemiplegic chorea (athetosis) is unlikely to cause difficulty in diagnosis. Sydenham's chorea does not occur in adults (except as chorea gravidarum), hence Huntington's chorea and the hemichorea following vascular lesions in the regio subthalamica or corpus Luysii can easily be distinguished without difficulty. A history of previous rheumatism or its association with the attack, and the presence of pericarditis or endocarditis, will serve to confirm the diagnosis.

General Management and Diet

The onset of chorea may be sudden or preceded by a period in which the child appears clumsy and drops things or else exhibits an unusual mental dullness. In treatment it is important to remember (a) that acute rheumatism is closely associated with the condition, and (b) that the basis of the involuntary movements, loss of power, speech, etc., is an encephalitis. The first point indicates the use of antirheumatic treatment, while the second imposes the necessity of quiet, the exclusion of anything likely to excite the patient, and the choice of a suitable nurse, professional or otherwise, who will not fuss or worry the child.

The child must be put to bed in a room which is well ventilated, not overheated (temperature 65° F.), and in which the light is diffused. The head of the bed should be towards the light. It may be necessary, should the involuntary movements be severe, to fix padded sides to the bed to prevent the child from falling out. In small children a wide bandage fixed across the body and secured to the frame of the bedstead may be helpful. In this as in other rheumatic conditions blanket sheets are useful in order to prevent chills, and the child's night-dress should be flannel and be changed as often as may be necessary if there is much perspiration. The room must be kept quiet, and other children and visitors excluded during the acute phase of the illness. It is important that the child should be nursed as a case of rheumatic fever, kept as flat as possible, and not allowed to exert itself in any way. It should be blanket-bathed and not allowed out of bed.

Diet should be as nourishing as circumstances permit. There are two factors to be considered in this connexion: (a) the patient's temperature, and (b) severity of the involuntary movements. If the patient's temperature is high the diet must be liquid: milk, junket, custard, glucose lemonade, and fruit are all suitable; but meat, soups, and meat extracts are best avoided. The milk may be citrated or given with soda-water.

Violent involuntary movements may make nasal feeding with a tube necessary, while in less severe cases a short rubber tube on the spout of a feeding cup will suffice. For a child of 10 years the quantity of milk should be not less than two pints in the twenty-four hours. In afebrile cases, in addition to the above, eggs, fish, chicken, vegetables, and so on may be allowed, but butcher's meat is best avoided. It is probable that for a time the child will require hand feeding.

Drug Therapy

As chorea is rheumatic in origin salicylates or aspirin should be prescribed. Aspirin appears to be more helpful than sodium salicylate. Ten grains of aspirin may be taken three times daily by a child from 8 to 14 years of age, with suitable alterations for those younger or older. A calcium compound of aspirin, which is easily soluble, has recently been put on the market, and is to be preferred as less likely to cause indigestion—dose: 10 to 15 grains freshly dissolved in water and given three times daily.

Sedatives may be necessary to ensure sleep and quiet the movements. Bromide and chloral, 10 grains of each, may be taken one, two, or three times daily, according to the patient's condition. In very severe cases 1/100 grain hyoscine may be injected subcutaneously and repeated as may be necessary, or the following mixture may be helpful:

R. Hyoscin. hydrobrom.	...	gr. 1/150-1/100
Acid hydrobrom. dil.	...	℥ xv
Aq. chlorof.	...	ad 3 ss

Urethane is another drug which is useful where there is great restlessness, in doses of 5 to 10 grains, repeated as may be necessary. Among other drugs which have proved helpful are arsenic and nirvanol. The former may be given as Fowler's solution in doses of 3 to 10 minims three times a day, a careful watch being kept for any symptoms of intolerance. If the child is febrile or has

any evidence of rheumatic infection of the heart or pericardium, arsenic should not be prescribed until the active infection has been checked by salicylates or aspirin. But in non-febrile cases with no evidence of cardiac infection arsenic may be used from the beginning of treatment. Nirvanol is a drug which has had a considerable trial, and in some cases has seemed to produce good results, but its use is not unattended with risk of toxic symptoms. It is prescribed in daily doses of 0.3 gram (total) until a rash appears with some pyrexia, this is accompanied by an eosinophilia. In no case should administration be continued for more than fourteen days. Conjunctivitis or oedema of the eyelids is an indication for immediate cessation.

Protein shock has been recommended in America as likely to curtail the duration of the disease. I have tried it in several cases in older children who are afebrile and without cardiac complications. It has certainly appeared beneficial, and is induced by intravenous injections of T.A.B. vaccine in a dose of ten millions, repeated with increasing dose (20, then 30 or 40 millions) every fifth day. These doses are smaller than those stated to yield good results in shortening the duration of the disease, but in my opinion they provoke sufficiently satisfactory febrile reactions. It is very important to ensure sleep, and the drugs already mentioned as sedatives will be found useful for this purpose. Bromide and chloral in the doses mentioned will as a rule serve the purpose, but if sleep is not obtained by this combination medinal, 3 to 5 grains, according to the patient's age, may be administered.

(Chorea gravidarum is treated in a similar manner to that described. The condition of the patient is rarely so severe as to necessitate termination of the pregnancy, but there are cases in which the physical and mental condition of the patient may make this procedure necessary.)

Importance of Convalescence

Convalescence must be prolonged; its length will be determined partly by the condition of the heart and partly by the exhaustion produced by the disease. Even if there is no evidence of endocarditis the heart is frequently dilated, and the child should be kept in bed until the apex beat has returned to its normal position. A liberal diet should be given, and the patient should, if possible, be sent for at least a month to some quiet place on the South or West Coast. The air on the East Coast is too "exciting" for a child recovering from chorea. During this period an iron tonic may usefully be administered, such as syrup. ferri phosphatis, 1 drachm, thrice daily after food; or ferri carb. sacc., 10 grains thrice daily; or a mixture such as:

R̄	Ferri et ammon. cit.	...	...	gr. x
	Liq. arsenicalis	...	...	℥ iij
	Aq. chlorof.	...	...	ad 3 ss
	3 ss t.d.s., p.c.			

If the child is of school age the question of the date of returning to school must be considered. The child should not return to school for a period of three months from the onset of convalescence, and must be allowed to go easy at first. There is always the possibility of a relapse to be considered, and this must be borne in mind when deciding the amount of school work which should be undertaken and also the proper management of the child during its life at home. But in no circumstances should the child be overloaded with school work. Harmful as this is at any time, it is particularly to be avoided in the case of a child who has recently had chorea. Finally, it is important to inspect the tonsils very carefully, and if there is evidence of definite infection they should be removed.

THE WORLD TOUR AND THE FUTURE

The following article, written on board s.s. "Rajputana" approaching Marseilles, has been received from Dr. S. Watson Smith.

I

In the course of the century of its existence, no single year can have been so important to the welfare of the British Medical Association as 1935, the year of the first World Tour.

In the past, there have been occasions when, dictated by a change of policy, the Association has appeared to reach a turning-point in its career; and we can look upon this tour as marking yet another, perhaps the most important, for the reasons that: (1) This year of the Melbourne Meeting marked the first occasion upon which the Association has travelled round the world to meet in succession representative members and friends in almost every part of the great British Empire, effecting a personal contact with those whose interests are also ours, and learning on the spot of questions and difficulties that beset them; (2) The tour has achieved the cementing of friendships and assisted in binding us all together in what is surely one of the greatest of all causes, so affording greater strength and power to the Association; (3) It has brought to light certain new facts as well as grievances that for some time have been fomenting in several overseas Branches, which it should now be possible to redress and, we hope, remedy, a better understanding being thereby produced as between these Branches of the Association and they with the parent body.

II

We recognize that we have reached our present position but slowly and with many changes: that the history of the Association is a study in compromise, and a study of the influences between which compromise has had to be made. Continuity of policy and administration have, in the very nature of things, been interrupted and discontinuous. But if we have some knowledge as to our position at home, how are we situated as to the position of our Association in the various parts of the Empire? Not only do we find that the British Medical Association stands in a position almost unassailable everywhere, having a greater influence and power now than might have been supposed; and more than ever have we found it recognized that the B.M.A. is for the public good and necessary to the wise and effective administration of all health services abroad as well as at home.

Wherever the tour has taken us, we have been received with an overwhelming enthusiasm and hospitality and have made new friendships which we trust will be lasting. The great extent of our influence as an association and as the mouthpiece of the profession can now be the better gauged and appreciated. So can our position in relation to the community and the State.

It is our considered opinion that at the present moment our representatives abroad have greater difficulties to face in finding and holding members and in maintaining the prestige of the Association than ever before. It seems clear that we can no longer talk of unity of organization or of consolidating the position we hold. Rather does it seem that, to be progressive, we must give greater attention to local conditions and needs and to organizing, granting at the same time a greater measure of autonomy and conceding what is reasonable to the various Branches and Divisions abroad.

We must take the large view, not interpreting our duty in any narrow sense; and in so doing it seems to me we should be assisting in one way and another in the uphill struggle taking place in certain areas. But we must not confuse the "virtue of breadth with the vice of uncer-