Pringle and Teacher found evidence of healing in some of their cases. This led us to examine those oesophagi where the appearances were in the least suggestive of ante-mortem digestion. In very many of these cases evidence of vital reaction in the form of haemorrhage from small vessels and leucocytic infiltration was found, so that ante-mortem digestion in its lesser degrees is probably a much commoner condition than is imagined.

The well-marked forms of ante-mortem digestion of the oesophagus are specially common in severe toxic and infective conditions, especially following operations. The slighter degrees mentioned were found to occur most commonly in cases where there had been a debilitating illness of no special kind. Very many of these cases show evidence of healing, seen macroscopically by the rounding off of the islets of epithelium left in the damaged area. This fact suggests that if treated early the condition may not be so hopeless as it might often seem. The rational treatment would appear to be to neutralize the acid in the oesophagus by frequent small doses of alkali and the soothing of the inflamed surfaces by some emollient such as gum arabic, olive oil, or liquid paraffin.

My thanks are due to Mr. Patrick, who has given his permission to use the clinical history of the case, and to Professor J. Shaw Dunn for his kind help at all times.

BIBLIOGRAPHY

Pringle and Teacher: *Brit. Journ. Surg.*, April, 1919, No. 24.
Pringle, Stewart, and Teacher: *Journ. Path. and Bact.*, 1921, p. 396.

MASTOID RADIOLOGY*

BY

W. O. LODGE, F.R.C.S.Ed.

HONORARY SURGEON TO THE OPHTHALMIC AND AURAL DEPARTMENT
OF THE ROYAL HALIFAX INFIRMARY

(With Special Plate)

A reliable source of information upon this subject is Dr. Mittermaier's *Die Krankheiten der Nasenbenhöhlen und des Ohres im Röntgenbild*, published by Georg Thieme of Leipzig in 1934. This is an atlas with 144 pages and 213 illustrations. A leading article in the *Lancet* of June 2nd, 1934, referred to contributions by Frey, Cottentot, and Thienpont, and summarized Witmaack's theories. For examining a number of my patients suffering from mastoid disease, my thanks are due to Dr. Franklyn and his assistants.

Illustrative Case

A man, aged 21, had had acute suppurative otitis for eight days, with copious muco-purulent discharge and slight mastoid tenderness. Temperature, 99° F.; pulse, 96. Fig. 1 (on Special Plate) is a skiagram of the healthy ear in Schüler's projection. The temporo-mandibular articulation, auditory meatus, and zygomatic and mastoid cells are visible. The sinus plate crosses the cells obliquely. Fig. 2 (on Special Plate) shows early Bezold's mastoiditis on the affected side. The cells are becoming disintegrated. The antral region is densely obscured. These changes are more apparent on comparing the two skiagrams as a whole than on minute examination of any one cell. Note, however, the triangular breach in the inner wall of the tip of the mastoid process.

At operation pus was found under pressure. Digastric muscle fibres were visible through the triangular erosion, and formed the inner wall of the abscess.

Schüler's method of projecting the temporal bone clear of overlying shadows is shown in Fig. 3. This diagram is a coronal section of the head with an arrow passing through the auditory meatus at an angle of 30 degrees, to indicate the axis of the central ray. For comparison, the other ear is projected at 150 degrees in the same plane.

* Read at a meeting of the Bradford Medico-Chirurgical Society on March 12th, 1935.

Stenver's projection is suitable for deep-seated affections. The central ray crosses a horizontal section of the head at 45 degrees, but the trajectory is elevated 10 degrees out of this plane to cast the shadow of the petrous temporal bone above that of the zygoma. A dozen anatomical points are recognizable in the skiagram.

Axial projections are informative in traumatic cases. The central ray crosses a sagittal section of the head as near 90 degrees as possible. In the skiagram the thickness of the mandibular shadow shows whether the projection was submento-vertical or vertico-submental.

Provided that obliquity of incidence is in the plane of the laminae of the Bucky diaphragm, x rays need not strike the film perpendicularly. A head clamp is not indispensable.

Sometimes, by having serial examinations, one is able to refrain from operation—for example, in pregnancy. When there are several mastoids on the operation list radiography conduces to speed, anatomical variations having been visualized beforehand.

FIG. 3.—Schüler's projection.

In well-pneumatized bones even the oedema of the overlying tissues due to a furuncle suffices to obscure the cells. In examples of arrested pneumatization, dating from otitis in infancy, the signs of inflammation are less distinctive.

Owing to a deposition of lime salts the mastoid cortex may appear as though its boundaries had been overlined with a J pen. This appearance suggests a previous attack of mastoiditis, which has subsided without operation. A contracted meatus implies chronicity. Such data are helpful, especially if the history is unreliable, in deciding whether a simple or a radical operation is indicated. If the semicircular canals show clearly in Schüler's projection cholesteatomata may be suspected. Cholesteatomatous cavities are best seen in stereoscopic skiagrams—they are smooth-walled.

If there is an erosion of the sinus plate Queckenstedt's test will show whether the sinus is thrombosed.

J. François (*Thèse de Paris*, 1935, No. 374) states that from the middle of the seventeenth century to the second part of the nineteenth severe epidemics of malaria, with a high fatality rate, occurred at Rochefort-sur-Mer. Since then there has been a progressive decline owing to draining of the marshes and other hygienic measures undertaken in the urban and suburban districts. The disease, however, is not entirely extinct but still persists, although it has undergone considerable attenuation. Benign tertian is the commonest form and larvate forms are frequent, especially in children, and are often overlooked. At the Naval Aviation Centre, between Rochefort and Soubise, four or five cases of indigenous malaria occur yearly in a military population of about 1,500 men. The persistence of the disease is partly due to the presence of swarms of anopheles and the large number of chronic cases of malaria and collections of stagnant water.

R. D. B. WRIGHT: STRANGULATION OF TRAUMATIC
DIAPHRAGMATIC HERNIA NINE MONTHS AFTER
ORIGINAL ACCIDENT

Post-mortem appearances. "A" points to tear in diaphragm, with coin in opening.

ALEXANDER LYALL: ANTE-MORTEM DIGESTION
OF THE OESOPHAGUS

Section of the oesophagus $\times 100$; (a) Layer of brown blood-stained necrotic material; (b) leucocytic infiltration.

W. O. LODGE: MASTOID RADIOLOGY

FIG. 1.—Normal mastoid region.

FIG. 2.—Bezold's mastoiditis.