PREMEDICATION *

BY

F. W. GREEN, M.B., CH.B.

HONORARY ANAESTHETIST, ROYAL MELBOURNE HOSPITAL, AND
EYE AND EAR HOSPITAL, MELBOURNE

The last ten or twelve years have shown an enormous increase in the number of drugs used for pre-anaesthetic medication, and a controversy has raged amongst anaesthetists as to which is the best premedicant, some supporting the claims of one, some of another, the choice depending on the personal predilection of the various administrators. From this difference of opinion three points emerge clearly:

1. That nervous and highly strung patients, and especially children, should receive some form of sedative drug treatment to ameliorate the psychic effect inevitable whilst waiting to be brought to the theatre.
2. That adequate premedication greatly decreases the amount of general anaesthetic needed subsequently.
3. That there is no form of premedicant treatment that is applicable to all cases. Each case must be considered as a separate entity and a suitable drug administered.

Towards the latter end of last and the beginning of the present century it was a common practice to give 0.016 gram (1/4 grain) of morphine as a preliminary to a chloroform anaesthesia often lasting two or more hours, without any apparent ill effects. What would be the remarks of a present-day surgeon should any such combination of anaesthetics be suggested?

For the purpose of this paper premedicant narcotic drugs may be divided into three groups according to the method of administration, and only those in common use will be discussed.

Drugs Given by the Mouth

(a) *Chloretone*.—In at least one of our leading public hospitals it is the custom to give chloretone 0.65 to 0.97 gram (10 to 15 grains) as a routine before all ether anaesthesias. The patients appear to be as comfortable as with a small dose of the other narcotics, and the after-vomiting is certainly less. Although the patients are rarely brought to the theatre asleep, occasionally there is complete forgetfulness of the period immediately prior to the operation.

(b) *The Barbiturates, Sodium Amytal, and Nembutal*.—If these drugs are used at all the dose should be sufficiently large to ensure that the patient is asleep before the general anaesthetic is commenced, as small doses are very uncertain and variable in their action. The use of sodium amytal should be confined to robust and virile types of patients, and the best results will be obtained by administering it in divided doses: 3 grains on the night before operation, 3 grains three hours before, and 6 grains two hours before. With this dose the patient will almost invariably be brought to the theatre asleep, and will remain so for several hours. The drawback to the use of this drug is the fall of blood pressure which it causes. This is very great in some cases, although much less with oral than with intravenous administration. The use of sodium amytal is very often followed by a period of excitement. This, however, can usually be controlled with morphine. Nembutal should be given in 3-grain doses half an hour before operation. It is quicker in its action than sodium amytal, the fall of blood pressure is less, and the period of narcosis much shorter. On the other hand, its sedative effect

is much less and its action very variable. It should be reserved for women and the less vigorous types of male patients. Pernocton, sodium alurate, pernoston, and various other barbiturates have not been used to any great extent in this country, but the foregoing remarks apply in some degree to them all. It is often the custom to combine the use of morphine with the barbiturates. This should never be done, as the long list of fatalities goes to prove.

Drugs Given Intravenously and Subcutaneously

(a) *The Barbiturates*.—Most of these when first introduced were administered by the intravenous route, but this procedure for various reasons is gradually being abandoned in favour of oral administration. The fall of blood pressure is greater than when the drug is given by the mouth, and patients with an idiosyncrasy to the drug are more liable to receive an overdose, since the administrator has little control once the injection has been made into the vein. Of late sodium evipan has come into popular use. It, however, causes a greater though more evanescent fall of blood pressure than the other barbiturates, and a degree of narcosis equivalent to the third stage of anaesthesia. It appears to be rather a potent drug to use for this purpose until it has been further tried out as an anaesthetic *per se*.

(b) *Morphine and its Derivatives*.—In years gone by it was a common practice to give a small dose of morphine whatever the nature of the surgical procedure and the physical condition of the patient may have been. However, as it is a respiratory depressant and increases the post-anaesthetic vomiting, its use is now becoming much less frequent. A still further deterrent, especially in teaching hospitals, is its effect on the pupil whereby the depth of anaesthesia is obscured. Hyoscine is often employed with morphine. It is doubtful if this practice is ever advisable on account of its depressant action on the respiratory centre. If used it should only be in cases where rebreathing is to be practised, as then the carbon dioxide acts as a stimulant. Even then it should never be used before the administration of ethylene, as very frequently a stoppage of respiration has occurred with this combination, ethylene being also a respiratory depressant.

Drugs Administered per Rectum

Drugs given by the rectal route have a much more consistent action than if given orally, as their absorption in the stomach is inhibited in nervous and frightened patients. Paraldehyde and avertin are the two narcotics which are commonly administered in this manner.

Paraldehyde is given in doses up to 0.56 c.cm. per kilo (1 drachm per stone) of body weight. One part of paraldehyde is shaken up with ten parts of water and injected one hour before operation. The main drawback to its use is the uncertainty of its action, the same dose producing widely different effects in different patients. It also appears to cause small haemorrhages in the mucous coat of the stomach, indicated by coffee-ground vomit.

Avertin appears to be the most generally suitable of all the premedicants, provided that the dose is small and in proportion to the age, physical condition, and weight of the patient. There is a fall of blood pressure, but if the amount given is within reasonable limits this is negligible and transitory in its duration. The dose should be from 0.05 to 0.1 c.cm. per kilo of body weight. Larger amounts than this should never be used under any circumstances. Half an hour after the injection the patient is brought to the theatre quietly asleep, and very little general anaesthetic will be necessary for one or two hours. After-excitement is negligible, but should it occur it can be easily controlled with a small dose of morphine.

* Read in opening a discussion in the Section of Pharmacology at the Annual Meeting of the British Medical Association, Melbourne, 1935.

Where muscular relaxation is required and a gas-oxygen anaesthetic is to be used some form of pre-medication is essential. The use of a suitable hypnotic will usually obviate the necessity of adding ether to the gas employed.

In certain operations in which it is essential that there should be no post-anaesthetic vomiting, a preliminary administration of a small dose of avertin followed by anaesthesia with nitrous oxide and oxygen is very satisfactory. Especially is this the case in some eye operations, where vomiting may absolutely ruin the result.

Value of Atropine

The foregoing remarks have referred only to sedative premedicant drugs, but before concluding some mention must be made of the use of atropine. This is the most valuable premedicant of all, and should be used before

all anaesthetic agents, and in all cases except where the basal metabolic rate is high. It inhibits the action of the vagus and diminishes mucous secretion. It should be given hypodermically in doses of 1/100 to 1/150 grain one hour before operation, except in the case of children, when it is better to give tincture of belladonna by the mouth to obviate the discomfort of the subcutaneous injection.

Summary

To sum up, for children and nervous patients some sedative is essential, but no one drug is suitable in all cases. Administration should never be done by the intravenous route, the rectal route being the most suitable and avertin the best of the drugs so used.

Atropine should be given except when the basal metabolic rate is high.

SINUSITIS IN CHILDREN*

BY

G. C. SCANTLEBURY, M.B., B.S.MELB., F.R.C.S.
SURGEON TO THE EAR AND THROAT DEPARTMENT, MELBOURNE
HOSPITAL

The importance of sinusitis in children has been realized only in comparatively recent years. Though full sinus development may not be completed until about the fifteenth year, the ethmoidal cells, and to a lesser extent the maxillary antrum, are definite structures at birth, and by the fifth or sixth year all sinuses are large enough to be the seat of acute or chronic disease. Acute disease may have serious (or fatal) local complications, giving rise to retrobulbar neuritis, orbital cellulitis or abscess, cavernous sinus thrombosis, brain abscess, meningitis, or otitis media with all its complications. These, however, are beyond the scope of a short paper, and it is with chronic and subacute sinusitis that I wish to deal.

The frequency of chronic sinusitis has been stressed by so many writers, and is such a commonplace with all rhinologists, that details need not be repeated. The picture of a child with this disease is typical in many cases. A pale-faced, easily fatigued child with a running nose, nasal obstruction, a history of frequent colds, a loose cough, worse during the night and in the morning, anorexia, deficient general development, intermittent or constant hoarseness, with sometimes headaches (Valentina Wasson¹ states that this occurs in 20 per cent. of cases in older children), and a history of attacks of otitis media and cervical adenitis, is common. Night sweats and persistent temperature are frequently observed.

Causative Factors

The causes of sinusitis are deficient hygiene, defective feeding, bad climatic conditions, allergic state, exposure to local infections, nasal abnormalities, and hypertrophied tonsils and adenoids. These causes are variously emphasized by different observers. All are agreed that deficient hygiene—such as bad ventilation, damp houses or clothing—are definite factors in the causation of the disease.

Defective feeding has received a lot of attention. R. G. Turner² considers sinusitis to be a vitamin A deficiency disease, and has demonstrated great improvement in cases treated with carotene. Excess of starch and meat is likely to produce an acidotic tendency, and is deprecated by Luongo,³ who prefers a diet of lacto-vegetarian food,

which should be as much as possible fresh and unprocessed. He also considers that lack of calcium is a factor. Rea Ashley⁴ considers hypothyroidism important in some cases, and has had good results by exhibiting small doses of thyroid extract or by using calcium gluconate and parathyroid.

Climatic conditions undoubtedly play a part. In Australia sinusitis is much more prevalent on the coast than in inland hilly country. Dr. E. M. Kent-Hughes⁵ states that in Armadale, New South Wales, the condition is uncommon, and that children from the coast are very quickly cured by residence in that district.

Exposure to local infections is seen in the very frequent occurrence of sinusitis following constant colds where children at school infect one another, and one is constantly struck by the presence of sinusitis in different members of one family. Influenza and infectious diseases such as scarlet fever, measles, and pertussis are common causes. Hoopple and Cave⁶ found sinusitis in all except one of a large series of cases of otitis media following scarlet fever. Nasal deformities, such as deviated septum, by interference with the normal drainage and ventilation of the nose, are considerable factors.

TONSILS AND ADENOIDS

Hypertrophy of tonsils and adenoids is placed last as a cause, not because it is necessarily the least important, but because it is the one about which there is probably some difference of opinion. In most textbooks and articles on the subject it is frequently given as the most important cause, but I have long been of opinion that the possibility of its being an effect should be kept in mind. Lymphoid tissue is firmly believed to have a protective effect in that it deals with infection, being, to a certain extent, "the first line of defence." If it becomes grossly enlarged and chronically infected we can consider that it is losing in the fight, and has become a factor on the other side by causing obstruction and by acting as a sponge to hold infection. I agree with Cantor,⁷ who does not advise removing tonsils and adenoids until he is satisfied that they are obstructing or "septic." The mere presence of tonsils and adenoids is not enough.

Allergic causes must always be kept in mind. Skin tests should be done by skilled allergists when practicable, and appropriate measures carried out.

Types Found in Children

In general, three types of chronic sinusitis are found in children: (1) suppurative; (2) hyperplastic; and (3) mixed suppurative and hyperplastic. There is probably never any infection of the sinuses without hyperplasia, and on the degree of hyperplasia depends, to a very large extent,

* Read in the Section of Laryngology at the Annual Meeting of the British Medical Association, Melbourne, 1935.