BILIARY PERITONITIS

BY

STEPHEN POWER, M.S., F.R.C.S.

SURGEON, ELTHAM HOSPITAL

A recent case of biliary peritonitis recalled to my mind two others, and suggested that the subject is one which is not commonly discussed in the textbooks and is almost ignored in current literature. My three cases are recorded in brief below.

Case Records

Mrs. A., aged 65, was admitted with a history of abdominal pain and uniform tenderness and distension, which made one suspect an obstruction. A slight icterus negated this diagnosis, and as her condition rapidly deteriorated she was not subjected to operation. She had had a cholecystectomy a year previously. At post-mortem examination the abdomen was found to be full of bile, and there were several stones in the common bile duct, though no perforation was discovered; it is not unlikely that it may have eluded the pathologist.

Mrs. C. was admitted with symptoms and signs indicating acute spreading cholecystitis. Laparotomy confirmed the diagnosis, but also revealed massive adhesions of old-standing disease and a leakage of bile. No perforation was seen, but circumstances did not favour investigation. Multiple calculi were removed and cholecystostomy performed. The gall-bladder was removed at a later date.

Mr. S., aged 70, appeared clinically to be a mild case of perforation. Actually he had a tense gall-bladder with extravasation. A point of leakage was not seen. Several small stones were present. Cholecystostomy was done. The fistula persisted, and one anticipated the necessity of further interference, but unfortunately a coronary thrombosis carried the patient off.

I have seen one other case at operation in which the bile was oozing from a pin-point perforation in the fundus. The haemorrhagic nature of the exudate was quite striking.

Aetiology

The condition of biliary peritonitis is not common, and occurs only about once in every hundred gall-bladder cases. Extravasation may be from the gall-bladder itself or from any of the ducts, as a result either of disease or of trauma. It must be remembered that the duodenal contents can be very similar in appearance. The extent of the lesion or lesions, for they can be multiple, varies from the microscopic to widespread sloughing cholecystitis, and is not necessarily located on the peritoneal aspect of the bile passages, for an instance of a large retro-peritoneal escape has been recorded. Localization is only likely in the event of long-standing inflammatory trouble, or after rupture due to injury. The exact mechanism of the process is uncertain, and probably varies, but the presence of stones is an important factor, as they are found in over 70 per cent. of cases.

The actual hole may be the result of pressure necrosis at the site of impaction, and this applies even when local tension is not marked. Small stones are found not infrequently lying in the depressions between the trabeculae of a "chronic" gall-bladder, producing minute ulcers. In a later stage a large calculus is often seen to be in contact with an area of much grosser ulceration. In the cases where stones are not present an infective ulceration can lead to destruction of all coats, and this is surprisingly common in typhoid fever. A mechanical bursting is possible with a cystic duct blockage, either by stone or by oedema. The gall-bladder in health is generally reputed to empty easily. Practical experiment or a consideration of the usual calibre of its duct makes it obvious that some little patience must always be expended on the emptying of the viscus, and that no great degree of inflammation is required to block it. However, if this

channel is effectively occluded it is unlikely that any big extravasation can ensue. The Germans postulate a condition of over-viscid bile to explain the syndrome of very definite colic without the presence of calculi. Whether this could ever raise intrabiliary tension sufficiently to cause leakage is extremely doubtful. Of considerable practical importance is the site of the perforation, and, fortunately for the surgeon, this is, in 90 per cent. of cases, in the gall-bladder itself, usually in the fundus, but sometimes in the pouch of Hartmann and sometimes midway between. In 3.3 per cent. it is in the cystic duct, in 4.4 per cent. in the common bile duct, and in only 1.1 per cent. in the hepatic ducts.

A very striking feature of many of these cases is that the gall-bladder is healthy, and no gross ulceration can be found. Rouillard and Schwob had a case of a woman of 83, not operated on, who on post-mortem examination was found to have a big extravasation with no signs of inflammation in the gall-bladder and no perforation, although a few stones were present. The organ was not under tension, but the cystic duct was narrow. To explain this case they quote Leriche's almost lyrical description of his findings in a similar one: "*La vésicule distendue laisse suinter une véritable rosée qui tombe dans l'abdomen et se renouvelle quand on l'essuie avec une compresse.*" The gall-bladder here was tense. Their observation was made after death. Yasse and Cornelis describe a similar happening. In this case a lax and healthy gall-bladder was seen to have on its lower surface a dark stain about one inch long, from which a drop of bile was suspended. There were no stones, and subsequent examination after cholecystectomy revealed no thinning or solution of continuity in the discoloured area. These instances leave no doubt that transudation of bile is not a pathological myth. The Continental literature provides numerous other examples.

One author describes a case which, in addition to extravasation, showed signs of acute pancreatitis. He raises the question of the relation between the two conditions. It is an undoubted fact that fat necrosis is an occasional concomitant of bile leakage, and this independently of pancreatic disease. The explanation seems to be that the bile salts increase the permeability of the pancreas and allow its enzymes to escape and act on fat, and even on the liver. Microscopy reveals no disease of the gland when this happens. Rewbridge conducted experiments on these lines, and found that bile or the salts thereof injected into an animal's peritoneal cavity produced fat necrosis, especially in the neighbourhood of the pancreas, but failed to do so if that gland had been previously removed. It has further been demonstrated that bile used in this way, even if sterile, will kill an animal, but experience teaches that it is relatively innocuous in man. The fatal type of peritonitis produced by morbid process is bacterial rather than chemical.

Symptomatology, Treatment, and Prognosis

Biliary extravasation is a disorder of later life, the average age of the patient being 50. Women are the more frequent subjects, and second attacks are not unknown. The signs and symptoms are such as might be expected. The patient usually commences with epigastric pain of sudden onset, but it may be preceded by the pain of colic or cholecystitis. The state of affairs resembles that found in rupture of a peptic ulcer, but is not so acute. Tenderness and muscular tension are widespread, especially on the right side of the abdomen, but there is usually absence of board-like rigidity, and there is a tendency to distension. Temperature and pulse are both raised, and there may be shifting dullness. The quantity of fluid is often enormous, as the gall-bladder has lost

its concentrating function, but, on the other hand, supply may be obstructed at its duct. A well-localized case can simulate a pure cholecystitis or even colic, but any spread of the physical signs should bring these diagnoses under suspicion. A little later in the history an appendicitis may be considered, through the tracking of exudate into the right iliac fossa. Still later it is necessary to distinguish from general peritonitis and intestinal obstruction. Jaundice is not as common as might be expected, and is late in onset and mild in character.

Operation is the only treatment for this condition, and it cannot be advised too early. Some traumatic cases are temporarily relieved by tapping, but the measure is not curative. Suture of the perforation is, in view of the nature of the disease, obviously hopeless, and cholecystectomy is the most satisfactory course to adopt. Local and suprapubic drainage is employed. If any of the ducts has been perforated the procedure can be suitably modified and a drain placed in its lumen. Peritoneal toilet is the only other requisite, and attention should be directed to the state of the lesser sac. On some occasions, with very ill patients or with considerably damaged gall-bladders, it may be impossible to remove the organ, and it then has to be left for a later date, the surgeon being content with cholecystostomy and removal of stones.

The prognosis depends on the degree of infectivity of the bile and the promptness of intervention. It would appear that bile exalts the virulence of the infection and diminishes the bactericidal properties of the exudate which it produces. In the traumatic group, with sterile bile, it seems possible to delay operation for as long as six weeks or more without a fatal result. The recovery rate is about 80 per cent., whereas in the pathological perforation group all die without operation, and only 50 per cent. recover with it. Even here, however, life is sustained sometimes for as long as twelve days without laparotomy, and some of the patients are still fit enough to go on the table. This is in marked contrast to the rapid deterioration after stomach or duodenal contents are liberated in the peritoneal cavity.

BIBLIOGRAPHY

- McWilliams: *Ann. of Surg.*, 1912, lv, 235.
 Fifield: *British Medical Journal*, 1926, ii, 635.
 Mentzer: *Arch. of Surg.*, 1934, xxix, 227.
 Rewbridge: *Arch. of Path.*, 1931, xii, 70.
 Rouillard and Schwob: *Bull. et Mém. Soc. Méd. des Hôp. de Paris*, 1930, xlvi, 1891.
 Yasse and Cornelis: *Le Scalpel*, April, 1935.

During the twelve months ended March 31st, 1935, the National Institute for the Deaf obtained new headquarters in Gower Street at a cost of over £12,000, of which about £7,500 remains to be found. A fine hall has been erected at the rear of the property to the memory of the founder of the Institute, Mr. Leo Bonn. In the annual report for 1934-5 the claim is renewed for the appointment by the Government of an advisory committee on the deaf and the passing of a Deaf Persons Act in view of the inadequate benefit derived from the Ministry of Health's Circular No. 1337. Six regional associations for the deaf cover the whole of England and Scotland. All are affiliated to the Institute, and have adopted a scheme ensuring closer mutual co-operation. These associations are of the greatest importance to the future development of work for the welfare of the deaf, since they bring together the public authorities upon whom that work must to a large extent ultimately depend and the voluntary societies actually engaged in it. The report includes accounts of the many and varied activities carried on by the National Institute, and the protest against the "ramp in aids to hearing" is repeated. The organization will attain its silver jubilee next June.

Clinical Memoranda

Melanoma of the Choroid with Secondary Deposits in the Liver, etc.

In the *British Medical Journal* of October 5th, 1935 (p. 624), Dr. Halls Dally describes a case of melanoma of the choroid with late secondary deposits in the liver. Full details of laboratory and post-mortem examinations are given. No mention, however, is made of the urine having been examined for the presence of melanin or its immediate precursors, an examination which was of value in the case recorded below, as well as in others in the experience of the writer.

CASE REPORT

On September 11th, 1935, I was asked by her medical attendant to see in consultation with him a 73-year-old lady in a town on the west coast of Cumberland. For seven or eight years she had had "bilious attacks," with pain in the right upper quadrant of the abdomen and occasional evanescent jaundice. These were attributed to biliary colic, and traces of bile were found in the urine in April, 1932, and in January, 1934. In June, 1935, there was tenderness in the right upper abdomen, and in August marked deterioration was noted in her general condition. On September 10th enlargement of the liver was definitely present, and there were signs of a right pleural effusion.

On June 24th, 1932, she had developed an acute glaucoma of the right eye, was seen by an ophthalmologist without delay, and had the eye removed. The pathological report on the eye was: "Melanoma of the choroid involving the ciliary body and extending through into the anterior chamber." There had been no suspicion of any local recurrence.

Seen by me on September 11th, she was of a good colour, with no jaundice, but showed signs of recent weight loss, and had atonic muscles. The right eye was artificial, the orbit apparently healthy. The tongue was coated, the teeth false, the throat healthy, and no enlarged glands were felt in the neck, axillae, or groins. The chest had an old bony deformity, and respiratory movement was restricted on the right side, in which was an effusion which extended up to the sixth rib behind. The heart was not appreciably displaced or diseased; the blood pressure was 156/84 mm. Hg. A bulge was present in the right upper abdomen. The abdominal wall moved poorly with respiration. In the mid-clavicular line the liver was felt to extend to a distance of five inches below the right costal margin. It was hard, irregularly nodular, and tender. The left lobe was similarly involved. The spleen was not enlarged. A finger in the rectum detected no disease of the bowel nor any secondary deposits of malignant disease in the pouch of Douglas.

The urine was not abnormally dark in colour, nor did its colour appreciably change on standing, but it contained a trace of albumin. The presence of melanin compounds was detected by the ferric chloride test, the ferric-chloride-in-hydrochloric-acid test, and by the Thormählen reaction.

The patient died on September 27th, 1935.

DISCUSSION

The rapidly increasing hepatic enlargement of irregular nodular type suggested secondary malignant disease, as did the pleural effusion. A primary focus had been removed three years before, and, although there had been no local recurrence, the finding of melanin bodies in the urine placed the diagnosis beyond reasonable doubt. There was thus no need to worry the patient by any more complicated investigations.

SUMMARY

There is described a case of melanoma of the choroid, removed by operation, and followed three years later by metastases in the liver and chest without demonstrable local recurrence.