ANTE-NATAL DIAGNOSIS OF QUADRUPLETS

BY

E. ULYSSES WILLIAMS, M.R.C.S.
SENIOR RADIOLOGIST, ROYAL FREE HOSPITAL

(With Special Plate)

During the last ten or twelve years the radiological diagnosis of twins or triplets, at any time after the sixth month of pregnancy, has come to be looked on as part of the ordinary routine work sent up from the maternity departments. It is thought, however, that the diagnosis of quadruplets two months before birth is still uncommon enough to merit a brief article.

A fairly extensive search of the literature on the subject has revealed only one other instance of the x -ray diagnosis of quadruplets before birth up to the present time. This was in Manchester in 1934, and in this case the skiagraphic picture, though establishing the fact of four foetuses being present, was not, so far as I can ascertain, reproduced in any medical journal.

Radiological Aspects

The following few notes record shortly the radiological aspects of my case.

On August 21st, 1935, Mrs. H., a multipara aged 34, was sent to the x -ray department of the Royal Free Hospital with a clinical diagnosis of "2 twins." The skiagram showed at once that quadruplets were present. One, hereafter called Foetus A, was a vertex with the head very low in the pelvis. Foetus B was a breech with flexed legs, lying well over to the left of the maternal abdomen. Foetuses C and D were lying head to tail transversely on the right side of the abdomen, both well above the level of the right iliac crest. The shadow of the head of the upper one, Foetus C, reached to the level of the eleventh rib.

All four foetuses were apparently well formed, A and D being equal in size, and smaller than B and C, which were again of equal size.

Clinical data as to the duration of pregnancy were conflicting, but from a skiagraphic standpoint it appeared that the general ossification was that of foetuses of about thirty weeks, but their size, as was to be expected, was only that of twenty-six weeks.

The suboccipito-bregmatic measurements at this time were: A, 2.8 inches; B, 2.7 inches; C, 2.75 inches; and D, 2.85 inches.

There did not seem to be any excess of liquor amnii.

On September 18th the patient was x -rayed again, and the quadruplets were seen to have grown considerably and to have altered their position.

Foetus A was still a vertex, with head well down in the pelvis, but B, C, and D were now all breeches, lying side by side in the upper part of the uterus, with their heads more or less on the same level. C was in the middle, with B and D having their backs to it, and the upper parts of the body curved symmetrically away. The general appearance, as can be seen from the illustration on the Plate, is strangely like three tulips in a vase.

The general ossification of the skeleton had progressed considerably, all the foetuses had grown well, and the heads now measured: A, 3.5 inches; B, 3.3 inches; C, 3.4 inches; and D, 3.5 inches.

It will be seen at once that the ratio of head diameters has been kept, but the actual increase of the size of the head shadows is 0.75 inch in each case. The usual increase in size of the foetal head in the last months of pregnancy is $2/5$ inch per month. The discrepancy of $3/10$ inch is accounted for by the fact of the great increase in diameter of the maternal abdomen. This causes the heads to be unavoidably further from the photographic film and therefore projected larger.

In conclusion, it was estimated on general skiagraphic appearances that the duration of pregnancy at the date

of the photograph reproduced (September 18th) was thirty-four weeks. The further rather hazardous opinion was made, and committed to paper, that the foetuses were all boys. On October 13th quadruplets were delivered, all boys, as recorded by Dr. Constantine in her clinical account of this case.

A CASE OF QUADRUPLETS

BY

M. C. E. CONSTANTINE, M.D.

(From the Gynaecological and Obstetrical Unit of the Royal Free Hospital)

As a quadruplet pregnancy is a comparatively rare event it was felt that a short account of the recent case at the Royal Free Hospital might be of some interest. Diddle and Burford,¹ collecting figures from the literature, recently gave the incidence of quadruplet labour as 1 in 654,455, from a total of 219,899,446 recorded births.

Case Record

History.—The patient, aged 34, had had three previous normal pregnancies and labours. The children are girls, 6, 4, and 2 years old respectively. There was no history of multiple pregnancy in the father's family for three generations. The wife, one of eleven children, was herself a twin, her twin brother having died in infancy. Her mother, one of three children, had had twin sisters, who died as infants. One of the patient's brothers had twin sons living, in addition to five other children. This family history is in keeping with the general opinion that the mother has more influence in producing multiple pregnancies, though cases such as those recorded by Tarnier,² Gartier,² Strassman,² and Peifer³ suggest a paternal influence also. There is some doubt when this pregnancy began. The last normal menstrual period lasted from about December 27th, 1934, to January 1st, 1935, but for seven days at the end of February there was an excessive loss, which the patient thought to be a miscarriage, as clots were passed. After this no further haemorrhage occurred. The small size of the infants, their general appearance, measurements, and muscular inertia suggest several weeks' prematurity, but radiosopic examination of the first two infants after death showed a degree of ossification approaching maturity. Mr. Ulysses Williams reports on these findings in the preceding article.

Examination.—The patient first attended the district ante-natal clinic on August 20th, when the abnormal size of the uterus suggested a multiple pregnancy. X -ray examination to confirm this showed four foetuses. The only other abnormality noted at this time was moderate oedema of feet and legs, the urine and blood pressure being normal. The oedema and general discomfort increased, and on September 5th the patient was admitted for rest and observation. There was no evidence of toxæmia, and, apart from pressure symptoms, the pregnancy continued normally until labour began on October 10th. Her girth on this date was $45\frac{1}{2}$ inches, and a head was palpable just above the pelvic brim.

Labour.—At 2.30 p.m. on October 10th there was a sudden discharge of blood-stained liquor, but pains did not begin until four hours later. The contractions were very irregular and slight for the next twenty-four hours, and at the end of this time the os was found to be two fingers dilated, with the first head in the pelvic cavity. Soon after this there was a further discharge of clear liquor, probably from rupture of a second bag of membranes. It was only about midday on October 12th that the contractions became regular and forceful. At 6 p.m. a second vaginal examination was made, and the cervix now found to be dilated except for a small posterior rim and an anterior lip, the head still in the oblique diameter. In spite of strong and frequent pains no further advance was made, and four hours later it was decided to deliver the first child while the uterus was still acting well.

Delivery.—After rotating the head an extremely easy forceps delivery was effected at 10.25 p.m. Contractions recommenced

J. MINDLINE: MASSIVE COLLAPSE OF LUNG COMPLICATING HAEMOPTYSIS

FIG. 1.—Commencing collapse at left base. No obvious lesion at left apex. (October 1st, 1934.)

FIG. 2.—Complete collapse of left lung. Note raising of left leaf of diaphragm, displacement of heart and trachea to left, and translucency of right lung, which can be seen extending as a relatively clear area to left of vertebral column. (October 4th, 1934.)

FIG. 3.—Re-expansion at left base almost complete. Left apex and mid-zone shows area of tuberculous infiltration. Heart and trachea have returned to normal position. (October 18th, 1934.)

E. ULYSSES WILLIAMS: ANTE-NATAL
DIAGNOSIS OF QUADRUPLTS

X-ray photograph, taken on September 18th, 1935, antero-posterior view.