Summary

Five cases of atypical megalocytic anaemia are described. In the first, in a boy of 11, the stomach showed normal acidity and the condition reacted well to liver therapy.

The second case demonstrated the development of a megalocytic from a microcytic anaemia in a woman with normal gastric acidity. The condition failed to react to liver or iron therapy until after blood transfusion. The condition was associated with tetany.

Cases III and IV appeared typical of pernicious anaemia; in Case III the anaemia was associated with soreness of the tongue, but analysis of the gastric contents in both cases showed the presence of free hydrochloric acid.

Case V cannot be definitely classified as of achrestic anaemia, owing to the fact that on admission to the hospital the patient was too ill to have a test meal. An anaemia at first reacting to iron but later becoming definitely megalocytic, it showed no response to iron or liver therapy, and only reacted slowly to blood transfusion. The anaemia was accompanied by menorrhagia, and during the more severe stages with mania. The patient died of bronchopneumonia, and the post-mortem findings were those of intense normoblastic activity in conjunction with fatty degeneration of the vital organs.

REFERENCE

¹ Wilkinson, J. F., and Israëls, M. C. G.: *British Medical Journal*, 1935, i, 139 and 194.

A CONTRIBUTION TO THE TREATMENT OF BURNS

BY

ALFRED C. TURNER, M.B., B.Ch.Belf.,
F.R.C.S.Ed.

SURGEON, THE ANGLO-IRANIAN OIL COMPANY

Ten years ago E. C. Davidson of Detroit introduced a new method of treatment of burns by means of tannic acid solution. The surgeon has thus been armed with a weapon completely replacing modes of treatment which were clumsy, unsatisfactory, and expensive, while mortality has been very considerably lowered and suffering greatly reduced. There will be no doubt in the minds of those experienced in the treatment of extensive burns of the first three degrees that the coagulating method is the ideal one. Certain aspects of the tannic acid technique are, however, unsatisfactory. An attempt has been made to overcome the worst of these by modifying the treatment while still retaining the basic ideal—the "coagulated" and desiccated surface.

The advantages and disadvantages of any particular line of treatment can only be ascertained by careful examination and analysis of a large series of cases. The observations which follow were made over a period of five years, during which the tannic acid method was used, and a one-year period in which modifications of this technique were adopted. The total number of cases of burns in this series, inclusive of all degrees sufficiently serious to render the patient unfit for work, was 2,696.

Disadvantages of Tannic Acid

1. *The Coagulum*.—In the case of tannic acid this is coarse, tough, and not transparent. Any pus formation beneath it is difficult to diagnose, and may spread over a large area rather than perforate an unyielding scab. In this way much of the young epithelium may be destroyed before any signs of damage are manifest. When the scab, floating on a bed of pus, is removed later a very coarse type of granulation tissue will be found beneath, which is difficult to control and ultimately tends to produce extensive scarring. These observations suggested to us that certain definite advantages could be claimed for a thin, transparent scab.

2. *Destructive Action*.—Tannic acid may well be described as "dirty," and is, in addition, most destructive to clothing and bed linen. This necessitates the use of a mackintosh to protect the bed. The mackintosh acts on the burnt area like jaconet used in a fomentation. Delay in evaporation takes place to such an extent as seriously to interfere with one of the main essentials of the treatment. The choice, therefore, lies between sacri-

ficing the bed linen or restricting the evaporation from the surface of the coagulum, which is so essential for its well-being.

3. *Instability in Solution*.—A further very important objection to tannic acid is that it rapidly becomes unstable in the aqueous solution required for use. This makes its employment most cumbersome both as a first-aid measure and also in hospital practice.

The above disadvantages suggested a search for an equally satisfactory coagulant which would not destroy linen and could be kept indefinitely in solution.

Process of Repair of a Burn

In the process of repair the exudate from the surface does not coagulate, or does so only in a very slight degree. Fibrin strands may be demonstrated, but are usually insufficient in quantity to form a firm coagulum. Many drugs are found to have the property of assisting in the formation of a coagulum. Our contention is that the ideal drug to employ is one with strong antiseptic and bactericidal properties, which will at the same time help to bind the crust. After a trial with several drugs, mercurochrome (di-sodium-dibromo-oxymercury fluorescein) was selected, as it appeared to possess the necessary qualities. It was given an extensive trial over a period of one year, and the following observations were made.

1. Mercurochrome does not precipitate protein, and can therefore be claimed as an effective antiseptic in the presence of protein.

2. In cases treated with this drug the crust formed is thin and transparent.

3. Bed linen is not in any way destroyed.

4. It is non-irritating to the tissues.

5. A 2 per cent. aqueous solution is stable for an indefinite period.

6. Epithelialization under the scab takes place rapidly. It was noteworthy in using some other drugs, particularly trypanflavine, that though the crust formed was excellent little or no epithelialization took place beneath it, and aseptic necrosis of the superficial living tissues occurred.

Technique and Results of Mercurochrome Treatment

The treatment of shock and exhaustion are dealt with on generally accepted lines and will not be further discussed here. With reference to the preliminary local treatment there are certain points of importance.

1. No general anaesthetic—not even nitrous oxide and oxygen—is given. Many cases of extensive burns require hours of preliminary work, which can only be accomplished by dealing with small areas at a time. Large doses of morphine or other opiates are recommended, and are usually quite effective.

2. All dead tissue is stripped from the burnt area and foreign material removed. The greater the care taken in this the better will be the end-result.

3. At this stage no antiseptic whatever is recommended. The denuded area should be swabbed over with cotton-wool soaked in normal saline at about 100° F. until all debris has been removed.

4. When the general toilet of the area has been completed it is swabbed over with a 2 per cent. aqueous solution of mercurochrome. The surface is then dried with the assistance of an electric hair drier, an ordinary electric radiator, or an electric cradle, provided the last-named has ventilation holes along the top. Desiccation takes place rapidly.

The following routine treatment is recommended: first day, four applications; second day, three applications; third day, two applications. The area is dried off after each application, and is always kept exposed to the atmosphere. In some cases where a large area is involved the patient is forced to lie on a portion of the burnt surface and the crust is likely to disintegrate. This feature is not associated in any way with the drug used, but in our opinion is the result of inadequate evaporation from the surface. The patient should therefore be frequently turned and the freest ventilation possible provided. Should there be any local collection of pus—and diagnosis of this is facilitated by the thinness and semi-transparency of the crust—the crust should be freely removed from the area, the area swabbed up with normal saline as before, and the mercurochrome treatment reapplied.

A very brief account of a recently treated case is given below.

F. B., aged 32, European, an electrical engineer, was admitted to hospital on the morning of August 19th, 1935, suffering from burns sustained by coming in contact with an electric arc at a voltage of 33,000. It is assumed that all the burns were caused by the actual flame of the arc.

The following were the areas and degrees of the burns sustained. *First degree.*—A large area of the chest above the nipples and the back of the neck. *Second degree.*—The entire face and front of the neck as far back on each side as the mastoid process; the left forearm; an area about three and a half inches in diameter over the left patella. *Third degree.*—The back of the left hand; the right thenar eminence; an area on the chest from left clavicle to the fourth interspace. There was also superficial burn of the lower half of both corneae.

The eyes were treated with cocaine, atropine, and liquid paraffin, and cleared up in thirty-six hours. The burnt areas were treated by the routine mercurochrome method. The patient's temperature was normal from the fourth day onwards. By the twelfth day the burns had cleared up, though the scab had not yet separated from the back of the left hand. There was no scarring, and the patient was able to shave on the eleventh day.

Conclusion

Our results have been most encouraging. The rapidity and reliability with which desiccation has taken place have been eminently satisfactory, and with adequate preliminary treatment we have been able to reduce sepsis to a minimum. On a general survey of our cases it would appear that patients treated with mercurochrome have less general reaction during convalescence than those treated with other agents, while the amount of pain and discomfort compare well with any other method. On no occasion have we seen any toxic effects following the use of this drug.

I wish to express my thanks to Dr. E. Jamieson, chief medical officer of the Anglo-Iranian Oil Company, for permission to publish this article.

BIBLIOGRAPHY

- Davidson, E. C.: *Surg., Gynecol. and Obstet.*, xli, No. 2.
 Wilkie, D. P. D., et al.: Report on Medical Treatment of Men Burned in Colliery Explosions. Mines Department, H.M. Government.
 Mitchiner, P. H.: *Med. Press and Circ.*, December 13th, 1933.

A NOTE ON CUTANEOUS CANCER IN COTTON MULE-SPINNERS

BY

E. D. IRVINE, M.D., D.P.H.

ASSISTANT MEDICAL OFFICER OF HEALTH, BLACKBURN

Dr. A. E. Somerford, in his interesting paper published in the *British Medical Journal* of June 29th, 1935 (p. 1305), concludes that non-scrotal cancer in cotton mule-spinners is not occupational in origin, and supports the view that the causal factor in the development of scrotal cancer in these workers is not exposure to mineral oil, but continued friction at this site.

A Local Inquiry

In 1930 I made an inquiry into the occupational incidence of cutaneous cancer in men in the Blackburn Registration District,¹ the census figures for which give the number at risk in the different occupations, this, of course, being of the greatest importance in the consideration of occupational diseases.

Details of patients suffering from this disease and who lived in this area were extracted from the death registers from 1837 and from the in-patient and out-patient registers of the hospitals serving the area (from 1865). The cases not ascertained by this means must be few in number. The cotton trade is the principal industry, employing almost all of the textile workers in the district, and many of the other industries are associated with, and to a great extent dependent on, it.

In all 625 cases were traced, 246 occurring in men engaged in the cotton trade. In addition, there were twenty-five* of epithelioma or cancer of the groin and groin glands, almost certainly secondary to cancer of a skin site (scrotum, penis, or leg), and a further twenty-five† of "epithelioma," the site of which was not stated. As further information about these cases was not available they have not been included in the following table, which summarizes the findings and shows the numbers employed in the different occupations as stated in the 1921 census for the district.

Occupation as described in Registers	Number of Men Employed (Census 1921)	Cases of Skin Cancer Traced									
		Rodent Ulcer	Squamous Epithelioma					Total Non-scrotal	Scrotal Epithelioma	Total	Percentage Non-scrotal
			Head and Neck	Lip	Penis	Others Non-scrotal					
Cotton mule-spinners	1,502	19	9	—	3	8	39	94	133	29.3	
Cotton weavers	10,917	13	13	7	7	8	48	3	51	94.1	
Other textile workers	10,050	18	21	4	6	7	56	6	62	90.3	
Others	52,621	64	106	85	44	44	343	36	379	90.5	
All occupied males	75,100	114	149	96	60	67	486	139	625	77.7	

The following table is intended to show the relative incidence of cutaneous cancer among spinners and other sections of the employed male population.

Occupation	Number of Cases traced per 1,000 Employed in 1921	
	Non-scrotal	Scrotal
Cotton mule-spinners ...	26.0	62.6
Cotton weavers ...	4.4	0.3
Other textile workers ...	5.6	0.6
Others ...	6.5	0.7
All occupied males ...	6.5	1.8

* Spinners 8, weaver 1, other textile workers 5, others 11.

† Spinner 1, weaver 1, other textile worker 1, others 22.