TREATMENT IN GENERAL PRACTICE

This article is one of a series on the management of some diseases of the nervous system met with in general practice.

DISSEMINATED SCLEROSIS

BY

W. RUSSELL BRAIN, D.M., F.R.C.P.

Disseminated sclerosis is probably due to an infection of the nervous system. Inflammatory patches in which the myelin sheaths of the nerve fibres are destroyed occur mainly in the white matter of the brain and spinal cord, including the optic nerves. These patches appear at intervals, a fresh patch leading to new symptoms or to an exacerbation of existing symptoms, while during the earlier phases of healing and scarring the patient improves, often strikingly. The appearance of patches may be so rapid that the patient dies within three months of the first symptom, or the intervals between the outbreaks may be so long that there is no substantial disablement at the end of twenty-five years. The nature of the symptoms and signs depends upon the situation of the patches in the nervous system. A common early symptom is acute unilateral retrobulbar neuritis, which leads to primary optic atrophy characterized by pallor of the optic disk, especially in its temporal half. The brunt of the damage usually falls upon the pyramidal tracts and the cerebellar connexions. The principal manifestations of the disease therefore are weakness of the upper motor neurone type and incoordination. These find expression in nystagmus, slurred speech, intention tremor, a spastic and ataxic gait, increased tendon reflexes, and bilateral extensor plantar responses. The abdominal reflexes are usually lost at an early stage. Impaired control over the bladder leading to precipitancy or delay in micturition is a common early symptom. Since the cause of the disease is unknown treatment is necessarily empirical, and it is clearly very difficult to judge the value of any method of treatment for a disorder which runs such a variable course.

General Management

Rest is essential during an exacerbation, and if this is at all severe the patient should be kept in bed until there are signs of improvement, which usually takes several weeks. In the intervals of remission the patient should be encouraged to live a normal life as far as his disability allows, always, however, avoiding carrying exertion to the point of fatigue. One should aim at postponing as long as possible the ultimately inevitable bed-ridden state, and suggestion and encouragement will be a great help in this. Anything which may be impairing the health should receive attention, and if there are septic foci in the nose, throat, or teeth these should be treated, but with reasonable conservatism, since the wholesale extraction of infected teeth in one operation, for example, may lead to an exacerbation of the disease.

Treatment with Drugs

Arsenic in both its organic and its inorganic forms has long been part of the routine treatment. Its value is a matter of personal impression. It may be prescribed in the form of 3 minims of liquor arsenicalis in half an ounce of water three times a day after meals. This

mixture should be given for from six to twelve months at a time, omitting it, however, for one week in every four. Larger doses are sometimes administered, but do not seem to be of greater benefit. Organic compounds of arsenic are given on the same lines as for the treatment of neurosyphilis, though usually in smaller doses and shorter courses. Novarsenobillon, silver salvarsan, and sulfarsenol all have their advocates. Novarsenobillon should be injected intravenously in doses of 0.45 gram for a male and 0.3 gram for a female patient. Silver salvarsan is prescribed in doses of 0.2 gram for a male and 0.15 gram for a female patient. Whichever drug is used, an intravenous injection is given weekly for four to six weeks, and this course is repeated at intervals of about six months. Sulfarsenol is administered in doses of 0.45 gram to a male and 0.3 gram to a female patient, either intravenously or into the deep subcutaneous tissues, in courses at the same intervals. During an exacerbation iodide may usefully be prescribed for a few weeks, either alone in the following mixture:

R.	Pot. iod.	...	...	...	...	gr. v-x
	Aq. menth. pip.	...	...	...	...	ad. 3 ss
					t.d.s.	

or with the addition of a drachm of liq. hydrarg. perchlor. to each dose.

Good results have recently been claimed for the prolonged administration of quinine hydrochloride in doses of 5 grains three times a day by the mouth. Liver extract has also been advocated, the equivalent of half a pound of liver being taken daily for an indefinite period.

Treatment By Artificial Pyrexia

Artificial pyrexia sometimes leads to striking temporary improvement, and is most likely to be effective in rapidly progressive cases, especially in young adults. Various methods of inducing pyrexia have been used, including malaria, the subcutaneous and intravenous injection of foreign protein, and physical agencies. A convenient and safe method is the intravenous injection of "pyrifer," a *B. coli* vaccine. This is supplied in graduated doses, and the patient receives a dose intravenously on alternate days three times a week, six to ten injections being given. The rise of temperature extends over several hours, and usually reaches about 103° F. The patient must be kept in bed during the course of injections.

Treatment of Special Symptoms

During an attack of acute retrobulbar neuritis the affected eye should be covered with a shade or a dark glass, and the eye should be rested as much as possible. Ataxy of gait may be diminished by suitable re-educational exercises, such as Fraenkel's exercises. When, owing to spasticity of the calf muscles, there is marked plantar flexion of the foot a surgical boot with a toe-raising spring may make walking much easier. Frequency and precipitancy of micturition may be relieved by small doses of belladonna, such as half a grain of dry extract in a pill two or three times a day. During the late stages of the disease the skin, the bladder, and the rectum will require the attention necessitated by severe degrees of paraplegia (see article by Dr. Henry Cohen, October 5th, p. 633).